



Vital and Health Statistics

From the CENTERS FOR DISEASE CONTROL AND PREVENTION / National Center for Health Statistics

Detailed Diagnoses and Procedures, National Hospital Discharge Survey, 1996



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics



Copyright information

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

Suggested Citation

Graves EJ, Kozak LJ. Detailed diagnoses and procedures. National Hospital Discharge Survey, 1996. National Center for Health Statistics. Vital Health Stat 13(138). 1998.

Library of Congress Catalog Card Number 90-13573

For sale by the U.S. Government Printing Office
Superintendent of Documents
Mail Stop: SSOP
Washington, DC 20402-9328
Printed on acid-free paper.

Vital and Health Statistics

Detailed Diagnoses and Procedures, National Hospital Discharge Survey, 1996

Series 13
Data From the National Health
Survey
No. 138

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics

Hyattsville, Maryland
September 1998
DHHS Publication No. 98-1799

National Center for Health Statistics

Edward J. Sondik, Ph.D., *Director*

Jack R. Anderson, *Deputy Director*

Jack R. Anderson, *Acting Associate Director for
International Statistics*

Lester R. Curtin, Ph.D., *Acting Associate Director for
Research and Methodology*

Jennifer H. Madans, Ph.D., *Acting Associate Director for
Analysis, Epidemiology, and Health Promotion*

P. Douglas Williams, *Acting Associate Director for Data
Standards, Program Development, and Extramural Programs*

Edward L. Hunter, *Associate Director for Planning, Budget,
and Legislation*

Jennifer H. Madans, Ph.D., *Acting Associate Director for
Vital and Health Statistics Systems*

Stephen E. Nieberding, *Associate Director for Management*

Charles J. Rothwell, *Associate Director for Data Processing
and Services*

Division of Health Care Statistics

Thomas McLemore, *Acting Division Director*

Robert Pokras, *Chief, Hospital Care Statistics Branch*

Cooperation of the U.S. Bureau of the Census

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies.

In accordance with specifications established by the National Center for Health Statistics, the U.S. Bureau of the Census, under a contractual arrangement, participated in planning the survey and collecting the data.

Contents

Abstract	1
Introduction	1
Methods	1
Data Source	1
Estimation	2
Use of Tables	2
References	3
Appendixes	137
Technical Notes	137
Abstract Form	137
Population Estimates	137
Reliability of Estimates	137
Changes in ICD-9-CM Codes	139

List of Detailed Tables

1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996	4
2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996	32
3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996	59
4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996	117

Appendix Tables

I. Civilian population, by sex, age, and geographic region: United States, 1996	137
II. Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics by selected characteristics: United States, 1996	139
III. Changes in diagnosis codes	140
IV. Changes in procedure codes	144

Appendix Figure

I. Medical abstract for the National Hospital Discharge Survey, 1996	138
--	-----

Abstract

Objectives

This report presents national estimates of the use of non-Federal short-stay hospitals in the United States during 1996. Estimates of first-listed diagnoses, all-listed diagnoses, days of care for first-listed diagnoses, and all-listed procedures are shown by sex and age of patient and geographic region of hospital.

Methods

The estimates are based on data collected through the National Hospital Discharge Survey for 1996. The survey has been conducted annually by the National Center for Health Statistics since 1965. In 1996, data were collected for approximately 282,000 discharges from 480 non-Federal short-stay hospitals. Diagnoses and procedures are presented according to their code number in the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*.

Keywords: *inpatients • diagnoses • procedures • ICD-9-CM*

Detailed Diagnoses and Procedures, National Hospital Discharge Survey, 1996

by Edmund J. Graves and Lola Jean Kozak, Division of Health Care Statistics

Introduction

The purpose of this report is to present data from the 1996 National Hospital Discharge Survey (NHDS). The survey has been conducted continuously by the National Center for Health Statistics (NCHS) since 1965. National estimates of hospital use derived from the NHDS are published for each calendar year by NCHS. This report is intended as a source for researchers and others who need data for detailed diagnostic and procedure categories. Estimates of specific diagnoses and procedures and days of care for first-listed diagnoses are shown by age and sex of patients and geographic region of the hospitals. Similar reports have been published with data for 1977-79 and 1983-95. Other data from the NHDS are also published in Series 13 of *Vital and Health Statistics*.

The NHDS is the principal source for national data on the characteristics of discharges from non-Federal short-stay hospitals. Data from the NHDS are used to examine important topics in public health (1-5) and for a variety of planning, administrative, and evaluation activities by governmental, scientific, academic, and commercial institutions.

Estimates of the number of procedures shown in this report are for inpatients only. Data on ambulatory surgery are collected by means of another survey conducted by NCHS, the National Survey of Ambulatory Surgery (NSAS). The NSAS was implemented in 1994 and covers ambulatory surgery provided in hospitals and freestanding ambulatory surgery centers (6). Data from NSAS are not included in the tables of this report.

Methods

Data Source

NHDS data are collected from a sample of inpatient records acquired from a national sample of hospitals. Because persons with multiple discharges during the year may be sampled more than once, the NHDS produces estimates of discharges, not persons. Only hospitals with an average length of stay of fewer than 30 days for all patients, general hospitals, or children's general hospitals are included in the survey. Federal, military, and Department of Veterans Affairs hospitals, as well as hospital units of institutions (such as prison hospitals) and hospitals with fewer than six beds staffed for patient use, are excluded.

NOTE: This report was prepared in the Division of Health Care Statistics. Elaine Wood, Hospital Care Statistics Branch, verified the data. Rong Cai of the OAO Corporation did the computer programming and produced estimated parameters for relative standard error equations. This report was edited by Gail V. Johnson and typeset by Annette F. Holman of the Publications Branch, Division of Data Services.

Beginning in 1988, the sampling frame for hospitals has been the *SMG Hospital Market Database* (7–9).

The NHDS had a two-stage design until 1988 when it was redesigned (10). Beginning in 1988, all hospitals with 1,000 beds or more or 40,000 discharges or more annually were included in the sample with certainty, while the remaining sample of hospitals was based on a stratified three-stage design. The first stage consisted of the selection of 112 primary sampling units (PSU's) that comprise a probability sample of PSU's used in the 1985–94 National Health Interview Survey (11). The second stage consisted of the selection of noncertainty hospitals from the sample PSU's. At the third stage, a systematic random sample of discharges within hospitals was selected.

For 1996, the sample consisted of 525 hospitals, 18 of which were out of scope (ineligible) because they had gone out of business or otherwise failed to meet the criteria for the NHDS universe. Of the 507 in-scope (eligible) hospitals, 480 (95 percent) responded to the survey. Data were collected for approximately 282,000 patient discharges from the 480 responding hospitals.

Two data collection procedures were used for the survey. One was a manual system in which sample selection and transcription of information from the hospital records to abstract forms were performed by the hospital's staff or by staff of the U.S. Bureau of the Census on behalf of NCHS. Completed forms were sent to NCHS for coding, editing, and post-survey weighting adjustments.

The other data collection procedure was an automated system in which NCHS purchased machine-readable medical record data from commercial organizations, State data systems, hospitals, or hospital associations. Records from these sources were systematically sampled by NCHS. In 1996, approximately 38 percent of respondent hospitals provided data through the automated system.

The medical abstract form for NHDS is shown as [figure I](#) in the appendix. This form and the automated data contain items that relate to the

personal characteristics of the patient. These items include birth date (or age), sex, race, ethnicity, marital status, ZIP code, and expected sources of payment. Administrative items such as admission and discharge dates, discharge status, and medical record number are also included. The medical information about patients includes diagnoses, surgical and nonsurgical procedures, and dates of surgery. (The medical record number, date of birth, and ZIP code are confidential information that are not available to the public.)

For the manual data collection, an ongoing quality control program is undertaken on the coding and keying of data from abstracts to machine-readable form. Approximately 5 percent of the abstracts are independently recoded by an NHDS coder, with discrepancies resolved by the chief coder. The overall error rate for records manually coded by NCHS for the 1996 data year was 3.0 percent for medical (ICD–9–CM) coding and keying and 0.6 percent for demographic coding and keying.

Estimation

Because they are sample data, the survey data must be inflated or weighted in order to produce national estimates. The estimation procedure produces essentially unbiased national estimates and has three basic components: (a) inflation by reciprocals of the probabilities of sample selection, (b) adjustment for nonresponse, and (c) population weighting ratio adjustments. These three components of the final weight are described in more detail in an earlier report (12). Information about the standard errors of statistics for the 1996 NHDS is in the appendix of this report.

Use of Tables

The tables in this report present diagnoses and procedures according to the detailed code numbers of the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) (13), which is updated periodically (see [appendix](#)). Estimates shown are the number of first-listed diagnoses ([table 1](#)), number of days of

care ([table 2](#)), number of all-listed diagnoses ([table 3](#)), and number of all-listed procedures ([table 4](#)) for discharges from short-stay hospitals.

The first-listed diagnosis ([table 1](#)) is the one specified as the principal diagnosis on the face sheet or discharge summary of the medical record. However, if the principal diagnosis is not specified, the first one listed on the face sheet or discharge summary of the medical record is used. Data for all-listed diagnoses ([table 3](#)) include the first-listed diagnosis and other diagnoses appearing in the medical record. A minimum of one and a maximum of seven diagnoses are coded.

Estimates of procedures, which consist of surgical and nonsurgical procedures, are also published. Estimates of all-listed procedures ([table 4](#)) include all occurrences of the procedure coded, regardless of their order in the medical record. Up to four procedures are coded for each discharge. The definitions of other terms related to the NHDS can be found in earlier reports (12).

[Tables 1–3](#) provide information by 3-digit, 4-digit, and 5-digit ICD–9–CM diagnostic codes. [Table 4](#) shows 2-, 3-, and 4-digit procedure codes. Only ICD–9–CM code numbers appear in [tables 1–4](#). Specific titles and more detailed information about the coding scheme can be found in appropriate volumes of the ICD–9–CM (13).

A specific code appears in [tables 1, 3, and 4](#) if the estimate of the total for the code is 5,000 or more. These estimates generally have a relative standard error (a measure of the reliability of the data) of less than 30 percent and a sample size greater than 30. The codes that appear in [table 2](#) for days of care are identical to those in [table 1](#) for first-listed diagnoses. Only an asterisk (*) is shown in [tables 1, 3, and 4](#) if the estimate of diagnoses or procedures for a sex or age group or region is less than 5,000. In [table 2](#) only an asterisk is shown for estimates of days of care derived from fewer than 5,000 estimated discharges.

In addition, all estimates of diagnoses and procedures less than 9,000 are preceded by an asterisk to indicate that they are based on a

relatively small sample of discharge records. These estimates are generally based on fewer than 60 records.

Days-of-care estimates derived from fewer than 9,000 estimated discharges are also preceded by asterisks.

The total for all codes presented at the beginning of each table includes estimates for codes not shown in the table. Likewise, the total for each diagnostic chapter or procedure category includes estimates for all the codes in the chapter or category, whether or not they are shown individually. Estimates for 5-digit codes may not add to the 4-digit total, 4-digit codes to the 3-digit total, or, for procedures, 3-digit codes to the 2-digit total because the totals include estimates for codes not shown or because the data were rounded to the nearest thousand for presentation.

Although 4- and 5-digit diagnostic codes are presented together under the 3-digit codes of which they are a part, the estimates for 4- and 5-digit codes cannot be added to obtain the estimate for the 3-digit code. For example, in [table 1](#), the estimate for first-listed diagnosis code 198 is 123,000. If the estimates for the 4- and 5-digit codes included in 198 are added, the total would be 137,000. This occurs because the estimate of 24,000 for code 198.89 is part of the estimate of 26,000 for code 198.8 and thus cannot be added to it. Similarly, estimates for 3- and 4-digit codes cannot be added together to obtain the estimate for a 2-digit code.

References

1. Bennett TA, Kotelchuck M, Cox CE, et al. Pregnancy-associated hospitalizations in the United States in 1991 and 1992: A comprehensive view of maternal morbidity. *Am J Obstet Gynecol* 178(2):346–54. 1998.
2. Parashar UD, Holman RC, Clarke MJ, et al. Hospitalizations associated with rotavirus diarrhea in the United States, 1993 through 1995: Surveillance based on the new ICD–9–CM rotavirus-specific diagnostic code. *J Infect Dis* 177(1):13–7. 1998.
3. Gillum RF, Gillum BS, Francis CK. Coronary revascularization and cardiac catheterization in the United States: Trends in racial differences. *J Am Coll Cardiol* 29(7):1557–62. 1997.
4. Pappas G, Hadden WC, Kozak LJ, Fisher GF. Potentially avoidable hospitalizations: Inequalities in rates between U.S. socioeconomic groups. *Am J Public Health* 87(5):811–6. 1997.
5. Davis H, Moore RM, Gergen PJ. Cost of hospitalizations associated with sickle cell disease in the United States. *Public Health Rep* 112(1):40–3. 1997.
6. McLemore T, Lawrence L. Plan and operation of the National Survey of Ambulatory Surgery. National Center for Health Statistics. *Vital Health Stat* 1(37). 1997.
7. SMG Marketing Group, Inc. Hospital Market Database. Chicago: Healthcare Information Specialists. 1987.
8. SMG Marketing Group, Inc. Hospital Market Database. Chicago: Healthcare Information Specialists. 1991.
9. SMG Marketing Group, Inc. Hospital Market Database. Chicago: Healthcare Information Specialists. 1994.
10. Haupt BJ, Kozak LJ. Estimates from two survey designs: National Hospital Discharge Survey. National Center for Health Statistics. *Vital Health Stat* 13(111). 1992.
11. Massey JT, Moore TF, Parsons VL, Tadros W. Design and estimation for the National Health Interview Survey, 1985–94. National Center for Health Statistics. *Vital Health Stat* 2(110). 1989.
12. Graves EJ. Detailed diagnoses and procedures, National Hospital Discharge Survey, 1993. National Center for Health Statistics. *Vital Health Stat* 13(122). 1995.
13. Public Health Service and Health Care Financing Administration. International Classification of Diseases, 9th Revision, Clinical Modification. Washington: Public Health Service. 4th ed. 1991.
14. Shah BV, Barnwell BG, Bieler GS. SUDAAN User's Manual: Software for Analysis of Correlated Data, Release 6.40. Research Triangle Park, NC: Research Triangle Institute. 1995.
15. Hansen MH, Hurwitz WN, Madow WG. Sample survey methods and theory, vol 1. New York: John Wiley & Sons. 1953.
16. Cochran WG. Sampling techniques, 3rd ed. New York: John Wiley & Sons. 1977.

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
All codes	30,545	12,110	18,435	2,207	10,325	6,294	11,718	6,665	7,107	11,085	5,688
001–139	845	412	434	153	225	135	332	197	173	316	160
003	*7	*	*	*	*	*	*	*	*	*	*
008	101	43	58	43	13	13	32	21	26	38	16
008.4	27	11	17	*	*	*	16	*8	*7	*8	*
008.45	24	9	14	*	*	*	15	*7	*6	*7	*
008.6	20	11	*9	20	*	*	—	*	10	*6	*
008.61	20	11	*8	19	*	*	—	*	9	*5	*
008.8	49	18	32	19	*8	*7	16	12	*8	23	*6
009	12	*	*7	*	*	*	*	*	*	*5	*
009.0	*6	*	*	—	*	*	*	*	*	*	*
011	*7	*	*	*	*	*	*	*	*	*	*
011.9	*6	*	*	*	*	*	*	*	*	*	*
034	10	*5	*5	*	*5	*	*	*	*	*6	*
034.0	9	*	*5	*	*5	*	*	*	*	*	*
038	355	156	199	19	43	57	236	67	83	145	60
038.0	24	10	14	*	*	*	12	*	*7	*8	*5
038.1	50	24	26	*	*6	13	29	*8	14	20	*8
038.2	10	*	*6	*	*	*	*	*	*	*	*
038.4	94	42	52	*	9	17	66	22	21	34	17
038.40	*8	*	*	*	*	*	*5	*	*	*	*
038.42	53	21	32	*	*5	11	36	14	14	15	11
038.43	*8	*5	*	*	*	*	*6	*	*	*	*
038.49	23	11	12	*	*	*	18	*	*	11	*
038.8	15	*7	*8	*	*	*	10	*	*	*6	*
038.9	157	67	90	9	22	17	109	24	36	73	24
041	11	*	*7	*	*	*	*5	*	*	*	*
042	101	72	29	*	73	24	*	39	11	32	19
047	21	11	10	10	*9	*	*	*	*	*8	*6
047.9	19	10	9	*8	*9	*	*	*	*	*7	*
053	20	*8	12	*	*	*	12	*	*6	*6	*
053.9	12	*	*8	*	*	*	*7	*	*	*	*
054	*7	*	*	*	*	*	*	*	*	*	*
070	19	*9	10	*	12	*	*	*5	*	*7	*
070.1	*6	*	*	*	*	*	*	*	*	*	*
070.5	*7	*	*	*	*	*	*	*	*	*	*
075	9	*	*	*	*5	*	*	*	*	*	*
078	*6	*	*	*	*	*	*	*	*	*	*
079	60	29	32	38	10	*6	*7	17	10	20	14
079.9	56	27	29	34	9	*6	*6	16	*9	18	13
079.99	56	27	29	34	9	*6	*6	16	*9	18	13
112	18	*8	10	*	*	*	10	*	*	*7	*5
112.4	*6	*	*	*	*	*	*5	*	*	*	*
117	*6	*	*	*	*	*	*	*	*	*	*
135	*8	*	*	—	*	*	*	*	*	*	*
136	12	*8	*	*	*5	*	*	*	*	*6	*
136.3	*8	*6	*	*	*	*	*	*	*	*	*
140–239	1,805	693	1,112	40	327	611	826	454	371	623	356
141	*5	*	*	—	*	*	*	*	*	*	*
150	11	*7	*	—	*	*	*8	*	*	*	*
151	25	17	*8	—	*	*6	18	*7	*	*7	*8

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
151.0	*6	*5	*	—	*	*	*	*	*	*	*
151.8	*6	*	*	—	*	*	*5	*	*	*	*
151.9	*5	*	*	—	*	*	*	*	*	*	*
153	108	49	59	*	*	29	76	23	26	43	16
153.1	*8	*	*5	*	*	*	*	*	*	*	*
153.2	*5	*	*	—	*	*	*	*	*	*	*
153.3	25	14	11	—	*	*8	17	*	*	10	*6
153.4	22	*8	13	—	*	*	17	*	*7	*6	*
153.6	18	*9	*9	—	*	*	13	*	*5	*7	*
153.8	*7	*	*	—	*	*	*6	*	*	*	*
153.9	16	*6	10	—	*	*	12	*	*	10	*
154	43	19	24	—	*6	*8	29	14	10	11	*8
154.0	14	*7	*7	—	*	*	11	*	*	*	*
154.1	25	10	15	—	*	*	17	*9	*7	*	*5
155	10	*	*5	*	*	*	*	*	*	*	*
155.0	*6	*	*	*	*	*	*	*	*	*	*
156	*7	*	*	—	—	*	*6	*	*	*	*
157	37	21	16	—	*	14	21	*9	*9	13	*7
157.0	11	*6	*5	—	*	*	*6	*	*	*	*
157.8	*8	*	*6	—	*	*	*6	*	*	*	*
157.9	15	11	*	—	*	*8	*7	*	*	*6	*
161	*8	*6	*	—	*	*	*	*	*	*	*
162	180	104	76	*	10	63	107	42	35	65	38
162.2	17	11	*6	—	*	*7	9	*	*5	*7	*
162.3	54	29	26	—	*	17	36	12	10	19	14
162.5	28	16	12	—	*	*8	18	*7	*5	10	*6
162.8	20	12	*8	—	*	*8	11	*9	*	*6	*
162.9	57	34	23	*	*	22	31	12	10	22	13
170	*6	*	*	*	*	*	*	*	*	*	*
171	*6	*	*	*	*	*	*	*	*	*	*
172	*5	*	*	—	*	*	*	*	*	*	*
173	*5	*	*	—	*	*	*	*	*	*	*
174	123	—	123	*	14	53	55	29	28	42	25
174.2	*6	—	*6	—	*	*	*	*	*	*	*
174.4	24	—	24	—	*	*9	12	*	*8	*7	*
174.8	31	—	31	—	*	13	14	12	*6	*7	*6
174.9	47	—	47	*	*6	23	19	*7	10	20	10
180	15	—	15	*	*7	*6	*	*	*	*8	*
180.9	*9	—	*9	*	*	*	*	*	*	*5	*
182	34	—	34	—	*	13	18	*8	10	9	*7
182.0	34	—	34	—	*	13	18	*8	10	9	*7
183	31	—	31	*	*	13	13	11	*	11	*
183.0	30	—	30	*	*	13	13	10	*	10	*
185	90	90	—	—	*	30	60	19	23	31	17
188	50	37	13	—	*	11	38	16	10	14	10
188.4	*5	*	*	—	—	*	*	*	*	*	*
188.8	17	13	*	—	*	*	12	*7	*	*	*
188.9	16	12	*	—	—	*	12	*	*	*	*
189	29	17	13	*	*	10	17	*6	*	14	*
189.0	23	13	10	*	*	*6	13	*6	*	10	*
189.1	*5	*	*	—	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
191	27	18	9	*	*7	11	*6	*7	*6	*7	*7
191.1	*6	*	*	*	*	*	*	*	*	*	*
191.2	*7	*	*	—	*	*	*	*	*	*	*
193	17	*	13	*	*7	*5	*	*6	*	*6	*
195	*6	*	*	*	*	*	*	*	*	*	*
196	20	*6	14	—	*	*8	*8	*6	*	*7	*
197	132	52	80	*	*9	46	75	38	29	42	22
197.0	30	13	17	*	*	9	15	11	*7	*8	*
197.2	30	11	19	—	*	10	17	*9	*8	*8	*6
197.6	17	*	13	—	*	*6	9	*5	*	*5	*
197.7	43	20	23	*	*	17	24	12	*7	16	*8
198	123	53	70	*	11	43	67	36	30	39	18
198.3	38	15	23	*	*	15	19	9	11	12	*5
198.5	49	25	25	*	*	17	29	16	11	17	*6
198.8	26	*9	17	*	*	*7	14	*8	*6	*5	*6
198.89	24	*9	15	*	*	*7	14	*8	*5	*	*6
199	14	*	10	*	*	*	*7	*	*	*7	*
199.0	*5	*	*	*	*	*	*	*	*	*	*
199.1	*9	*	*7	—	*	*	*	*	*	*	*
200	17	*8	*9	*	*	*	10	*	*	*5	*6
200.0	9	*5	*	*	*	*	*5	*	*	*	*
201	*7	*	*	*	*	*	*	*	*	*	*
201.9	*5	*	*	*	*	*	*	*	*	*	*
202	39	20	19	*	*	13	21	10	*9	12	*7
202.8	33	16	16	*	*	11	18	*8	*8	10	*7
202.80	17	*8	*9	*	*	*5	10	*	*	*5	*
202.83	*6	*	*	—	*	*	*	*	*	*	*
203	18	10	*8	—	*	*6	10	*5	*	*	*
203.0	17	10	*7	—	*	*6	10	*5	*	*	*
203.00	17	10	*7	—	*	*6	10	*	*	*	*
204	26	13	13	*9	*	*	*9	10	*	*8	*6
204.0	16	*8	*7	*9	*	*	*	*	*	*5	*
204.00	13	*7	*5	*7	*	*	*	*	*	*	*
204.1	9	*	*	*	—	*	*7	*	*	*	*
204.10	9	*	*	—	—	*	*7	*	*	*	*
205	24	11	13	*	*	*9	9	*7	*	*9	*
205.0	17	*8	10	*	*	*6	*7	*	*	*7	*
205.00	15	*7	*8	*	*	*	*6	*	*	*5	*
205.1	*6	*	*	*	*	*	*	*	*	*	*
205.10	*6	*	*	*	*	*	*	*	*	*	*
210	*8	*	*	*	*	*	*	*	*	*	*
210.2	*7	*	*	—	*	*	*	*	*	*	*
211	29	13	16	*	*	*7	19	9	*	11	*
211.3	23	10	13	—	*	*6	16	*8	*	*8	*
218	222	—	222	—	125	91	*7	43	42	85	51
218.0	31	—	31	—	17	14	*	*7	*5	9	*9
218.1	64	—	64	—	35	28	*	18	14	17	16
218.2	23	—	23	—	12	10	*	*	*	12	*
218.9	104	—	104	—	60	39	*	15	20	46	23
220	38	—	38	*	18	15	*	*7	*9	15	*8
225	17	*6	11	*	*	*	*8	*6	*	*	*
225.2	10	*	*8	*	*	*	*5	*	*	*	*
226	11	*	11	*	*7	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
227	19	*7	12	*	*5	*7	*6	*	*	*5	*6
227.1	11	*	*7	—	*	*	*	*	*	*	*
227.3	*5	*	*	*	*	*	*	*	*	*	*
228	*7	*	*	*	*	*	*	*	*	*	*
228.0	*5	*	*	*	*	*	*	*	*	*	*
233	25	*	24	—	11	*7	*6	*	*6	9	*
233.0	*8	—	*8	—	*	*	*	*	*	*	*
233.1	15	—	15	—	9	*	*	*	*	*6	*
235	11	*	*7	*	*	*	*7	*	*	*	*
236	*6	*	*	—	*	*	*	*	*	*	*
237	*6	*	*	*	*	*	*	*	*	*	*
238	19	12	*7	*	*	*	14	*	*	*7	*
238.7	13	9	*	*	*	*	11	*	*	*	*
239	10	*5	*5	*	*	*	*6	*	*	*	*
240-279	1,272	519	753	155	250	305	563	267	299	496	210
241	11	*	*8	*	*	*	*	*	*	*	*
241.1	*6	*	*5	*	*	*	*	*	*	*	*
242	14	*	11	*	*6	*	*	*	*	9	*
242.0	*7	*	*5	—	*	*	*	*	*	*6	*
242.00	*6	*	*	—	*	*	*	*	*	*5	*
242.9	*5	*	*	—	*	*	*	*	*	*	*
244	18	*	16	*	*	*7	*8	*	*	*8	*
244.9	17	*	14	*	*	*6	*7	*	*	*7	*
250	503	233	270	24	120	161	198	101	112	204	86
250.0	140	56	84	*7	27	52	53	25	26	65	23
250.00	39	16	23	*	*	16	18	*7	*7	17	*9
250.01	29	9	19	*	*5	9	12	*6	*6	11	*6
250.02	35	15	21	*	*7	15	13	*	*7	19	*
250.03	37	17	20	*	11	12	10	*8	*7	19	*
250.1	100	48	52	13	55	23	*8	19	24	42	15
250.10	10	*	*5	*	*5	*	*	*	*	*5	*
250.11	35	17	18	*7	22	*	*	9	*6	12	*7
250.12	*7	*	*	—	*	*	*	*	*	*	*
250.13	49	25	24	*6	26	13	*	*7	15	21	*6
250.2	10	*5	*	—	*	*	*	*	*	*5	*
250.3	*7	*	*	*	*	*	*	*	*	*	*
250.4	38	14	24	*	*7	16	14	*6	*5	14	13
250.40	14	*5	*9	—	*	*6	*6	*	*	*	*6
250.41	13	*	*8	*	*	*6	*	*	*	*	*
250.43	*8	*	*	—	*	*	*	*	*	*	*
250.5	*8	*	*	—	*	*	*	*	*	*	*
250.51	*5	*	*	—	*	*	*	*	*	*	*
250.6	30	14	17	—	*7	11	12	*7	*8	11	*
250.60	*9	*	*	—	*	*	*	*	*	*	*
250.61	11	*	*6	—	*	*	*	*	*	*	*
250.63	*7	*	*	—	*	*	*	*	*	*	*
250.7	66	34	32	—	*	24	39	12	16	25	13
250.70	23	14	9	—	*	*7	16	*	*	10	*6
250.71	23	11	12	—	*	11	11	*	*7	*6	*6
250.72	*9	*	*	—	*	*	*7	*	*	*	*
250.73	11	*	*7	—	*	*	*6	*	*	*6	*
250.8	91	48	43	*	11	25	54	24	23	32	11
250.80	28	17	11	—	*	*6	21	*8	*6	11	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
250.81	30	16	14	*	*5	10	14	*9	*8	9	*
250.82	17	*7	10	—	*	*	12	*	*	*6	*
250.83	16	*8	*8	*	*	*	*8	*	*	*6	*
250.9	13	*6	*6	*	*	*	*	*	*	*	*
250.93	*6	*	*	*	*	*	*	*	*	*	*
251	*9	*	*	*	*	*	*	*	*	*	*
251.2	*6	*	*	*	*	*	*	*	*	*	*
252	*6	*	*	*	*	*	*	*	*	*	*
252.0	*6	*	*	—	*	*	*	*	*	*	*
253	*7	*	*	*	*	*	*	*	*	*	*
253.6	*5	*	*	—	*	*	*	*	*	*	*
255	9	*	*	*	*	*	*	*	*	*	*
255.4	*6	*	*	*	*	*	*	*	*	*	*
263	15	*6	10	*	*	*	12	*	*	*6	*
263.9	13	*	*8	*	*	*	10	*	*	*	*
272	25	12	14	*	*7	12	*6	*6	*6	*9	*
272.0	11	*5	*6	*	*	*7	*	*	*	*	*
272.4	11	*	*6	—	*	*	*	*	*	*	*
274	9	*6	*	—	*	*	*6	*	*	*	*
274.0	*7	*	*	—	*	*	*	*	*	*	*
275	15	*7	*9	*	*	*	9	*	*	*	*
275.4	13	*6	*7	*	*	*	*9	*	*	*	*
276	575	213	362	118	76	87	293	119	149	223	85
276.0	11	*	*7	*	*	*	*7	*	*	*	*
276.1	61	20	41	*	*	11	42	*7	20	22	12
276.2	10	*	*6	*	*	*	*	*	*	*	*
276.5	434	162	272	110	58	58	208	98	107	169	60
276.6	*6	*	*	*	*	*	*	*	*	*	*
276.7	14	*6	*8	*	*	*	*9	*	*	*	*
276.8	33	*9	25	*	*7	*6	20	*5	*7	15	*6
277	14	*9	*5	*5	*7	*	*	*7	*	*	*
277.0	9	*6	*	*	*5	—	—	*5	*	*	*
277.00	9	*6	*	*	*5	—	—	*5	*	*	*
278	16	*	12	*	10	*6	*	*	*	*	*5
278.0	15	*	12	*	9	*	*	*	*	*	*5
278.00	*7	*	*	—	*	*	*	*	*	*	*
278.01	*8	*	*7	*	*6	*	*	*	*	*	*
280–289	333	143	191	53	88	65	128	83	80	114	57
280	42	13	29	*	*8	*8	25	11	9	16	*5
280.0	25	10	16	—	*	*	17	*6	*	10	*
280.9	16	*	13	*	*	*	*8	*5	*	*5	*
281	10	*	*7	*	*	*	*5	*	*	*6	*
281.9	*6	*	*	*	*	*	*	*	*	*	*
282	61	29	32	15	42	*	*	18	17	20	*6
282.6	56	27	29	14	40	*	*	17	16	18	*5
282.62	50	24	26	*9	39	*	*	15	14	17	*
283	*6	*	*	*	*	*	*	*	*	*	*
284	23	10	13	*	*	*8	9	*6	*7	*7	*
284.8	20	*9	11	*	*	*8	*7	*5	*	*7	*
285	68	24	44	*	9	16	40	14	17	21	16
285.1	18	*	13	*	*	*	13	*	*5	*	*
285.9	48	18	30	*	*6	13	26	9	11	16	11

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
286	17	10	*6	*	*	*	*6	*	*	*	*
286.0	*5	*	*	*	*	*	*	*	*	*	*
287	34	14	20	*8	*	*7	14	*8	*7	11	*8
287.3	13	*5	*8	*	*	*	*	*	*	*	*
287.4	*5	*	*	*	*	*	*	*	*	*	*
287.5	13	*	*8	*	*	*	*8	*	*	*6	*
288	49	23	26	9	*8	13	19	13	11	16	*8
288.0	44	22	22	*8	*8	12	16	11	11	14	*7
288.8	*5	*	*	*	*	*	*	*	*	*	*
289	23	13	10	*9	*7	*	*5	*	*	10	*
289.2	12	*7	*5	*6	*6	*	—	*	*	*5	*
290–319	1,943	1,004	939	81	1,145	401	315	555	516	669	202
290	70	26	44	—	*	*5	64	11	22	31	*7
290.0	20	*7	13	—	*	*	19	*5	*6	*6	*
290.2	20	*7	13	—	—	*	18	*	*9	10	*
290.20	*7	*	*	—	—	*	*7	*	*	*	—
290.21	13	*	9	—	—	*	12	*	*7	*	*
290.3	11	*	*7	—	—	*	10	*	*	*	*
290.4	16	*7	*9	—	—	*	14	*	*	*9	*
290.40	*6	*	*	—	—	*	*6	*	*	*	*
291	90	73	17	—	50	31	*8	23	34	21	12
291.0	22	19	*	—	9	10	*	*7	*	*8	*
291.8	59	47	12	—	36	19	*	13	28	10	*8
292	59	35	24	*	38	10	11	19	18	15	*8
292.0	24	12	12	—	18	*	*	10	*8	*	*
292.1	*7	*6	*	*	*	*	*	*	*	*	*
292.8	23	14	*9	*	11	*6	*6	*8	*5	*7	*
292.81	10	*	*	*	*	*	*	*	*	*	*
292.84	*8	*7	*	—	*7	*	*	*	*	*	*
293	29	13	16	*	*6	*7	16	10	*7	*6	*6
293.0	15	*6	9	*	*	*	12	*	*	*	*
293.8	13	*7	*6	—	*6	*	*	*7	*	*	*
293.83	10	*6	*	—	*	*	*	*7	*	*	*
294	20	*8	12	*	*	*	17	*	*	10	*
294.8	11	*5	*6	*	*	*	10	*	*	*6	*
295	257	147	110	*	167	68	20	92	63	70	31
295.3	75	46	29	*	47	21	*7	32	10	20	13
295.30	15	*8	*7	*	9	*	*	*	*	*	*
295.32	18	12	*6	—	10	*6	*	*9	*	*	*
295.34	41	26	15	—	27	11	*	20	*	12	*
295.6	22	12	11	—	11	9	*	*8	*7	*5	*
295.62	*6	*	*	—	*	*	*	*	*	*	*
295.64	14	*8	*6	—	*8	*6	*	*6	*	*	*
295.7	116	63	54	*	76	30	10	37	36	29	14
295.70	59	29	31	*	38	17	*	17	17	19	*6
295.74	51	32	19	—	35	11	*5	18	18	*8	*7
295.9	33	20	13	*	25	*6	*	12	*8	10	*
295.90	17	11	*6	*	14	*	*	*	*6	*7	*
295.94	13	*7	*6	—	9	*	*	10	*	*	*
296	597	223	374	32	352	118	95	151	169	230	47
296.2	204	87	116	17	122	31	35	46	54	88	15
296.20	91	41	50	*6	55	17	13	25	32	25	*8
296.22	23	9	13	*	17	*	*	*	*	18	*
296.23	53	21	32	*	29	*7	13	9	13	28	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
296.24	35	15	20	*	19	*6	*8	9	*7	16	*
296.3	216	70	146	*8	126	44	38	63	66	79	*8
296.30	67	23	44	*	38	15	11	20	26	19	*
296.32	18	*	13	*	13	*	*	*	*8	*7	*
296.33	85	27	59	*	52	16	13	24	20	37	*
296.34	43	14	29	*	20	10	12	16	12	14	*
296.4	43	17	26	*	21	14	*7	12	12	10	*9
296.40	23	*9	14	*	11	*7	*	*	*7	*6	*6
296.44	15	*6	*9	*	*7	*	*	*6	*	*	*
296.5	42	13	29	*	25	11	*	*9	12	16	*5
296.50	24	*7	17	*	13	*7	*	*	*8	*9	*
296.53	*8	*	*6	*	*	*	*	*	*	*	*
296.54	*8	*	*	—	*7	*	*	*	*	*	*
296.6	45	17	29	*	31	*9	*	10	10	20	*6
296.60	25	11	15	*	17	*	*	*	*8	11	*
296.63	*7	*	*	*	*5	*	*	*	*	*	*
296.64	10	*	*7	*	*7	*	*	*	*	*	*
296.7	30	11	19	*	18	*6	*	*7	11	9	*
296.8	11	*6	*5	*	*6	*	*	*	*	*5	*
296.89	*5	*	*	*	*	*	*	*	*	*	*
297	*7	*	*	—	*	*	*	*	*	*	*
298	81	40	40	*	45	13	21	27	12	27	15
298.9	74	38	36	*	41	12	19	24	11	25	15
300	71	29	41	*	38	16	14	20	19	22	*9
300.0	28	12	16	*	10	*7	11	*8	*7	10	*
300.00	17	*8	10	*	*	*	*8	*	*	*6	*
300.01	9	*	*5	*	*6	*	*	*	*	*	*
300.1	10	*	*8	*	*7	*	*	*	*	*	*
300.11	*7	*	*5	*	*	*	*	*	*	*	*
300.4	21	*8	12	*	12	*	*	*7	*7	*6	*
300.9	*6	*	*	*	*	*	*	*	*	*	*
301	13	*7	*7	*	9	*	*	*5	*	*	*
301.8	*7	*	*	—	*6	*	*	*	*	*	*
303	206	158	48	*	140	56	10	43	53	88	22
303.0	44	35	10	—	27	16	*	10	13	14	*8
303.00	12	9	*	—	*8	*	*	*	*	*5	*
303.01	32	25	*6	—	18	12	*	*8	9	*9	*6
303.9	162	123	39	*	113	40	*8	34	40	74	14
303.90	53	41	12	*	35	14	*	*	*9	37	*
303.91	107	80	26	*	77	25	*	29	31	37	9
304	115	74	41	*	94	18	*	46	18	45	*6
304.0	31	20	11	—	25	*5	*	18	*	*7	*
304.00	*6	*	*	—	*	*	—	*	*	*	—
304.01	25	16	*9	—	20	*	*	17	*	*	*
304.2	33	22	11	—	28	*	*	13	*	15	*
304.20	10	*6	*	—	*8	*	*	*	*	*8	*
304.21	22	16	*6	—	19	*	*	13	*	*6	*
304.7	18	12	*6	—	17	*	*	11	*	*	*
304.71	15	9	*6	—	14	*	*	11	*	*	*
304.8	14	9	*5	—	12	*	*	*	*	11	*
304.80	*6	*	*	—	*	*	*	*	*	*	*
304.81	*8	*5	*	—	*7	*	*	*	*	*7	*
304.9	10	*6	*	—	*7	*	*	*	*	*5	*
304.90	9	*6	*	—	*7	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
305	67	43	24	*	38	22	*5	17	13	28	9
305.0	28	20	*8	*	14	9	*	9	*	11	*
305.00	20	14	*6	*	11	*6	*	*7	*	*7	*
305.01	*7	*6	*	—	*	*	*	*	*	*	*
305.1	15	10	*	—	*7	*8	*	*	*	*8	*
305.6	*7	*	*	—	*6	*	—	*	*	*	*
305.9	9	*6	*	*	*6	*	*	*	*	*6	*
305.90	*7	*	*	*	*	*	*	*	*	*5	*
307	10	*	*8	*	*6	*	*	*	*	*	*
308	*8	*	*7	*	*	*	*	*	*	*	*
309	96	45	51	*8	72	10	*5	37	28	20	11
309.0	35	16	19	*	29	*	*	15	*8	*9	*
309.2	13	*6	*7	*	9	*	*	*5	*	*	*
309.28	9	*	*5	*	*7	*	*	*	*	*	*
309.4	21	12	10	*	14	*	*	10	*	*	*
309.8	13	*	9	*	*9	*	*	*	*5	*	*
309.81	12	*	9	*	*8	*	—	*	*5	*	*
309.9	*9	*	*5	*	*7	*	*	*	*	*	*
310	10	*	*5	*	*	*	*7	*	*	*	*
310.9	*7	*	*	—	*	*	*6	*	*	*	*
311	79	34	45	*7	51	14	*8	26	20	29	*6
312	26	20	*6	*7	18	*	*	12	*9	*	*
312.3	14	10	*	*	11	*	*	*9	*	*	*
312.30	*8	*	*	*	*7	*	—	*6	*	*	—
312.9	*8	*7	*	*	*6	—	—	*	*	*	*
314	14	11	*	10	*	—	*	*	*7	*	*
314.0	13	10	*	10	*	—	*	*	*7	*	—
314.01	11	*9	*	*9	*	—	—	*	*6	*	—
320–389	512	225	287	89	138	99	187	116	100	198	97
320	*6	*	*	*	*	*	*	*	*	*	*
322	*7	*	*	*	*	*	*	*	*	*	*
322.9	*6	*	*	*	*	*	*	*	*	*	*
323	*5	*	*	*	*	*	*	*	*	*	*
331	36	17	19	*	*	*	26	*6	*8	16	*7
331.0	16	*9	*7	—	—	*	16	*	*	10	*
331.3	*7	*	*	*	*7	*	*	*	*	*	*
331.4	11	*5	*6	*	*	*	*	*	*	*	*
332	20	*7	13	*	*	*	18	*	*	*9	*
332.0	20	*7	13	*	*	*	18	*	*	*9	*
333	9	*5	*	*	*	*	*	*	*	*	*
337	*6	*	*	—	*	*	*	*	*	*	*
340	20	*	15	—	9	10	*	*	*5	*8	*
341	*6	*5	*	*	*	—	*	—	*	*	*
342	17	*9	*8	*	*	*5	*8	*	*	10	*
342.9	15	*8	*7	*	*	*	*8	*	*	*9	*
342.90	14	*8	*6	*	*	*	*8	*	*	*9	*
344	11	*8	*	*	*	*	*	*	*	*	*
345	59	26	32	18	21	*7	12	15	12	20	12
345.1	12	*6	*7	*	*	*	*	*	*	*	*
345.10	10	*	*5	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
345.3	17	*9	*8	*	*	*	*6	*	*	*7	*
345.4	*6	*	*	*	*	*	*	*	*	*	*
345.5	*5	*	*	*	*	*	*	*	*	*	*
345.9	14	*6	*8	*5	*	*	*	*	*	*	*
345.90	11	*5	*6	*	*	*	*	*	*	*	*
346	37	*7	31	*	23	11	*	*9	*8	17	*
346.8	*6	*	*	*	*	*	*	*	*	*	*
346.9	25	*	20	*	16	*7	*	*	*	13	*
346.90	16	*	12	*	12	*	*	*	*	*9	*
346.91	*8	*	*8	—	*	*	—	*	*	*	*
348	32	15	17	*	*9	*7	12	*6	*8	11	*7
348.1	*8	*	*	*	*	*	*	*	*	*	*
348.3	10	*5	*	*	*	*	*7	*	*	*	*
348.8	*5	*	*	*	*	*	*	*	*	*	*
349	14	*6	*8	*	*8	*	*	*	*	*7	*
349.0	*7	*	*	*	*5	*	*	*	*	*	*
353	*7	*	*	*	*	*	*	*	*	*	*
355	*6	*	*	*	*	*	*	*	*	*	*
357	*7	*5	*	*	*	*	*	*	*	*	*
357.0	*6	*	*	*	*	*	*	*	*	*	*
359	*6	*	*	*	*	*	*	*	*	*	*
361	13	*6	*7	*	*	*	*7	*	*	*9	*
361.0	10	*	*6	*	*	*	*6	*	*	*7	*
362	*9	*	*5	*	*	*	*	*	*	*	*
366	22	9	13	—	*	*	19	*6	*	*8	*8
366.1	*6	*	*	—	—	*	*5	*	*	*	*
366.9	10	*6	*	—	*	*	9	*	—	*7	*
368	*5	*	*	*	*	*	*	*	*	*	*
376	10	*5	*	*5	*	*	*	*	*	*	*
376.0	*9	*5	*	*5	*	*	*	*	*	*	*
376.01	*9	*5	*	*5	*	*	*	*	*	*	*
381	10	*6	*	*9	*	*	*	*	*	*	*
382	20	12	*7	18	*	*	*	*	*	*9	*
382.9	18	12	*6	17	*	*	*	*	*	*8	*
386	29	9	20	*	*	*7	19	*5	*8	11	*
386.1	*5	*	*	*	*	*	*	*	*	*	*
386.3	18	*7	11	*	*	*	13	*	*	*7	*
386.30	15	*5	10	*	*	*	10	*	*	*6	*
390–459	6,107	3,101	3,006	25	413	1,706	3,963	1,388	1,477	2,235	1,008
396	14	*5	*8	*	*	*	9	*	*	*	*
398	*8	*	*	*	*	*	*7	*	*	*	*
398.9	*8	*	*	*	*	*	*7	*	*	*	*
398.91	*8	*	*	—	—	*	*7	*	*	*	*
401	199	80	119	*	33	80	85	46	40	83	28
401.0	30	10	20	—	*7	12	11	*6	*	14	*6
401.9	165	68	97	*	27	66	72	38	36	69	22
402	99	34	65	*	*	21	74	23	21	39	17
402.0	*7	*	*	—	*	*	*	*	*	*	*
402.9	90	31	59	*	*	19	69	21	20	34	16
402.90	14	*	10	—	*	*6	*8	*	*	*6	*
402.91	76	27	49	*	*	13	61	18	17	27	14

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
403	85	42	43	*	19	30	35	14	16	41	13
403.0	*5	*	*	—	*	*	*	*	*	*	*
403.9	79	40	39	*	17	28	33	14	15	37	13
403.91	75	39	36	*	17	27	31	13	14	35	13
404	34	15	19	—	*	10	22	*8	*8	12	*6
404.9	31	14	16	—	*	*9	19	*8	*7	11	*5
404.93	23	10	13	—	*	*7	15	*7	*	*7	*
410	825	496	329	*	42	282	500	182	201	299	144
410.0	33	20	13	—	*	13	19	*7	9	10	*6
410.01	30	19	12	—	*	12	17	*7	*9	9	*5
410.1	151	92	59	—	9	60	82	37	33	57	24
410.11	139	86	53	—	*8	54	76	35	33	50	21
410.12	*8	*	*	—	*	*	*	*	*	*	*
410.2	19	10	9	—	*	*6	11	*5	*	*6	*
410.21	19	10	9	—	*	*6	11	*5	*	*6	*
410.3	19	12	*7	—	*	*7	11	*	*	*6	*
410.31	19	12	*6	—	*	*7	10	*	*	*6	*
410.4	176	109	67	*	13	73	90	36	48	65	26
410.41	168	103	65	*	13	69	86	36	47	61	24
410.42	*6	*	*	—	*	*	*	*	*	*	*
410.5	19	10	*9	—	*	*7	12	*6	*	*	*5
410.51	18	9	*9	—	*	*6	11	*5	*	*	*5
410.7	276	162	113	*	12	80	184	63	68	92	53
410.70	*6	*	*	—	—	*	*	*	*	*	*
410.71	259	152	107	*	11	75	173	61	67	83	48
410.72	11	*7	*	—	*	*	*7	*	*	*6	*
410.8	22	14	*9	—	*	*	17	*6	*	10	*
410.81	22	14	*8	—	*	*	17	*6	*	10	*
410.9	105	62	43	*	*	29	73	15	24	47	19
410.90	18	10	*8	—	*	*8	10	*	*	*8	*
410.91	83	49	34	*	*	20	60	11	21	37	14
411	265	135	130	*	18	91	155	80	63	87	35
411.1	231	117	114	—	16	81	134	70	56	74	31
411.8	33	18	15	*	*	9	20	9	*7	12	*
411.81	12	*9	*	—	*	*5	*	*	*	*	*
411.89	21	*9	12	*	*	*	16	*6	*	9	*
412	9	*5	*	—	*	*	*	*	*	*	*
413	98	47	50	—	9	39	49	17	31	38	11
413.9	95	46	49	—	*9	38	48	16	30	38	11
414	1,066	642	424	*	48	421	594	249	247	378	192
414.0	1,041	626	416	*	47	414	578	244	242	367	189
414.00	180	94	86	*	*7	67	105	35	31	85	29
414.01	815	498	317	*	39	329	446	202	195	265	152
414.02	47	34	13	—	*	18	28	*7	16	17	*8
414.8	13	*8	*5	*	*	*	*8	*	*	*	*
414.9	11	*8	*	*	*	*	*7	*	*	*7	*
415	67	27	41	*	11	16	40	15	17	18	16
415.1	65	26	39	*	11	15	38	14	17	18	16
415.11	*7	*	*	—	*	*	*	*	*	*	*
415.19	58	24	34	*	9	14	35	13	16	16	12
416	*9	*	*6	*	*	*	*	*	*	*	*
420	15	9	*6	*	*	*	*6	*	*	*	*
420.9	15	9	*6	*	*	*	*6	*	*	*	*
420.90	9	*5	*	*	*	*	*	*	*	*	*
420.91	*5	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
421	16	*8	*9	*	*	*	*8	*5	*	*5	*
421.0	16	*8	*8	*	*	*	*8	*	*	*	*
423	12	*8	*	*	*	*6	*	*	*	*6	*
423.9	10	*5	*	*	*	*	*	*	*	*	*
424	85	42	44	*	10	21	53	19	22	29	15
424.0	35	11	24	*	*8	*9	18	*7	*8	13	*7
424.1	49	30	19	*	*	13	34	12	13	16	*8
425	32	19	13	*	*	15	12	*8	*8	12	*
425.4	28	17	12	*	*	14	11	*6	*7	12	*
426	58	31	28	*	*6	10	42	14	16	20	9
426.0	24	15	*9	*	*	*	22	*5	*7	*8	*
426.1	16	*8	*9	*	*	*	13	*	*	*6	*
426.13	*9	*	*	*	*	*	*7	*	*	*	*
427	618	290	328	*6	45	140	427	151	149	213	105
427.0	35	13	22	*	*	10	22	*8	*8	15	*
427.1	45	31	13	*	*	15	24	16	10	9	9
427.3	325	154	171	*	17	73	234	77	84	114	50
427.31	286	133	153	*	14	61	211	68	72	102	45
427.32	38	21	17	*	*	13	22	9	12	12	*5
427.4	10	*6	*	*	*	*5	*	*	*	*	*
427.41	9	*6	*	*	*	*	*	*	*	*	*
427.5	17	*8	*9	*	*	*	11	*	*	*8	*
427.6	23	11	11	*	*	*5	13	*7	*	*6	*5
427.69	18	10	*7	*	*	*	11	*	*	*5	*5
427.8	153	60	92	*	12	27	111	36	37	54	26
427.81	61	23	38	*	*	*6	54	13	12	28	*8
427.89	92	37	54	*	11	21	56	23	24	26	18
427.9	10	*	*6	*	*	*	*7	*	*	*	*
428	877	381	496	*	29	154	692	189	226	339	123
428.0	870	377	493	*	29	153	686	187	224	337	122
428.1	*5	*	*	*	*	*	*	*	*	*	*
429	22	12	10	*	*	*	15	*	*	10	*
429.2	9	*	*	—	*	*	*6	*	*	*	*
429.4	*5	*	*	*	*	*	*	*	*	*	*
430	24	*6	18	*	*6	12	*6	*	*6	*8	*6
431	63	29	34	*	*	12	46	11	14	26	11
432	20	13	*7	*	*	*	13	*	*	*8	*5
432.1	18	12	*7	*	*	*	12	*	*	*7	*
433	166	92	74	*	*	34	127	40	36	69	21
433.1	122	67	55	*	*	25	93	29	26	52	15
433.10	104	59	45	*	*	21	79	24	23	45	12
433.11	18	*8	10	—	*	*	14	*5	*	*8	*
433.3	37	21	17	—	*	*7	30	10	*8	14	*6
433.30	31	17	13	—	*	*5	25	*8	*6	12	*
433.31	*7	*	*	—	*	*	*5	*	*	*	*
434	302	141	161	*	9	52	242	72	75	103	52
434.0	17	*6	11	—	*	*	14	*6	*	*6	*
434.01	15	*5	9	—	*	*	12	*5	*	*	*
434.1	41	18	24	—	*	*7	34	11	10	10	11
434.10	*7	*	*	—	*	*	*6	*	*	*	*
434.11	35	15	20	—	*	*6	28	9	*8	*7	10
434.9	244	117	127	*	*7	43	193	55	60	88	41
434.90	*5	*	*	*	*	*	*	*	*	*	*
434.91	239	114	125	*	*7	41	190	53	59	86	40

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
435	179	73	106	*	*7	37	136	42	47	65	25
435.3	*9	*5	*	—	*	*	*7	*	*	*	*
435.8	*6	*	*	—	*	*	*	*	*	*	*
435.9	161	64	97	*	*6	33	122	38	42	58	23
436	158	66	93	—	*	25	130	20	40	66	32
437	28	10	18	*	*	*8	17	*	*7	10	*5
437.1	*8	*	*6	*	—	*	*7	*	*	*	*
437.3	*6	*	*	*	*	*	*	*	*	*	*
438	15	*8	*7	—	*	*	11	*	*	*5	*
440	137	77	60	*	*	38	95	32	30	56	19
440.2	119	68	51	—	*	32	83	30	25	48	15
440.20	15	10	*6	—	*	*	10	*	*	*8	*
440.21	22	15	*7	—	*	*8	14	*	*	*8	*5
440.22	16	*9	*7	—	*	*	11	*	*	*	*
440.23	18	*8	10	—	*	*6	12	*6	*	*7	*
440.24	47	25	22	—	*	9	37	12	10	20	*
440.3	*7	*	*	—	*	*	*	*	*	*	*
441	58	44	14	*	*	9	47	13	12	23	9
441.0	*9	*	*	—	*	*	*6	*	*	*	*
441.01	*6	*	*	—	*	*	*	*	*	*	*
441.3	9	*8	*	—	*	*	*7	*	*	*	*
441.4	36	30	*6	*	—	*	31	*8	*8	16	*
442	15	10	*6	—	*	*	10	*	*	*5	*
442.3	*8	*6	*	—	*	*	*6	*	*	*	*
443	23	11	12	—	*	*	17	*6	*	12	*
443.9	19	*9	10	—	*	*	15	*	*	11	*
444	53	26	26	*	*6	14	33	12	12	19	9
444.2	40	20	21	—	*	*8	28	10	10	15	*6
444.21	*7	*	*5	—	*	*	*	*	*	*	*
444.22	33	18	15	—	*	*7	23	*9	*9	12	*
444.8	*6	*	*	*	*	*	*	*	*	*	*
444.81	*6	*	*	*	*	*	*	*	*	*	*
446	10	*	*6	*	*	*	*	*	*	*	*
447	17	*9	*9	*	*	*7	*8	*6	*	*5	*
447.1	11	*6	*	*	*	*	*6	*	*	*	*
451	46	20	27	*	*8	13	25	11	15	16	*6
451.0	11	*	*6	—	*	*	*	*	*6	*	*
451.1	25	10	15	*	*	*6	15	*6	*7	*9	*
451.11	*9	*	*6	*	*	*	*	*	*	*	*
451.19	16	*8	*8	—	*	*	10	*	*	*5	*
451.2	*7	*	*	—	*	*	*	*	*	*	*
453	141	64	76	*	22	37	81	28	42	45	27
453.8	135	62	73	*	21	35	79	27	40	42	26
454	18	*6	12	*	*	*	9	*	*5	*	*
454.2	*9	*	*6	—	*	*	*6	*	*	*	*
454.9	*5	*	*	—	*	*	*	*	*	*	*
455	24	13	12	—	9	*7	*8	*5	*6	10	*
455.2	12	*7	*5	—	*	*	*6	*	*	*	*
456	*6	*	*	—	*	*	*	*	*	*	*
458	62	30	32	*	*	13	45	13	19	18	12
458.0	33	16	17	—	*	*9	23	*7	10	11	*5
458.9	28	13	15	*	*	*	21	*6	*8	*7	*7

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
459	11	*5	*6	*	*	*	*8	*	*	*	*
460–519	3,238	1,544	1,694	653	450	584	1,550	667	797	1,228	546
461	10	*	*	*	*	*	*	*	*	*	*
462	10	*	*6	*	*6	*	*	*	*	*6	*
463	14	*	*9	*	*7	*	*	*	*	*6	*
464	24	15	*9	19	*	*	*	*	*6	*8	*5
464.4	17	11	*6	17	*	—	—	*	*5	*6	*
465	34	15	18	22	*	*	*5	*8	*7	12	*7
465.9	32	15	17	21	*	*	*5	*7	*6	12	*7
466	226	118	108	147	17	21	41	43	56	86	41
466.0	95	40	55	18	17	20	40	16	22	46	12
466.1	131	78	53	129	*	*	*	27	34	40	30
470	*6	*	*	—	*	*	*	*	*	*	*
473	14	*6	*8	*	*6	*	*	*	*	*5	*
474	27	14	13	20	*6	*	—	17	*	*6	*
474.0	13	*5	*8	*9	*	*	—	*6	*	*	*
474.1	13	*9	*	12	*	*	—	10	*	*	*
474.10	9	*5	*	*9	*	—	—	*7	*	*	*
475	10	*	*6	*	*7	*	*	*	*	*	*
478	28	17	11	*8	11	*6	*	*8	*	11	*5
478.2	10	*7	*	*	*	*	*	*	*	*	*
478.29	*8	*6	*	*	*	*	*	*	*	*	*
478.7	*6	*	*	*	*	*	*	*	*	*	*
480	42	23	20	34	*	*	*	*5	11	12	13
480.1	23	12	11	23	*	—	—	*	*7	*7	*7
480.9	18	9	*9	10	*	*	*	*	*	*5	*6
481	54	30	24	*	13	15	23	11	15	19	9
482	286	147	139	12	23	36	215	35	82	131	38
482.0	20	9	11	*	*	*	15	*	*	11	*
482.1	43	24	19	*	*5	*7	25	*7	11	19	*6
482.2	29	10	19	*	*	*	20	*	13	*6	*6
482.3	14	*8	*7	*	*	*	11	*	*	*6	*
482.30	*6	*	*	*	*	*	*	*	*	*	*
482.39	*5	*	*	*	*	*	*	*	*	*	*
482.4	43	22	21	*	*	*	34	*5	*7	22	*9
482.8	121	65	56	*	*6	12	101	10	39	62	10
482.83	78	45	33	*	*	*7	66	*5	33	35	*
482.89	37	17	20	*	*	*	30	*	*5	23	*
482.9	15	*9	*6	*	*	*	9	*5	*	*5	*
483	*8	*	*	*	*	*	*	*	*	*	*
483.0	*7	*	*	*	*	*	*	*	*	*	*
485	39	14	25	16	*	*	15	*	15	16	*
486	773	357	416	123	95	116	440	163	203	275	132
487	21	9	12	*	*6	*	*9	*	*	10	*
487.0	*6	*	*	*	*	*	*	*	*	*	*
487.1	15	*8	*7	*	*	*	*6	*	*	*8	*
490	18	*6	12	*6	*	*	*5	*	*	*9	*
491	404	176	228	*	21	108	274	83	110	166	45
491.2	401	175	226	*	20	107	272	83	110	165	44
491.20	17	*6	11	*	*	*7	*8	*	*	*9	*
491.21	384	169	216	*	18	101	264	80	107	156	41

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
492	31	18	13	—	*	11	20	*8	*9	9	*5
492.8	28	16	12	—	*	9	19	*7	*8	*8	*
493	474	195	279	195	132	88	59	126	116	146	86
493.0	37	22	15	31	*5	*	*	10	*	*	21
493.00	13	*7	*6	11	*	*	*	*	*	*	*8
493.01	24	14	9	20	*	*	—	*8	*	*	13
493.2	71	20	51	*	11	27	33	19	15	25	12
493.20	48	13	35	*	*6	19	23	11	10	19	*8
493.21	23	*7	16	—	*6	*8	10	*8	*5	*6	*
493.9	364	153	211	164	115	59	26	97	95	119	52
493.90	190	68	122	64	67	37	22	48	44	68	31
493.91	173	85	88	100	47	22	*	49	52	50	22
494	*7	*	*	*	*	*	*	*	*	*	*
496	130	65	65	*	*	41	87	28	25	53	24
507	156	87	69	*6	12	12	126	27	32	68	28
507.0	154	86	67	*	12	12	125	27	31	67	28
510	11	*7	*	*	*	*	*5	*	*	*	*
510.9	10	*6	*	*	*	*	*	*	*	*	*
511	68	32	36	*	*8	15	44	13	15	24	16
511.0	11	*	*6	—	*	*	*5	*	*	*	*
511.8	9	*	*	*	*	*	*7	*	*	*	*
511.9	46	22	24	*	*6	10	31	10	10	16	11
512	38	22	16	*	16	10	12	*8	*8	17	*5
512.1	*6	*	*	*	*	*	*	*	*	*	*
512.8	29	18	12	*	14	*8	*7	*6	*5	14	*
513	*6	*5	*	*	*	*	*	*	*	*	*
513.0	*5	*	*	*	*	*	*	*	*	*	*
515	19	9	10	*	*	*	11	*	*5	*8	*
516	11	*	*7	*	*	*	*7	*	*	*	*
516.3	*5	*	*	*	*	*	*	*	*	*	*
516.8	*6	*	*	*	*	*	*	*	*	*	*
518	209	106	103	*9	22	60	118	37	39	87	45
518.0	*7	*	*	*	*	*	*	*	*	*	*
518.4	10	*	*6	*	*	*	*7	*	*	*	*
518.8	180	94	87	*7	17	52	105	30	35	77	38
518.81	164	86	78	*	15	48	98	27	31	72	34
518.82	*9	*	*	*	*	*	*	*	*	*	*
518.89	*7	*	*	*	*	*	*	*	*	*	*
519	14	*7	*7	*6	*	*	*	*	*	*	*
519.1	*7	*	*	*	*	*	*	*	*	*	*
520–579	2,906	1,295	1,612	205	746	757	1,198	629	690	1,050	536
524	16	*6	10	*	14	*	*	*	*5	*	*
524.1	*7	*	*	*	*6	*	*	*	*	*	*
524.10	*7	*	*	*	*6	*	*	*	*	*	*
527	13	*	*9	*	*	*	*8	*	*	*	*
527.2	*8	*	*5	*	*	*	*	*	*	*	*
528	11	*6	*6	*	*	*	*	*	*	*	*
530	186	89	97	26	35	58	66	38	39	80	29
530.1	43	18	25	*	12	13	15	11	*8	18	*6
530.10	16	*7	10	*	*	*6	*	*	*	*8	*
530.11	20	9	11	*	*5	*6	*8	*	*	*7	*
530.19	*6	*	*	*	*	*	*	*	*	*	*
530.2	18	11	*7	*	*	*	10	*	*	*6	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
530.3	9	*	*	*	*	*	*7	*	*	*	*
530.5	10	*	*6	*	*	*6	*	*	*	*5	*
530.7	15	11	*	*	*	*6	*6	*	*	*7	*
530.8	86	39	47	23	13	28	22	15	19	39	13
530.81	80	37	43	23	12	26	20	14	18	35	13
531	91	46	45	*	10	27	53	15	27	29	21
531.0	27	11	16	*	*	*8	16	*	9	*8	*6
531.00	25	10	15	*	*	*8	14	*	*8	*8	*6
531.4	42	24	18	*	*	11	27	*8	11	14	9
531.40	42	23	18	*	*	11	27	*7	11	14	9
531.5	*6	*	*	—	*	*	*	*	*	*	*
531.50	*5	*	*	—	*	*	*	*	*	*	*
531.9	*9	*	*	*	*	*	*	*	*	*	*
531.90	*8	*	*	*	*	*	*	*	*	*	*
532	75	46	29	*	14	20	40	15	18	30	12
532.0	20	12	*7	—	*	*6	10	*	*6	*8	*
532.00	19	12	*7	—	*	*6	10	*	*6	*8	*
532.4	32	22	11	*	*6	*8	19	*6	*7	13	*7
532.40	32	22	11	*	*6	*7	19	*6	*7	13	*7
532.9	*7	*	*	—	*	*	*	*	*	*	*
532.90	*6	*	*	—	*	*	*	*	*	*	*
533	23	9	14	*	*	*8	11	*	*9	10	*
533.4	*8	*	*6	*	*	*	*	*	*	*	*
533.40	*7	*	*6	*	*	*	*	*	*	*	*
533.9	*8	*	*	—	*	*	*	*	*	*	*
533.90	*8	*	*	—	*	*	*	*	*	*	*
535	166	76	90	*	38	42	83	32	33	75	25
535.0	44	20	24	*	11	10	22	*8	12	19	*5
535.00	29	12	17	*	*9	*6	12	*6	*8	13	*
535.01	15	*8	*7	*	*	*	10	*	*	*6	*
535.1	12	*	*8	*	*	*	*6	*	*	*6	*
535.10	*6	*	*	*	*	*	*	*	*	*	*
535.11	*6	*	*	*	*	*	*	*	*	*	*
535.3	10	*7	*	*	*	*	*	*	*	*	*
535.31	*6	*	*	*	*	*	*	*	*	*	*
535.4	44	19	26	*	9	*8	27	*9	*7	19	10
535.40	21	*5	15	*	*6	*5	9	*	*	9	*
535.41	24	13	11	*	*	*	17	*5	*	9	*6
535.5	47	20	27	*	10	15	22	*7	9	25	*6
535.50	32	11	21	*	*8	10	13	*	*7	18	*
535.51	15	9	*6	*	*	*	*8	*	*	*7	*
535.6	*8	*	*	*	*	*	*	*	*	*	*
536	26	11	14	*	*8	*8	*7	*7	*	11	*
536.2	*8	*	*6	*	*	*	*	*	*	*	*
536.3	*6	*	*	*	*	*	*	*	*	*	*
536.8	9	*	*	*	*	*	*	*	*	*	*
537	16	*8	*8	*	*	*	*9	*	*	*6	*
537.0	10	*6	*	*	*	*	*5	*	*	*	*
540	217	134	83	57	115	30	15	40	49	66	63
540.0	45	28	17	14	17	*8	*6	9	10	12	14
540.1	26	17	9	*	10	*7	*5	*	*6	10	*7
540.9	146	89	57	38	88	15	*	27	33	44	42
541	*9	*	*5	*	*7	*	*	*	*	*	*
550	57	48	*9	9	*9	11	28	22	10	16	*9
550.1	23	19	*	*	*	*	14	*7	*5	*7	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
550.10	17	14	*	*	*	*	*9	*6	*	*	*
550.9	33	29	*	*5	*7	*8	13	14	*	9	*5
550.90	19	16	*	*	*	*6	*7	9	*	*6	*
550.91	*5	*	*	*	*	*	*	*	*	*	*
550.92	*8	*8	*	*	*	*	*	*	*	*	*
552	40	14	27	*	*7	11	21	11	11	12	*7
552.1	10	*	*6	*	*	*	*	*	*	*	*
552.2	20	*7	13	*	*	*6	10	*	*6	*7	*
552.20	*6	*	*	—	*	*	*	*	*	*	*
552.21	12	*	*9	*	*	*	*6	*	*	*	*
553	75	28	47	*	15	26	33	20	17	30	*8
553.1	*6	*	*	*	*	*	*	*	*	*	*
553.2	45	17	28	*	*7	17	21	13	11	17	*5
553.20	10	*	*7	*	*	*	*	*	*	*	*
553.21	33	14	18	*	*	13	15	10	*9	11	*
553.3	22	*8	14	*	*5	*7	10	*	*6	10	*
555	34	12	22	*	19	9	*	10	*8	10	*6
555.0	9	*	*7	—	*	*	*	*	*	*	*
555.1	*9	*	*5	*	*	*	*	*	*	*	*
555.2	*6	*	*	*	*	*	*	*	*	*	*
555.9	10	*	*5	*	*7	*	*	*	*	*	*
556	28	13	15	*	14	*8	*6	*6	*8	*8	*5
556.9	16	*7	*9	*	*8	*	*	*	*6	*6	*
557	37	13	23	*	*	*8	27	*8	11	11	*6
557.0	21	*7	14	*	*	*	15	*5	*5	*7	*
557.1	*7	*	*	—	*	*	*6	*	*	*	*
557.9	*8	*	*6	—	*	*	*6	*	*	*	*
558	178	72	106	59	47	24	48	36	42	81	19
558.9	172	69	103	59	46	22	44	34	41	79	18
560	262	108	154	*8	35	62	157	55	63	87	57
560.1	49	18	30	*	*	10	34	*8	*9	20	11
560.2	15	*9	*6	*	*	*	11	*	*	*7	*
560.3	26	11	15	*	*	*	18	*8	*5	*9	*
560.39	25	11	14	*	*	*	17	*7	*5	*8	*
560.8	88	35	53	*	13	24	49	17	26	24	21
560.81	65	26	40	*	11	19	34	12	18	17	17
560.89	22	*9	13	*	*	*5	15	*5	*7	*7	*
560.9	82	33	49	*	13	23	45	17	20	27	18
562	210	89	121	*	21	54	135	51	50	78	31
562.1	209	88	121	*	21	53	134	51	50	78	31
562.10	23	9	14	—	*	*7	14	*6	*6	10	*
562.11	120	53	68	*	18	39	62	33	29	40	19
562.12	47	20	27	—	*	*	42	*8	11	19	*9
562.13	18	*6	12	—	*	*	16	*	*	*8	*
564	48	19	29	*6	14	*8	21	9	13	17	9
564.0	24	13	11	*	*	*	13	*	*6	*7	*6
564.1	14	*	12	*	*8	*	*	*	*	*6	*
565	*7	*	*	*	*	*	*	*	*	*	*
566	19	14	*	*	*8	*6	*	*	*	*6	*
567	18	*8	11	*	*	*7	*6	*	*7	*	*
567.2	11	*	*6	*	*	*	*	*	*	*	*
568	13	*	10	*	*6	*	*	*	*	*	*
568.0	12	*	9	*	*5	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
569	73	26	47	*	10	15	46	16	19	26	12
569.3	15	*5	9	*	*	*	9	*	*5	*	*
569.4	*7	*	*	*	*	*	*	*	*	*	*
569.6	11	*5	*6	*	*	*	*7	*	*	*	*
569.69	9	*	*5	*	*	*	*5	*	*	*	*
569.8	29	10	20	*	*	*	22	*	*7	12	*5
569.83	*7	*	*	—	*	*	*5	*	*	*	*
569.85	11	*	*7	—	*	*	*8	*	*	*	*
570	*8	*	*	*	*	*	*	*	*	*	*
571	83	52	30	*	22	37	23	15	12	36	20
571.1	11	*9	*	—	*5	*6	*	*	*	*	*
571.2	35	27	*9	—	11	19	*6	*7	*6	13	10
571.3	*5	*	*	—	*	*	*	*	*	*	*
571.5	26	11	15	*	*	*9	14	*	*	15	*
572	29	15	14	*	*	13	10	*7	*	10	*7
572.2	19	9	9	*	*	*8	*8	*	*	*7	*
573	9	*	*5	*	*	*	*	*	*	*	*
574	386	115	271	*	118	126	139	98	84	131	74
574.0	131	47	83	*	36	43	51	33	25	43	29
574.00	116	40	76	*	34	37	44	29	23	38	26
574.01	15	*7	*8	—	*	*5	*7	*	*	*5	*
574.1	182	46	136	*	61	66	54	51	41	60	29
574.10	176	44	132	*	58	65	52	50	39	58	29
574.11	*5	*	*	*	*	*	*	*	*	*	*
574.2	36	9	27	*	11	10	14	*7	10	11	*8
574.20	35	*9	26	*	10	10	14	*6	9	11	*8
574.3	*8	*	*5	—	*	*	*	*	*	*	*
574.4	11	*	*9	*	*	*	*	*	*	*	*
574.40	*6	*	*	*	*	*	*	*	*	*	*
574.5	19	*8	11	—	*	*	12	*	*	10	*
574.50	10	*	*6	—	*	*	*5	*	*	*	*
574.51	10	*	*	—	*	*	*7	*	*	*5	*
575	61	23	38	*	24	17	20	11	20	22	*9
575.0	30	14	16	—	10	*8	12	*	9	10	*6
575.1	25	*8	16	*	10	*8	*6	*	*9	9	*
576	29	12	17	*	*5	*9	14	*	*6	12	*6
576.1	12	*6	*6	*	*	*5	*	*	*	*	*
576.2	11	*	*6	*	*	*	*6	*	*	*	*
577	184	83	101	*	72	58	51	33	57	63	30
577.0	146	66	80	*	56	45	44	26	45	52	24
577.1	30	14	15	*	13	11	*6	*5	11	*8	*5
577.2	*5	*	*	*	*	*	*	*	*	*	*
578	134	58	76	*	10	26	96	28	34	44	29
578.0	13	*6	*7	*	*	*	*9	*	*	*	*
578.1	21	10	11	*	*	*	14	*	*	*8	*5
578.9	101	42	59	*	*7	21	73	21	25	34	20
579	*5	*	*	*	*	*	*	*	*	*	*
580–629	1,673	526	1,146	69	590	413	600	334	383	666	290
581	*9	*	*	*	*	*	*	*	*	*	*
581.9	*7	*	*	*	*	*	*	*	*	*	*
584	80	40	40	*	10	15	55	17	17	33	13
584.9	74	36	38	*	10	14	50	16	16	30	12
585	35	20	16	*	*6	11	18	9	*6	13	*7

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
586	10	*	*5	*	*	*	*7	*	*	*	*
590	125	26	99	17	57	25	26	22	31	48	24
590.1	80	16	64	10	39	12	19	15	22	26	17
590.10	80	16	64	10	39	12	19	15	22	26	17
590.8	42	9	33	*6	17	12	*7	*7	*8	21	*6
590.80	42	9	33	*6	17	12	*7	*7	*8	21	*6
591	13	*6	*7	*	*5	*	*	*	*	*	*
592	176	107	69	*	85	64	26	34	43	72	26
592.0	46	21	24	*	22	17	*6	10	9	19	*7
592.1	128	85	44	*	63	46	19	25	33	52	19
593	37	18	19	*7	*8	9	12	9	11	10	*6
593.4	12	*5	*7	*	*	*	*	*	*	*	*
593.7	*6	*	*	*	*	*	—	*	*	*	*
593.70	*6	*	*	*	*	*	—	*	*	*	*
593.8	*5	*	*	*	*	*	*	*	*	*	*
593.9	*8	*6	*	*	*	*	*6	*	*	*	*
594	*8	*6	*	*	*	*	*	*	*	*	*
594.1	*6	*	*	—	*	*	*	*	*	*	*
595	23	9	14	*	*	*5	16	*6	*8	*7	*
595.0	11	*	*8	*	*	*	*7	*	*5	*	*
596	16	13	*	*	*	*	12	*	*	*	*
596.0	*7	*7	*	*	*	*	*6	*	*	*	*
598	*8	*7	*	*	*	*	*	*	*	*	*
598.9	*5	*	*	*	*	*	*	*	*	*	*
599	332	112	220	24	36	50	222	68	75	142	47
599.0	297	88	208	23	29	45	199	61	68	126	41
599.7	28	19	10	*	*7	*	17	*5	*6	12	*
600	106	106	—	—	*	21	85	23	31	31	20
601	*7	*7	—	—	*	*	*	*	*	*	*
604	12	12	—	*	*5	*	*	*	*	*	*
604.9	11	11	—	*	*	*	*	*	*	*	*
604.90	*7	*7	—	*	*	*	*	*	*	*	*
607	*9	*9	—	*	*	*	*	*	*	*	*
607.8	*7	*7	—	*	*	*	*	*	*	*	*
607.84	*6	*6	—	—	*	*	*	*	*	*	*
608	*8	*8	—	*	*	*	*	*	*	*	*
611	28	*	27	*	17	*7	*	9	*	*8	*6
611.1	17	*	16	*	13	*	*	*7	*	*	*
614	67	—	67	*	59	*5	*	13	12	30	12
614.0	*6	—	*6	*	*6	*	*	*	*	*	*
614.1	*7	—	*7	*	*5	*	*	*	*	*	*
614.2	*5	—	*5	—	*	*	—	*	*	*	*
614.3	*9	—	*9	*	*8	*	—	*	*	*	*
614.6	20	—	20	—	18	*	*	*	*	*8	*
614.9	16	—	16	*	15	*	—	*	*	*7	*
615	*7	—	*7	—	*6	*	—	*	*	*	*
616	*8	—	*8	*	*5	*	*	*	*	*	*
617	85	—	85	*	65	20	*	11	21	39	14
617.0	46	—	46	—	30	15	*	*5	12	22	*7
617.1	14	—	14	*	12	*	—	*	*	*	*
617.3	18	—	18	—	16	*	*	*	*5	*8	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
618	152	—	152	*	34	60	57	23	33	63	32
618.0	41	—	41	*	*5	16	19	*6	*9	17	10
618.1	11	—	11	—	*5	*	*	*	*	*	*
618.2	20	—	20	—	*7	*9	*	*	*6	*6	*
618.3	21	—	21	—	*	*6	11	*6	*5	*6	*
618.4	16	—	16	—	*7	*5	*	*	*	*7	*
618.5	24	—	24	—	*	11	12	*	*	12	*6
618.8	13	—	13	—	*5	*5	*	*	*	*9	*
620	73	—	73	*	51	14	*	17	12	32	13
620.0	*5	—	*5	*	*	*	—	*	*	*	*
620.1	21	—	21	*	14	*	*	*	*	10	*
620.2	39	—	39	*	28	*7	*	11	*5	16	*7
621	16	—	16	—	*	*8	*	*6	*	*5	*
621.0	*5	—	*5	—	*	*	*	*	*	*	*
621.3	*8	—	*8	—	*	*5	*	*	*	*	*
622	12	—	12	—	*9	*	*	*	*	*6	*
622.1	12	—	12	—	*8	*	*	*	*	*6	*
625	92	—	92	*	50	28	14	12	21	39	21
625.3	10	—	10	*	*8	*	—	*	*	*	*
625.6	51	—	51	—	14	23	14	*6	12	18	15
625.9	24	—	24	*	22	*	*	*	*	16	*
626	75	—	75	*	48	25	*	*8	17	36	13
626.2	52	—	52	—	35	17	—	*7	11	26	*8
626.8	21	—	21	*	12	*7	*	*	*5	9	*
627	11	—	11	—	*	*6	*	*	*	*5	*
627.1	*9	—	*9	—	*	*	*	*	*	*	*
630–677	536	—	536	*	534	*	—	118	116	206	96
632	16	—	16	*	16	—	—	*	*	*6	*
633	38	—	38	—	38	—	—	*7	*8	17	*6
633.1	35	—	35	—	35	—	—	*6	*7	16	*6
634	29	—	29	*	29	—	—	12	*	10	*
634.9	23	—	23	*	23	—	—	11	*	*6	*
634.91	18	—	18	*	18	—	—	*9	*	*	*
635	10	—	10	*	10	*	—	*	*	*	*
635.9	*8	—	*8	—	*8	*	—	*	*	*	*
637	*8	—	*8	—	*8	*	—	*	*	*	*
640	*7	—	*7	—	*7	—	—	*	*	*	*
640.0	*6	—	*6	—	*6	—	—	*	*	*	*
640.03	*6	—	*6	—	*6	—	—	*	*	*	*
641	20	—	20	—	20	—	—	*5	*6	*	*
641.1	*5	—	*5	—	*5	—	—	*	*	*	*
641.13	*5	—	*5	—	*5	—	—	*	*	*	*
641.2	*6	—	*6	—	*6	—	—	*	*	*	*
641.23	*6	—	*6	—	*6	—	—	*	*	*	*
642	23	—	23	*	23	*	—	*6	*6	*9	*
642.3	*8	—	*8	—	*8	—	—	*	*	*	*
642.33	*8	—	*8	—	*8	—	—	*	*	*	*
642.4	*6	—	*6	—	*6	*	—	*	*	*	*
642.43	*5	—	*5	—	*5	*	—	*	*	*	*
643	34	—	34	—	34	—	—	*7	*8	16	*
643.0	10	—	10	—	10	—	—	*	*	*5	*
643.03	10	—	10	—	10	—	—	*	*	*5	*
643.1	20	—	20	—	20	—	—	*	*	*9	*
643.13	20	—	20	—	20	—	—	*	*	*9	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
644	133	—	133	*	133	*	—	24	29	50	30
644.0	106	—	106	*	106	*	—	18	23	39	25
644.03	106	—	106	*	106	*	—	18	23	39	25
644.1	27	—	27	—	27	—	—	*5	*5	11	*
644.13	27	—	27	—	27	—	—	*5	*5	11	*
646	57	—	57	*	57	—	—	9	13	24	10
646.6	40	—	40	*	40	—	—	*7	*8	16	*9
646.63	38	—	38	*	38	—	—	*6	*7	15	*9
646.8	13	—	13	—	13	—	—	*	*	*6	*
646.83	13	—	13	—	13	—	—	*	*	*6	*
647	*8	—	*8	—	*8	*	—	*	*	*	*
648	77	—	77	*	76	*	—	19	18	29	11
648.0	9	—	9	—	9	—	—	*	*	*	*
648.03	9	—	9	—	9	—	—	*	*	*	*
648.4	*8	—	*8	*	*8	—	—	*	*	*	*
648.43	*7	—	*7	*	*7	—	—	*	*	*	*
648.8	13	—	13	—	13	—	—	*	*	*6	*
648.83	13	—	13	—	13	—	—	*	*	*5	*
648.9	39	—	39	*	38	*	—	10	*9	14	*6
648.93	37	—	37	*	37	*	—	10	*9	14	*5
654	*9	—	*9	—	*9	*	—	*	*	*	*
658	9	—	9	—	9	—	—	*	*	*	*
659	10	—	10	—	10	—	—	*	*	*	*
659.8	*6	—	*6	—	*6	—	—	*	*	*	*
659.83	*6	—	*6	—	*6	—	—	*	*	*	*
670	*7	—	*7	—	*7	—	—	*	*	*	*
670.0	*7	—	*7	—	*7	—	—	*	*	*	*
670.04	*7	—	*7	—	*7	—	—	*	*	*	*
674	*6	—	*6	—	*6	—	—	*	*	*	*
680–709	452	236	216	45	126	120	162	107	115	145	84
681	20	12	*8	*	*6	*6	*5	*	*6	*7	*
681.0	10	*7	*	*	*5	*	*	*	*	*	*
681.00	9	*6	*	*	*	*	*	*	*	*	*
681.1	10	*5	*	*	*	*	*	*	*	*	*
681.10	9	*5	*	*	*	*	*	*	*	*	*
682	304	164	140	26	90	86	102	73	79	94	58
682.0	20	12	*8	*7	*6	*6	*	*	*	*6	*5
682.1	*5	*	*	*	*	*	*	*	*	*	*
682.2	21	11	10	*	*7	*5	*6	*	*5	*7	*6
682.3	44	23	20	*	23	10	*8	12	11	12	*9
682.4	19	14	*5	*	*8	*	*6	*6	*	*7	*
682.5	*6	*	*	*	*	*	*	*	*	*	*
682.6	138	68	71	*7	30	45	57	34	37	47	21
682.7	42	25	17	*	*8	12	21	11	14	*9	*8
682.9	*5	*	*	*	*	*	*	*	*	*	*
686	*6	*	*	*	*	*	*	*	*	*	*
693	*7	*	*	*	*	*	*	*	*	*	*
693.0	*6	*	*	*	*	*	*	*	*	*	*
695	*7	*	*	*	*	*	*	*	*	*	*
695.1	*5	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
707	69	30	39	*	10	15	43	13	18	25	13
707.0	32	16	16	*	*5	*7	19	*6	*7	15	*
707.1	34	14	20	*	*	*8	22	*7	11	9	*6
708	*5	*	*	*	*	*	*	*	*	*	*
709	*6	*	*	*	*	*	*	*	*	*	*
710–739	1,506	638	868	37	370	430	668	300	357	560	288
710	23	*	19	*	12	*5	*	*	*	11	*
710.0	17	*	15	*	10	*	*	*	*	*8	*
711	15	*8	*7	*	*6	*	*	*	*	*7	*
711.0	15	*8	*6	*	*5	*	*	*	*	*7	*
711.06	*5	*	*	*	*	*	*	*	*	*	*
714	33	11	22	*	*5	9	18	*8	*5	11	9
714.0	30	9	21	*	*	9	17	*7	*	10	*9
715	395	145	250	—	13	90	293	68	116	131	80
715.1	34	12	21	—	*	*7	23	*5	10	13	*6
715.15	9	*	*6	—	*	*	*7	*	*	*	*
715.16	23	*8	15	—	*	*	16	*	*6	10	*
715.3	147	58	89	—	*	37	108	28	47	46	25
715.35	41	19	23	—	*	11	30	*8	15	10	*8
715.36	100	37	63	—	*	24	75	19	31	34	16
715.8	*5	*	*	—	*	*	*	*	*	*	*
715.89	*5	*	*	—	*	*	*	*	*	*	*
715.9	203	71	132	—	*6	42	155	34	57	68	45
715.95	56	22	33	—	*	13	40	12	16	16	12
715.96	133	44	89	—	*	27	104	20	36	48	29
716	21	11	11	*	*	*9	*7	*	*	*9	*
716.1	*6	*	*	—	*	*	*	*	*	*	*
716.9	10	*5	*	*	*	*5	*	*	*	*	*
716.96	*5	*	*	*	*	*	*	*	*	*	*
717	21	13	*8	*	16	*	*	11	*	*	*
717.8	14	9	*	*	13	*	—	*7	*	*	*
717.83	11	*7	*	*	10	*	—	*5	*	*	*
718	23	11	13	*	14	*	*	10	*5	*6	*
718.3	9	*	*	*	*6	*	*	*	*	*	*
718.31	*5	*	*	*	*	*	*	*	*	*	*
719	31	9	22	*	*8	*	16	*7	*	14	*
719.4	19	*5	14	*	*7	*	*7	*	*	*9	*
719.45	10	*	*8	*	*	*	*	*	*	*6	*
719.7	*5	*	*5	—	—	*	*5	*	*	*	*
721	58	28	30	*	*9	24	24	*7	14	27	10
721.0	19	10	*9	*	*	10	*5	*	*	10	*
721.1	12	*6	*5	—	*	*6	*	*	*	*6	*
721.3	21	*8	13	*	*	*6	13	*	*7	*7	*
722	340	183	157	*	156	131	52	63	71	144	62
722.0	55	31	24	—	28	25	*	*9	12	28	*6
722.1	217	120	97	*	108	77	31	44	47	88	38
722.10	215	119	96	*	107	77	31	43	46	87	38
722.4	*8	*	*6	—	*	*	*	*	*	*	*
722.5	21	9	12	—	*	*8	9	*	*	*7	*8
722.52	21	9	12	—	*	*8	9	*	*	*7	*8
722.7	16	10	*6	—	*5	*7	*	*	*	*6	*
722.71	9	*6	*	—	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
722.73	*6	*	*	—	*	*	*	*	*	*	*
722.8	10	*	*6	—	*	*	*	*	*	*6	*
722.83	9	*	*5	—	*	*	*	*	*	*5	*
722.9	*7	*	*	—	*	*	*	*	*	*	*
722.93	*6	*	*	—	*	*	*	*	*	*	*
723	21	12	9	*	*8	*8	*	*	*	*8	*
723.0	*9	*6	*	—	*	*	*	*	*	*	*
723.4	*7	*	*	—	*	*	*	*	*	*	*
724	121	50	72	*	23	35	62	22	32	42	26
724.0	57	26	31	*	*	16	37	13	13	19	12
724.02	55	25	30	*	*	16	35	13	13	19	11
724.2	25	*6	19	*	*7	9	*9	*	*7	9	*6
724.4	15	*8	*7	—	*6	*	*	*	*	*5	*
724.5	13	*	10	*	*	*	*7	*	*	*	*
726	45	23	22	*	13	16	14	13	10	9	12
726.1	12	*6	*6	—	*	*	*5	*	*	*	*5
726.10	11	*6	*5	—	*	*	*5	*	*	*	*5
726.2	12	*6	*7	—	*	*5	*	*5	*	*	*
726.5	*6	*	*6	—	*	*	*5	*	*	*	*
726.6	*5	*	*	*	*	*	*	*	*	*	*
727	22	9	13	*	*9	*5	*	*6	*6	*7	*
727.0	11	*5	*6	*	*7	*	*	*	*	*	*
727.05	*6	*	*	*	*	*	*	*	*	*	*
728	19	*8	10	*	*6	*	*8	*	*	*7	*
728.8	11	*5	*6	*	*	*	*	*	*	*	*
728.89	*5	*	*	*	*	*	*	*	*	*	*
729	35	14	21	*	11	11	12	*	10	13	*7
729.1	*9	*	*5	*	*	*	*	*	*	*	*
729.5	14	*8	*7	*	*	*6	*	*	*	*	*
729.8	*6	*	*	*	*	*	*	*	*	*	*
730	53	31	23	*	*9	17	24	14	13	18	*8
730.0	21	13	*8	*	*	*7	9	*7	*	*7	*
730.07	11	*8	*	*	*	*	*	*	*	*	*
730.1	*8	*	*	*	*	*	*	*	*	*	*
730.2	22	12	10	*	*	*6	10	*6	*	*9	*
730.27	11	*6	*5	*	*	*	*7	*	*	*	*
732	*7	*	*	*5	*	*	—	*	*	*	*
733	175	48	126	*	32	40	102	33	36	75	31
733.1	89	15	74	*	*	10	75	14	19	38	18
733.13	52	*9	43	*	*	*	47	10	12	22	*8
733.14	14	*	11	*	*	*	11	*	*	*5	*
733.15	*8	*	*7	—	*	*	*6	*	*	*	*
733.19	*8	*	*7	—	*	*	*8	*	*	*	*
733.4	23	12	12	*	*7	9	*7	*5	*	11	*
733.42	19	10	9	*	*	*9	*6	*	*	9	*
733.6	17	*5	12	*	*	*8	*	*	*6	*5	*
733.8	33	13	20	*	13	10	10	*6	*5	15	*7
733.81	*6	*	*	*	*	*	*	*	*	*	*
733.82	28	10	17	—	11	*9	*8	*	*	12	*6
733.9	*5	*	*	*	*	*	*	*	*	*	*
735	*9	*	*7	—	*	*	*	*	*	*	*
736	*9	*	*5	*	*	*	*	*	*	*	*
737	*9	*	*6	*	*	*	*	*	*	*	*
737.3	*8	*	*6	*	*	*	*	*	*	*	*
737.30	*6	*	*5	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
738	13	*5	*8	*	*	*	*	*	*	*	*
738.4	10	*	*6	*	*	*	*	*	*	*	*
740–759	167	92	75	114	25	19	*8	64	28	39	36
742	*7	*	*	*	*	*	*	*	*	*	*
745	25	12	12	20	*	*	*	14	*	*5	*
745.2	*6	*	*	*5	*	*	—	*	*	*	*
745.4	*6	*	*	*5	*	—	—	*	*	*	*
745.5	*6	*	*	*	*	*	*	*	*	*	*
746	15	*8	*7	*9	*	*	*	*5	*	*	*
746.8	*6	*	*	*	*	*	*	*	*	*	*
747	17	*8	9	11	*	*	*	*7	*	*	*
749	10	*6	*	10	*	—	—	*	*	*	*
749.2	*5	*	*	*	*	—	—	*	*	*	*
750	13	10	*	12	—	*	*	*	*5	*	*
750.5	11	*9	*	11	—	—	—	*	*	*	*
751	12	*6	*5	9	*	*	*	*	*	*	*
753	10	*	*	*	*	*	*	*	*	*	*
754	*9	*	*	*8	*	*	—	*	*	*	*
755	*6	*	*	*	*	*	*	*	*	*	*
756	19	12	*7	*8	*	*7	*	*	*	*8	*
756.1	11	*7	*	*	*	*7	*	*	*	*6	*
756.12	*6	*	*	*	*	*	*	*	*	*	*
759	*9	*	*	*	*	*	*	*	*	*	*
760–779	152	90	63	152	—	—	—	26	37	41	49
765	20	12	*8	20	—	—	—	*	*	*7	*7
765.1	17	10	*7	17	—	—	—	*	*	*6	*5
769	14	*8	*5	14	—	—	—	*	*	*	*6
770	32	20	13	32	—	—	—	*6	10	*7	10
770.1	*6	*	*	*6	—	—	—	*	*	*	*
770.6	*7	*	*	*7	—	—	—	*	*	*	*
770.8	15	9	*6	15	—	—	—	*	*	*	*
771	13	*8	*5	13	—	—	—	*	*	*	*
771.8	11	*6	*	11	—	—	—	*	*	*	*
773	*7	*	*	*7	—	—	—	*	*	*	*
773.1	*5	*	*	*5	—	—	—	*	*	*	*
774	37	20	17	37	—	—	—	*5	*9	13	10
774.6	33	19	14	33	—	—	—	*	*7	12	10
778	*8	*5	*	*8	—	—	—	*	*	*	*
778.4	*5	*	*	*5	—	—	—	*	*	*	*
779	*8	*	*	*8	—	—	—	*	*	*	*
780–799	302	140	162	52	114	75	61	69	53	134	46
780	56	24	33	23	17	*7	9	17	10	22	*8
780.2	12	*6	*7	*	*	*	*	*	*	*5	*
780.3	28	10	17	13	10	*	*	9	*	10	*
780.6	10	*	*5	*8	*	*	*	*	*	*	*
784	*8	*	*	*	*	*	*	*	*	*	*
786	85	47	39	*5	32	37	11	17	21	33	14
786.0	*6	*	*	*	*	*	*	*	*	*	*
786.5	78	42	36	*	31	36	11	16	19	31	12

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
786.50	63	33	30	*	23	32	*9	12	17	24	11
786.52	*7	*	*	*	*	*	*	*	*	*	*
786.59	*5	*	*	—	*	*	*	*	*	*	*
788	*8	*	*	*	*	*	*	*	*	*	*
789	44	16	29	9	28	*6	*	10	11	16	*7
789.0	44	15	28	*8	28	*6	*	10	11	16	*6
789.00	10	*	*6	*	*5	*	*	*	*	*	*
789.03	18	*6	12	*	13	*	*	*7	*	*5	*
799	82	37	45	*6	28	17	32	16	*5	51	10
799.9	79	35	44	*5	26	16	31	16	*	49	9
800–999	2,550	1,277	1,273	223	865	485	977	550	572	905	523
800	9	*8	*	*	*	*	*	*	*	*	*
801	25	16	10	*6	12	*	*	*6	*	*8	*6
801.0	*8	*	*	*	*	*	*	*	*	*	*
801.2	*8	*5	*	*	*5	*	*	*	*	*	*
802	43	34	10	*	31	*	*	10	*8	14	11
802.2	17	15	*	*	13	*	*	*	*	*	*6
802.4	*8	*7	*	*	*6	*	*	*	*	*	*
802.6	*6	*	*	*	*	*	*	*	*	*	*
803	*7	*6	*	*	*	*	*	*	*	*	*
805	64	31	33	*	22	*8	33	11	19	20	14
805.0	13	*9	*	*	*7	*	*	*	*	*	*6
805.02	*5	*	*	*	*	*	*	*	*	*	*
805.2	15	*8	*8	*	*5	*	*8	*	*5	*5	*
805.4	31	14	17	*	*8	*	18	*5	10	11	*5
806	*8	*	*	*	*	*	*	*	*	*	*
807	40	22	18	*	*8	12	20	*7	10	15	*8
807.0	33	18	14	*	*7	10	16	*6	*7	12	*7
807.01	*5	*	*	*	*	*	*	*	*	*	*
807.02	*6	*	*	—	*	*	*	*	*	*	*
807.03	*7	*5	*	—	*	*	*	*	*	*	*
807.09	*6	*	*	—	*	*	*	*	*	*	*
808	49	18	32	*	12	*9	28	11	13	19	*7
808.0	10	*5	*5	*	*	*	*	*	*	*	*
808.2	24	*	19	*	*	*	16	*	*6	*9	*
808.4	*9	*7	*	*	*	*	*	*	*	*	*
808.8	*6	*	*6	*	*	*	*6	*	*	*	*
812	63	21	42	14	*9	14	27	13	15	19	17
812.0	29	*5	24	*	*	*9	18	*6	*6	11	*6
812.00	10	*	*9	*	*	*	*	*	*	*6	*
812.01	10	*	*9	*	*	*	*7	*	*	*	*
812.09	*6	*	*	—	*	*	*	*	*	*	*
812.2	9	*	*6	*	*	*	*6	*	*	*	*
812.21	*6	*	*	*	*	*	*	*	*	*	*
812.4	21	9	12	12	*	*	*	*5	*5	*	*7
812.41	13	*5	*7	9	*	*	*	*	*	*	*5
813	55	29	26	11	22	9	13	13	11	18	13
813.0	11	*7	*	*	*6	*	*	*	*	*	*
813.2	*7	*	*	*	*	*	*	*	*	*	*
813.23	*6	*	*	*	*	*	*	*	*	*	*
813.4	20	*7	13	*	*7	*	*7	*	*	*7	*
813.42	11	*	*7	*	*	*	*	*	*	*	*
813.44	*6	*	*	*	*	*	*	*	*	*	*
813.5	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
815	*6	*	*	—	*	*	*	*	*	*	*
816	14	12	*	*	*7	*	*	*6	*	*	*
816.1	10	9	*	*	*	*	*	*	*	*	*
816.11	*5	*5	*	*	*	*	*	*	*	*	*
820	349	80	269	*	*9	22	316	64	91	127	66
820.0	113	25	87	*	*	*8	103	21	35	34	23
820.02	*6	*	*	—	—	*	*6	*	*	*	*
820.03	9	*	*7	*	*	*	*8	*	*	*	*
820.09	91	18	73	—	*	*6	85	19	28	28	16
820.2	161	39	122	*	*	9	146	32	43	59	27
820.20	9	*	*6	*	*	*	*8	*	*	*	*
820.21	139	31	107	*	*	*8	126	26	36	52	24
820.22	13	*	*8	*	*	*	11	*	*	*5	*
820.8	71	14	57	*	*	*	65	11	13	32	16
821	50	25	25	11	15	*6	19	10	13	20	*6
821.0	31	18	13	9	10	*	9	*6	*8	13	*
821.00	*7	*	*	*	*	*	*	*	*	*	*
821.01	24	15	*9	*7	*9	*	*5	*	*7	9	*
821.2	17	*6	12	*	*	*	10	*	*	*7	*
821.23	*8	*	*7	*	*	*	*5	*	*	*	*
822	18	*7	11	*	*6	*	*8	*	*	*7	*
822.0	16	*	11	*	*	*	*7	*	*	*6	*
823	69	41	29	*8	30	15	16	13	16	26	14
823.0	22	11	12	*	*8	*6	*7	*	*6	*7	*
823.00	18	9	9	*	*7	*6	*5	*	*	*6	*
823.2	17	13	*	*	10	*	*	*	*	*7	*
823.20	*6	*	*	*	*	*	*	*	*	*	*
823.22	10	*8	*	*	*6	*	*	*	*	*	*
823.8	20	*9	11	*	*7	*	*6	*	*	*8	*
823.80	*6	*	*	*	*	*	*	*	*	*	*
823.82	12	*6	*6	*	*	*	*	*	*	*	*
823.9	*5	*	*	*	*	*	*	*	*	*	*
824	107	44	63	*	46	32	25	22	24	41	21
824.0	*7	*	*	*	*	*	*	*	*	*	*
824.2	*8	*	*	*	*	*	*	*	*	*	*
824.4	31	*9	22	*	12	*9	9	*7	*5	13	*6
824.6	29	10	19	*	11	11	*7	*6	*7	11	*5
824.8	24	13	11	*	12	*	*6	*5	*5	*8	*5
825	20	12	*7	*	11	*6	*	*5	*	*6	*
825.0	10	*6	*	—	*5	*	*	*	*	*	*
825.2	*7	*	*	*	*	*	*	*	*	*	*
836	10	*7	*	*	*8	*	*	*	*	*	*
840	39	24	16	—	*6	20	14	*8	13	12	*7
840.4	37	22	15	—	*	19	13	*7	12	12	*7
844	26	14	12	*	18	*	*	*8	*6	*8	*
844.2	18	9	*9	*	15	*	*	*	*	*7	*
845	*6	*	*	—	*	*	*	*	*	*	*
845.0	*6	*	*	—	*	*	*	*	*	*	*
846	*8	*	*	—	*	*	*	*	*	*	*
846.0	*6	*	*	—	*	*	*	*	*	*	*
847	15	*9	*7	*	*6	*	*5	*	*	*6	*
847.2	*7	*	*	—	*	*	*	*	*	*	*
850	38	24	13	*7	19	*5	*6	17	*6	9	*6
850.0	12	*8	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
850.1	14	*8	*6	*	*8	*	*	*7	*	*	*
850.5	*9	*6	*	*	*6	*	*	*	*	*	*
851	15	11	*	*	*7	*	*	*	*	*	*5
851.8	*9	*6	*	*	*	*	*	*	*	*	*
852	35	21	15	*	*7	*	22	*5	*6	16	*8
852.2	29	18	12	*	*5	*	19	*	*5	14	*7
852.20	13	*8	*	*	*	*	10	*	*	*8	*
852.21	*9	*	*5	*	*	*	*6	*	*	*	*
853	*6	*	*	*	*	*	*	*	*	*	*
853.0	*6	*	*	*	*	*	*	*	*	*	*
854	47	31	16	*9	21	*6	12	*6	14	13	14
854.0	46	31	15	*9	20	*6	11	*6	14	12	14
854.00	13	*8	*	*	*	*	*5	*	*	*	*
854.01	10	*7	*	*	*	*	*	*	*	*	*
854.02	12	9	*	*	*7	*	*	*	*	*	*
854.06	*7	*	*	*	*	*	*	*	*	*	*
860	27	19	*7	*	15	*8	*	*	*6	12	*
860.0	13	*9	*	*	*7	*	*	*	*	*6	*
861	13	*7	*5	*	*7	*	*	*	*	*6	*
861.2	*8	*	*	*	*	*	*	*	*	*	*
861.21	*8	*	*	*	*	*	*	*	*	*	*
863	12	11	*	*	*8	*	*	*	*	*	*
864	*8	*6	*	*	*6	*	*	*	*	*	*
864.0	*6	*	*	*	*	*	*	*	*	*	*
865	12	*7	*5	*	*7	*	*	*	*	*	*
865.0	10	*6	*	*	*6	*	*	*	*	*	*
871	*9	*8	*	*	*	*	*	*	*	*	*
873	24	16	*8	*	13	*	*5	*	*5	10	*
873.0	*5	*	*	*	*	*	*	*	*	*	*
873.4	12	*7	*	*	*5	*	*	*	*	*5	*
873.42	*7	*	*	*	*	*	*	*	*	*	*
879	10	*8	*	*	*9	*	*	*	*	*	*
879.2	*5	*	*	—	*	*	*	*	*	*	*
881	*8	*6	*	*	*6	*	*	*	*	*	*
882	*6	*	*	*	*	*	*	*	*	*	*
883	*8	*	*	*	*6	*	*	*	*	*	*
883.2	*5	*	*	*	*	*	*	*	*	*	*
886	10	10	*	*	*	*5	*	*	*	*	*
886.0	10	9	*	*	*	*	*	*	*	*	*
891	14	9	*	*	*8	*	*	*	*	*6	*
891.0	*6	*	*	*	*	*	*	*	*	*	*
891.1	*7	*6	*	*	*	*	*	*	*	*	*
892	*5	*	*	*	*	*	*	*	*	*	*
905	*7	*	*	—	*	*	*	*	*	*	*
920	*7	*	*	*	*	*	*	*	*	*	*
921	*5	*	*	*	*	—	*	*	*	*	*
922	17	9	*8	*	*6	*	*7	*	*	*7	*
922.1	*8	*	*	—	*	*	*	*	*	*	*
924	23	10	13	*	*6	*	14	*	*6	*7	*5
924.0	12	*6	*6	—	*	*	9	*	*	*	*
924.01	11	*5	*5	—	*	*	9	*	*	*	*
924.8	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
935	11	*5	*6	*6	*	*	*	*	*	*	*
935.1	10	*	*	*6	*	*	*	*	*	*	*
942	*5	*	*	*	*	*	*	*	*	*	*
944	*5	*	*	*	*	*	*	*	*	*	*
945	*6	*	*	*	*	*	*	*	*	*	*
946	10	*8	*	*	*	*	*	*	*	*7	*
946.2	*6	*	*	*	*	*	*	*	*	*	*
952	*6	*	*	*	*	*	*	*	*	*	*
958	*7	*6	*	*	*	*	*	*	*	*	*
959	15	*7	*8	*	*6	*	*	*	*	*6	*
965	51	16	35	*	37	*6	*	12	10	17	11
965.0	13	*6	*7	*	*9	*	*	*	*	*	*
965.1	10	*	*7	*	*7	*	*	*	*	*	*
965.4	22	*	17	*	17	*	*	*5	*5	*8	*
966	11	*6	*5	*	*5	*	*	*	*	*	*
966.1	*6	*	*	*	*	*	*	*	*	*	*
967	*7	*	*6	*	*	*	*	*	*	*	*
968	*8	*6	*	*	*7	*	*	*	*	*	*
969	57	23	33	*	39	10	*	*9	18	18	11
969.0	19	*8	11	*	12	*	*	*	*6	*6	*
969.4	25	12	14	*	18	*	*	*	10	*7	*
972	10	*	*6	*	*	*	*5	*	*	*	*
972.1	*5	*	*	*	*	*	*	*	*	*	*
977	12	*7	*6	*	10	*	*	*	*	*7	*
977.9	*7	*	*	*	*6	*	*	*	*	*	*
980	*9	*6	*	*	*7	*	*	*	*	*	*
980.0	*7	*5	*	*	*6	*	—	*	*	*	*
987	*6	*	*	*	*	*	*	*	*	*	*
989	*8	*5	*	*	*	*	*	*	*	*	*
989.5	*6	*	*	*	*	*	*	*	*	*	*
991	*6	*	*	*	*	*	*	*	*	*	*
994	*7	*	*	*	*	*	*	*	*	*	*
995	23	11	12	*	*	*7	*7	*5	*7	*8	*
995.1	*5	*	*	*	*	*	*	*	*	*	*
995.2	*8	*	*	*	*	*	*	*	*	*	*
996	358	184	174	27	81	97	154	89	67	127	75
996.0	27	15	12	*	*	*6	18	*8	*5	*8	*5
996.01	19	*9	11	*	*	*	15	*6	*	*6	*
996.1	11	*	*6	*	*	*	*	*	*	*	*
996.2	19	11	*7	11	*	*	*	*6	*5	*	*
996.3	*7	*5	*	*	*	*	*	*	*	*	*
996.39	*5	*	*	*	*	*	*	*	*	*	*
996.4	65	28	37	*	11	17	36	16	14	23	12
996.5	14	*	10	*	*	*	*	*	*	*	*
996.59	10	*	*7	*	*	*	*	*	*	*	*
996.6	87	48	39	*6	22	21	38	19	19	33	15
996.62	39	22	16	*	13	9	13	10	*7	14	*7
996.64	*7	*	*	—	*	*	*6	*	*	*	*
996.66	10	*6	*	*	*	*	*7	*	*	*	*
996.67	10	*	*6	*	*	*	*	*	*	*	*
996.69	13	*7	*6	*	*	*	*	*	*	*	*
996.7	111	57	54	*	25	35	48	27	16	39	29

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
996.73	49	25	24	*	13	16	19	11	*6	18	15
996.74	33	17	16	*	*	*9	18	*8	*	12	*9
996.77	*9	*	*	—	*	*	*	*	*	*	*
996.78	*9	*	*	*	*	*	*	*	*	*	*
996.8	18	11	*7	*	9	*6	*	*6	*	*9	*
996.81	12	*7	*	*	*7	*	*	*	*	*7	*
997	102	52	49	*	24	30	43	19	20	40	22
997.1	11	*5	*6	—	*	*	*6	*	*	*	*
997.2	11	*6	*	—	*	*	*	*	*	*	*
997.3	10	*	*5	*	*	*	*	*	*	*	*
997.4	44	20	24	*	14	11	16	9	9	16	9
997.5	*8	*5	*	*	*	*	*	*	*	*	*
997.6	12	*9	*	*	*	*	*7	*	*	*6	*
997.62	*9	*6	*	*	*	*	*5	*	*	*	*
998	175	83	92	10	54	45	66	42	35	62	35
998.1	37	17	19	*	11	*9	14	11	*8	11	*7
998.2	*8	*	*	*	*	*	*	*	*	*	*
998.3	13	*5	*7	*	*	*	*	*	*	*6	*
998.5	96	46	50	*	27	27	38	21	21	35	19
998.8	17	*7	9	*	*7	*	*	*	*	*5	*
998.89	16	*7	9	*	*7	*	*	*	*	*5	*
999	*8	*	*	*	*	*	*	*	*	*	*
V01–V82	4,246	177	4,069	59	3,919	88	180	740	944	1,458	1,105
V10	*6	*	*	—	*	*	*	*	*	*	*
V22	12	—	12	—	12	—	—	*	*	*	*
V22.2	*7	—	*7	—	*7	—	—	*	*	*	*
V24	*6	—	*6	—	*6	—	—	*	*	*	*
V25	*6	—	*6	—	*6	—	—	*	*	*	*
V25.2	*6	—	*6	—	*6	—	—	*	*	*	*
V27	3,830	—	3,830	10	3,817	*	—	624	785	1,384	1,036
V27.0	3,184	—	3,184	*8	3,173	*	—	539	617	1,145	882
V27.1	18	—	18	*	18	—	—	*	*	*8	*
V27.2	40	—	40	*	40	*	—	9	*6	13	11
V27.9	585	—	585	*	584	*	—	72	158	216	138
V30	*8	*	*	*8	—	—	—	*	*	*	*
V30.1	*8	*	*	*8	—	—	—	*	*	*	*
V45	*8	*	*	*	*	*	*	*	*	*	*
V45.8	*7	*	*	*	*	*	*	*	*	*	*
V54	*6	*	*	*	*	*	*	*	*	*	*
V55	18	10	*7	*	*	*	*6	*6	*5	*	*
V55.3	12	*7	*5	*	*	*	*	*	*	*	*
V57	153	60	93	*	12	24	116	27	86	17	22
V57.1	36	13	23	*	*	*	29	14	12	*	*
V57.8	117	47	70	*	10	19	87	13	74	12	17
V57.89	117	47	70	*	10	19	87	13	74	12	17
V58	112	51	61	21	21	33	36	44	36	19	12
V58.0	*6	*	*	*	*	*	*	*	*	*	*
V58.1	96	45	51	19	19	29	29	39	30	16	10
V58.4	*6	*	*	*	*	*	*	*	*	*	*
V58.49	*6	*	*	*	*	*	*	*	*	*	*
V70	9	*6	*	*	*6	*	*	*	*	*	*
V70.7	*9	*6	*	*	*6	*	*	*	*	*	*
V71	19	11	*8	*	*7	*	*	*7	*	*	*5
V71.4	*7	*	*	*	*	*	*	*	*	*	*
V71.8	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
All codes	159,883	69,928	89,955	10,064	39,645	33,521	76,652	41,343	35,574	57,392	25,574
001-139	6,242	3,205	3,037	594	1,903	1,105	2,640	1,752	1,090	2,226	1,173
003	*36	*	*	*	*	*	*	*	*	*	*
008	374	153	221	103	36	58	178	98	87	126	62
008.4	184	71	113	*	*	*	123	*64	*41	*52	*
008.45	168	67	102	*	*	*	116	*62	*36	*45	*
008.6	50	30	*20	49	*	*	—	*	22	*14	*
008.61	48	29	*19	48	*	*	—	*	22	*14	*
008.8	128	42	87	41	*16	*17	54	29	*21	57	*21
009	39	*	*20	*	*	*	*	*	*	*16	*
009.0	*24	*	*	—	*	*	*	*	*	*	*
011	*85	*	*	*	*	*	*	*	*	*	*
011.9	*62	*	*	*	*	*	*	*	*	*	*
034	26	*11	*14	*	*14	*	*	*	*	*14	*
034.0	22	*	*14	*	*14	*	*	*	*	*	*
038	3,006	1,338	1,668	109	389	524	1,983	702	590	1,226	488
038.0	201	80	120	*	*	*	103	*	*44	*55	*40
038.1	582	288	294	*	*52	168	350	*122	129	238	*93
038.2	75	*	*45	*	*	*	*	*	*	*	*
038.4	821	352	469	*	76	145	586	256	145	278	142
038.40	*64	*	*	*	*	*	*39	*	*	*	*
038.42	363	154	210	*	*29	74	255	118	89	97	59
038.43	*103	*70	*	*	*	*	*69	*	*	*	*
038.49	251	101	150	*	*	*	187	*	*	94	*
038.8	125	*51	*74	*	*	*	88	*	*	*47	*
038.9	1,160	512	648	39	198	129	794	183	227	573	177
041	106	*	*72	*	*	*	*39	*	*	*	*
042	1,201	915	286	*	933	224	*	551	102	298	250
047	66	32	34	28	*26	*	*	*	*	*25	*14
047.9	58	28	30	*26	*23	*	*	*	*	*22	*
053	111	*41	70	*	*	*	59	*	*29	*33	*
053.9	62	*	*43	*	*	*	*31	*	*	*	*
054	*50	*	*	*	*	*	*	*	*	*	*
070	86	*43	44	*	52	*	*	*27	*	*34	*
070.1	*22	*	*	*	*	*	*	*	*	*	*
070.5	*37	*	*	*	*	*	*	*	*	*	*
075	28	*	*	*	*17	*	*	*	*	*	*
078	*50	*	*	*	*	*	*	*	*	*	*
079	168	82	86	96	26	*24	*21	46	28	59	35
079.9	152	74	78	84	25	*23	*20	41	*26	51	34
079.99	152	74	78	84	25	*23	*20	41	*26	51	34
112	167	*63	104	*	*	*	107	*	*	*44	*53
112.4	*54	*	*	*	*	*	*47	*	*	*	*
117	*71	*	*	*	*	*	*	*	*	*	*
135	*44	*	*	—	*	*	*	*	*	*	*
136	109	*72	*	*	*59	*	*	*	*	*53	*
136.3	*83	*62	*	*	*	*	*	*	*	*	*
140-239	11,182	4,947	6,235	309	1,520	3,512	5,841	3,256	2,201	3,796	1,929
141	*42	*	*	—	*	*	*	*	*	*	*
150	98	*64	*	—	*	*	*71	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD–9–CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
151	272	184	*88	—	*	*90	170	*91	*	*75	*63
151.0	*79	*67	*	—	*	*	*	*	*	*	*
151.8	*63	*	*	—	*	*	*49	*	*	*	*
151.9	*60	*	*	—	*	*	*	*	*	*	*
153	1,011	438	573	*	*	238	743	229	227	426	129
153.1	*84	*	*54	*	*	*	*	*	*	*	*
153.2	*41	*	*	—	*	*	*	*	*	*	*
153.3	227	126	102	—	*	*59	160	*	*	96	*44
153.4	214	*67	147	—	*	*	169	*	*63	*73	*
153.6	191	*94	*97	—	*	*	158	*	*43	*83	*
153.8	*67	*	*	—	*	*	*54	*	*	*	*
153.9	109	*43	66	—	*	*	81	*	*	62	*
154	478	187	291	—	*75	*75	329	195	100	117	*66
154.0	160	*86	*74	—	*	*	118	*	*	*	*
154.1	281	86	194	—	*	*	196	*125	*65	*	*42
155	87	*	*57	*	*	*	*	*	*	*	*
155.0	*57	*	*	*	*	*	*	*	*	*	*
156	*77	*	*	—	—	*	*55	*	*	*	*
157	364	213	152	—	*	167	188	*82	*81	142	*59
157.0	118	*68	*50	—	*	*	*71	*	*	*	*
157.8	*83	*	*65	—	*	*	*57	*	*	*	*
157.9	136	114	*	—	*	*92	*40	*	*	*82	*
161	*93	*67	*	—	*	*	*	*	*	*	*
162	1,311	759	552	*	74	431	805	327	277	480	227
162.2	115	59	*56	—	*	*29	84	*	*27	*38	*
162.3	446	243	202	—	*	142	290	114	84	157	91
162.5	221	146	75	—	*	*56	142	*55	*45	82	*39
162.8	154	73	*82	—	*	*48	85	*63	*	*50	*
162.9	340	211	129	*	*	134	192	75	75	148	42
170	*37	*	*	*	*	*	*	*	*	*	*
171	*33	*	*	*	*	*	*	*	*	*	*
172	*26	*	*	—	*	*	*	*	*	*	*
173	*21	*	*	—	*	*	*	*	*	*	*
174	362	—	362	*	54	145	162	104	66	122	69
174.2	*14	—	*14	—	*	*	*	*	*	*	*
174.4	53	—	53	—	*	*17	29	*	*15	*17	*
174.8	91	—	91	—	*	36	40	38	*12	*24	*17
174.9	171	—	171	*	*28	71	71	*40	30	67	35
180	59	—	59	*	*27	*20	*	*	*	*30	*
180.9	*30	—	*30	*	*	*	*	*	*	*18	*
182	156	—	156	—	*	48	93	*42	48	41	*26
182.0	155	—	155	—	*	48	92	*40	48	41	*26
183	169	—	169	*	*	72	77	59	*	50	*
183.0	167	—	167	*	*	70	75	57	*	49	*
185	390	390	—	—	*	135	255	88	103	143	55
188	270	203	67	—	*	61	201	102	47	78	44
188.4	*23	*	*	—	—	*	*	*	*	*	*
188.8	114	87	*	—	*	*	72	*56	*	*	*
188.9	89	71	*	—	—	*	73	*	*	*	*
189	208	123	85	*	*	56	132	*63	*	81	*
189.0	169	99	70	*	*	*41	109	*59	*	61	*
189.1	*30	*	*	—	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD–9–CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
191	210	145	65	*	*58	71	*60	*53	*50	*71	*35
191.1	*57	*	*	*	*	*	*	*	*	*	*
191.2	*49	*	*	—	*	*	*	*	*	*	*
193	44	*	34	*	*16	*12	*	*13	*	*17	*
195	*48	*	*	*	*	*	*	*	*	*	*
196	91	*38	53	—	*	*46	*31	*33	*	*28	*
197	866	313	553	*	*56	251	544	240	184	293	148
197.0	167	70	97	*	*	38	103	40	*39	*67	*
197.2	212	75	137	—	*	67	123	*71	*51	*51	*39
197.6	126	*	94	—	*	*45	73	*42	*	*37	*
197.7	235	101	133	*	*	75	149	64	*37	89	*45
198	951	390	561	*	62	327	544	339	199	282	131
198.3	295	111	183	*	*	119	156	82	83	103	*27
198.5	415	189	226	*	*	123	263	174	83	118	*41
198.8	153	*60	93	*	*	*46	82	*56	*25	*33	*39
198.89	141	*58	83	*	*	*42	77	*53	*22	*	*38
199	87	*	62	*	*	*	*33	*	*	*44	*
199.0	*34	*	*	*	*	*	*	*	*	*	*
199.1	*52	*	*46	—	*	*	*	*	*	*	*
200	153	*78	*75	*	*	*	75	*	*	*45	*43
200.0	95	*53	*	*	*	*	*51	*	*	*	*
201	*86	*	*	*	*	*	*	*	*	*	*
201.9	*68	*	*	*	*	*	*	*	*	*	*
202	262	130	133	*	*	91	132	83	*72	79	*28
202.8	225	105	120	*	*	80	112	*63	*68	67	*27
202.80	116	*53	*63	*	*	*44	59	*	*	*33	*
202.83	*35	*	*	—	*	*	*	*	*	*	*
203	194	94	*100	—	*	*74	100	*48	*	*	*
203.0	189	94	*95	—	*	*69	100	*43	*	*	*
203.00	188	94	*95	—	*	*69	100	*	*	*	*
204	247	109	138	*88	*	*	*83	107	*	*67	*63
204.0	153	*67	*86	*88	*	*	*	*	*	*36	*
204.00	113	*59	*54	*58	*	*	*	*	*	*	*
204.1	92	*	*	*	—	*	*67	*	*	*	*
204.10	92	*	*	—	—	*	*67	*	*	*	*
205	320	168	152	*	*	*107	99	*131	*	*95	*
205.0	219	*114	106	*	*	*62	*88	*	*	*80	*
205.00	201	*105	*96	*	*	*	*83	*	*	*64	*
205.1	*98	*	*	*	*	*	*	*	*	*	*
205.10	*98	*	*	*	*	*	*	*	*	*	*
210	*13	*	*	*	*	*	*	*	*	*	*
210.2	*11	*	*	—	*	*	*	*	*	*	*
211	167	79	88	*	*	*34	116	59	*	44	*
211.3	127	61	66	—	*	*26	97	*38	*	*32	*
218	665	—	665	—	368	278	*19	150	125	247	143
218.0	92	—	92	—	48	43	*	*27	*16	26	*23
218.1	195	—	195	—	106	85	*	64	40	46	45
218.2	67	—	67	—	35	30	*	*	*	38	*
218.9	311	—	311	—	179	120	*	50	59	136	66
220	129	—	129	*	52	56	*	*25	*27	54	*24
225	114	*43	71	*	*	*	*50	*41	*	*	*
225.2	69	*	*51	*	*	*	*34	*	*	*	*
226	16	*	15	*	*8	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD–9–CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
227	78	*27	51	*	*14	*42	*22	*	*	*32	*15
227.1	27	*	*20	—	*	*	*	*	*	*	*
227.3	*37	*	*	*	*	*	*	*	*	*	*
228	*27	*	*	*	*	*	*	*	*	*	*
228.0	*24	*	*	*	*	*	*	*	*	*	*
233	67	*	63	—	24	*25	*18	*	*14	31	*
233.0	*15	—	*15	—	*	*	*	*	*	*	*
233.1	43	—	43	—	20	*	*	*	*	*22	*
235	58	*	*36	*	*	*	*43	*	*	*	*
236	*18	*	*	—	*	*	*	*	*	*	*
237	*35	*	*	*	*	*	*	*	*	*	*
238	140	97	*43	*	*	*	101	*	*	*56	*
238.7	108	83	*	*	*	*	76	*	*	*	*
239	44	*22	*22	*	*	*	*25	*	*	*	*
240–279	6,766	2,903	3,864	454	1,072	1,738	3,503	1,731	1,455	2,488	1,092
241	22	*	*19	*	*	*	*	*	*	*	*
241.1	*16	*	*15	*	*	*	*	*	*	*	*
242	44	*	32	*	*15	*	*	*	*	27	*
242.0	*15	*	*9	—	*	*	*	*	*	*11	*
242.00	*13	*	*	—	*	*	*	*	*	*11	*
242.9	*24	*	*	—	*	*	*	*	*	*	*
244	66	*	56	*	*	*19	*33	*	*	*36	*
244.9	57	*	47	*	*	*16	*28	*	*	*28	*
250	3,175	1,602	1,572	79	573	1,126	1,397	793	663	1,133	585
250.0	624	268	357	*27	94	217	286	144	110	296	74
250.00	162	88	74	*	*	50	95	*23	*24	87	*27
250.01	142	63	79	*	*15	59	58	*34	*22	66	*20
250.02	148	55	94	*	*27	50	70	*	*33	62	*
250.03	172	62	110	*	35	58	64	*52	*30	82	*
250.1	448	249	199	43	197	150	*58	111	80	155	103
250.10	50	*	*19	*	*19	*	*	*	*	*24	*
250.11	126	67	60	*28	74	*	*	45	*22	34	*25
250.12	*36	*	*	—	*	*	*	*	*	*	*
250.13	235	141	94	*14	93	92	*	*48	47	80	*60
250.2	67	*31	*	—	*	*	*	*	*	*23	*
250.3	*64	*	*	*	*	*	*	*	*	*	*
250.4	276	110	166	*	*58	117	97	*38	*35	107	97
250.40	98	*42	*56	—	*	*37	*46	*	*	*	*48
250.41	105	*	*61	*	*	*54	*	*	*	*	*
250.43	*55	*	*	—	*	*	*	*	*	*	*
250.5	*64	*	*	—	*	*	*	*	*	*	*
250.51	*32	*	*	—	*	*	*	*	*	*	*
250.6	217	109	109	—	*45	71	101	*47	*65	56	*
250.60	*75	*	*	—	*	*	*	*	*	*	*
250.61	67	*	*37	—	*	*	*	*	*	*	*
250.63	*49	*	*	—	*	*	*	*	*	*	*
250.7	714	395	319	—	*	313	364	202	184	206	122
250.70	204	133	71	—	*	*78	123	*	*	73	*44
250.71	311	167	144	—	*	164	135	*	*75	*71	*58
250.72	*74	*	*	—	*	*	*57	*	*	*	*
250.73	124	*	*67	—	*	*	*49	*	*	*45	*
250.8	645	360	286	*	77	191	376	183	132	240	90
250.80	218	128	90	—	*	*50	160	*62	*36	75	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
250.81	193	129	65	*	*26	62	104	*59	*48	59	*
250.82	140	*54	86	—	*	*	69	*	*	*64	*
250.83	94	*49	*45	*	*	*	*43	*	*	*41	*
250.9	55	*26	*29	*	*	*	*	*	*	*	*
250.93	*24	*	*	*	*	*	*	*	*	*	*
251	*45	*	*	*	*	*	*	*	*	*	*
251.2	*33	*	*	*	*	*	*	*	*	*	*
252	*23	*	*	*	*	*	*	*	*	*	*
252.0	*21	*	*	—	*	*	*	*	*	*	*
253	*54	*	*	*	*	*	*	*	*	*	*
253.6	*43	*	*	—	*	*	*	*	*	*	*
255	41	*	*	*	*	*	*	*	*	*	*
255.4	*28	*	*	*	*	*	*	*	*	*	*
263	167	*51	116	*	*	*	142	*	*	*49	*
263.9	152	*	*110	*	*	*	134	*	*	*	*
272	62	24	38	*	*12	26	*24	*11	*21	*23	*
272.0	25	*10	*15	*	*	*16	*	*	*	*	*
272.4	31	*	*21	—	*	*	*	*	*	*	*
274	60	*43	*	—	*	*	*29	*	*	*	*
274.0	*46	*	*	—	*	*	*	*	*	*	*
275	77	*27	*50	*	*	*	51	*	*	*	*
275.4	66	*25	*41	*	*	*	*46	*	*	*	*
276	2,602	919	1,683	271	275	411	1,646	658	592	1,041	312
276.0	57	*	*41	*	*	*	*43	*	*	*	*
276.1	301	91	210	*	*	42	230	*43	74	123	60
276.2	46	*	*28	*	*	*	*	*	*	*	*
276.5	1,953	709	1,244	240	215	288	1,210	539	425	775	214
276.6	*23	*	*	*	*	*	*	*	*	*	*
276.7	61	*29	*32	*	*	*	*40	*	*	*	*
276.8	126	*27	100	*	*20	*30	76	*26	*26	62	*12
277	127	*73	*53	*58	*57	*	*	*71	*	*	*
277.0	98	*55	*	*	*50	—	—	*55	*	*	*
277.00	98	*55	*	*	*50	—	—	*55	*	*	*
278	52	*	43	*	32	*15	*	*	*	*	*17
278.0	50	*	41	*	31	*	*	*	*	*	*17
278.00	*17	*	*	—	*	*	*	*	*	*	*
278.01	*32	*	*27	*	*24	*	*	*	*	*	*
280-289	1,647	664	983	174	522	301	651	449	348	577	273
280	170	56	114	*	*28	*33	107	49	26	78	*18
280.0	95	38	57	—	*	*	66	*23	*	51	*
280.9	72	*	55	*	*	*	*40	*25	*	*25	*
281	37	*	*27	*	*	*	*15	*	*	*18	*
281.9	*20	*	*	*	*	*	*	*	*	*	*
282	342	145	197	47	271	*	*	94	102	126	*21
282.6	316	135	181	44	259	*	*	89	90	122	*16
282.62	301	127	174	*34	256	*	*	84	86	116	*
283	*39	*	*	*	*	*	*	*	*	*	*
284	173	66	107	*	*	*50	61	*44	*40	*46	*
284.8	152	*51	101	*	*	*50	*53	*40	*	*38	*
285	262	94	168	*	48	41	164	70	57	80	55
285.1	88	*	66	*	*	*	70	*	*20	*	*
285.9	163	66	97	*	*39	31	85	39	34	55	36

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
286	81	46	*35	*	*	*	*44	*	*	*	*
286.0	*19	*	*	*	*	*	*	*	*	*	*
287	186	57	129	*20	*	*37	85	*42	*23	61	*60
287.3	66	*23	*42	*	*	*	*	*	*	*	*
287.4	*44	*	*	*	*	*	*	*	*	*	*
287.5	67	*	*47	*	*	*	*43	*	*	*33	*
288	281	134	147	44	*40	79	118	81	63	93	*44
288.0	260	128	132	*39	*39	76	106	70	62	88	*40
288.8	*19	*	*	*	*	*	*	*	*	*	*
289	76	40	36	*21	*13	*	*28	*	*	33	*
289.2	20	*10	*10	*10	*10	*	—	*	*	*8	*
290-319	16,490	7,986	8,504	908	8,845	3,361	3,375	5,533	3,850	5,538	1,569
290	764	327	436	—	*	*64	699	140	191	303	*129
290.0	243	*75	169	—	*	*	228	*61	*48	*39	*
290.2	212	*98	114	—	—	*	197	*	*82	106	*
290.20	*96	*	*	—	—	*	*96	*	*	*	—
290.21	116	*	65	—	—	*	101	*	*53	*	*
290.3	94	*	*48	—	—	*	94	*	*	*	*
290.4	193	*100	*94	—	—	*	163	*	*	*113	*
290.40	*59	*	*	—	—	*	*57	*	*	*	*
291	494	365	129	—	270	161	*62	112	163	116	103
291.0	117	101	*	—	46	57	*	*34	*	*45	*
291.8	283	225	59	—	166	91	*	62	129	50	*43
292	339	209	130	*	190	65	82	113	80	113	*33
292.0	128	68	60	—	97	*	*	53	*36	*	*
292.1	*31	*26	*	*	*	*	*	*	*	*	*
292.8	165	106	*59	*	64	*42	*58	*55	*32	*69	*
292.81	81	*	*	*	*	*	*	*	*	*	*
292.84	*48	*40	*	—	*34	*	*	*	*	*	*
293	231	102	129	*	*50	*75	107	120	*46	*40	*25
293.0	89	*38	52	*	*	*	74	*	*	*	*
293.8	136	*64	*73	—	*47	*	*	*82	*	*	*
293.83	110	*53	*	—	*	*	*	*78	*	*	*
294	197	*70	128	*	*	*	176	*	*	116	*
294.8	116	*46	*69	*	*	*	106	*	*	*66	*
295	3,268	1,689	1,579	*	2,046	924	286	1,396	685	848	339
295.3	995	580	415	*	626	278	*90	533	106	246	110
295.30	176	*81	*95	*	125	*	*	*	*	*	*
295.32	229	161	*68	—	125	*83	*	*139	*	*	*
295.34	582	330	252	—	371	158	*	322	*	156	*
295.6	282	131	151	—	143	124	*	*124	*98	*55	*
295.62	*73	*	*	—	*	*	*	*	*	*	*
295.64	179	*87	*93	—	*94	*77	*	*73	*	*	*
295.7	1,484	707	777	*	890	442	142	533	361	386	204
295.70	751	335	417	*	458	236	*	277	177	211	*86
295.74	667	355	312	—	393	180	*94	237	174	*155	*102
295.9	366	213	153	*	286	*56	*	161	*73	116	*
295.90	186	110	*76	*	150	*	*	*	*54	*90	*
295.94	161	*91	*70	—	119	*	*	134	*	*	*
296	5,751	2,075	3,676	356	2,963	1,063	1,369	1,818	1,493	2,062	377
296.2	1,826	818	1,008	163	943	273	447	519	462	738	107
296.20	764	358	406	*61	419	131	154	261	245	210	*48
296.22	193	69	124	*	150	*	*	*	*	151	*
296.23	504	222	282	*	217	*62	185	125	127	223	*
296.24	351	163	189	*	151	*68	*97	109	*75	145	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
296.3	2,089	623	1,466	*112	1,047	361	568	715	547	766	*60
296.30	579	198	381	*	270	108	177	222	187	158	*
296.32	169	*	133	*	133	*	*	*	*53	*97	*
296.33	852	248	604	*	427	145	206	281	182	353	*
296.34	466	130	336	*	202	87	162	197	120	141	*
296.4	555	186	369	*	285	135	*122	233	128	95	*98
296.40	250	*100	150	*	109	*69	*	*	*73	*49	*79
296.44	182	*59	*123	*	*74	*	*	*83	*	*	*
296.5	413	134	279	*	212	124	*	*107	120	156	*30
296.50	214	*56	158	*	94	*82	*	*	*69	*82	*
296.53	*86	*	*61	*	*	*	*	*	*	*	*
296.54	*92	*	*	—	*67	*	*	*	*	*	*
296.6	426	162	264	*	275	*82	*	115	100	167	*44
296.60	226	95	130	*	160	*	*	*	*65	98	*
296.63	*56	*	*	*	*35	*	*	*	*	*	*
296.64	125	*	*85	*	*71	*	*	*	*	*	*
296.7	290	85	205	*	144	*53	*	*72	109	77	*
296.8	99	*39	*60	*	*34	*	*	*	*	*43	*
296.89	*40	*	*	*	*	*	*	*	*	*	*
297	*73	*	*	—	*	*	*	*	*	*	*
298	714	320	394	*	410	112	174	297	81	228	108
298.9	675	307	369	*	389	106	162	283	72	215	105
300	355	139	215	*	164	86	64	118	82	114	*40
300.0	97	43	54	*	35	*19	41	*30	*19	38	*
300.00	50	*22	28	*	*	*	*26	*	*	*19	*
300.01	39	*	*19	*	*23	*	*	*	*	*	*
300.1	52	*	*41	*	*34	*	*	*	*	*	*
300.11	*29	*	*24	*	*	*	*	*	*	*	*
300.4	120	*57	63	*	58	*	*	*37	*34	*46	*
300.9	*38	*	*	*	*	*	*	*	*	*	*
301	121	*87	*34	*	87	*	*	*71	*	*	*
301.8	*39	*	*	—	*33	*	*	*	*	*	*
303	1,417	1,052	365	*	991	318	88	317	373	584	143
303.0	158	119	39	—	91	56	*	39	50	42	*26
303.00	41	32	*	—	*25	*	*	*	*	*21	*
303.01	116	86	*30	—	66	43	*	*36	36	*21	*23
303.9	1,259	934	326	*	900	262	*77	277	322	542	117
303.90	331	262	69	*	202	79	*	*	*29	261	*
303.91	922	666	255	*	695	182	*	251	290	281	99
304	781	521	260	*	628	135	*	252	111	389	*29
304.0	150	100	50	—	120	*27	*	82	*	*42	*
304.00	*40	*	*	—	*	*	—	*	*	*	—
304.01	110	72	*38	—	89	*	*	78	*	*	*
304.2	247	167	80	—	207	*	*	87	*	132	*
304.20	98	*57	*	—	*76	*	*	*	*	*82	*
304.21	148	109	*40	—	130	*	*	77	*	*50	*
304.7	123	83	*40	—	107	*	*	59	*	*	*
304.71	83	49	*34	—	74	*	*	58	*	*	*
304.8	103	72	*31	—	91	*	*	*	*	85	*
304.80	*37	*	*	—	*	*	*	*	*	*	*
304.81	*64	*45	*	—	*61	*	*	*	*	*54	*
304.9	81	*51	*	—	*58	*	*	*	*	*55	*
304.90	74	*51	*	—	*51	*	*	*	*	*	*
305	303	200	103	*	171	84	*38	101	30	148	24
305.0	141	106	*34	*	64	44	*	69	*	55	*
305.00	91	61	*30	*	47	*32	*	*46	*	*33	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
305.01	*48	*44	*	—	*	*	*	*	*	*	*
305.1	32	20	*	—	*12	*16	*	*	*	*21	*
305.6	*30	*	*	—	*25	*	—	*	*	*	*
305.9	70	*47	*	*	*48	*	*	*	*	*45	*
305.90	*50	*	*	*	*	*	*	*	*	*41	*
307	58	*	*51	*	*33	*	*	*	*	*	*
308	*28	*	*25	*	*	*	*	*	*	*	*
309	515	189	326	*63	330	53	*69	198	124	94	99
309.0	149	63	86	*	113	*	*	66	*29	*38	*
309.2	84	*20	*64	*	28	*	*	*26	*	*	*
309.28	35	*	*22	*	*23	*	*	*	*	*	*
309.4	108	54	53	*	71	*	*	57	*	*	*
309.8	123	*	94	*	*79	*	*	*	*38	*	*
309.81	119	*	93	*	*76	*	—	*	*38	*	*
309.9	*41	*	*28	*	*32	*	*	*	*	*	*
310	69	*	*41	*	*	*	*41	*	*	*	*
310.9	*45	*	*	—	*	*	*38	*	*	*	*
311	465	192	273	*55	259	101	*50	188	107	149	*21
312	244	204	*40	*124	110	*	*	147	*46	*	*
312.3	105	80	*	*	72	*	*	*52	*	*	*
312.30	*46	*	*	*	*37	*	—	*35	*	*	—
312.9	*55	*49	*	*	*20	—	—	*	*	*	*
314	153	124	*	125	*	—	*	*	*66	*	*
314.0	152	124	*	124	*	—	*	*	*66	*	—
314.01	127	*103	*	*110	*	—	—	*	*53	*	—
320-389	2,599	1,212	1,388	337	641	565	1,057	588	560	1,026	426
320	*56	*	*	*	*	*	*	*	*	*	*
322	*48	*	*	*	*	*	*	*	*	*	*
322.9	*42	*	*	*	*	*	*	*	*	*	*
323	*31	*	*	*	*	*	*	*	*	*	*
331	269	110	159	*	*	*	196	*56	*43	142	*28
331.0	126	*51	*74	—	—	*	125	*	*	84	*
331.3	*44	*	*	*	*	*	*	*	*	*	*
331.4	86	*29	*57	*	*	*	*	*	*	*	*
332	121	*49	72	*	*	*	110	*	*	*55	*
332.0	118	*48	70	*	*	*	108	*	*	*54	*
333	48	*29	*	*	*	*	*	*	*	*	*
337	*28	*	*	—	*	*	*	*	*	*	*
340	103	*	67	—	41	56	*	*	*39	*36	*
341	*36	*26	*	*	*	—	*	—	*	*	*
342	192	*102	*90	*	*	*75	*94	*	*	111	*
342.9	168	*92	*76	*	*	*	*84	*	*	*93	*
342.90	153	*83	*70	*	*	*	*83	*	*	*83	*
344	162	*119	*	*	*	*	*	*	*	*	*
345	307	157	150	62	87	*47	111	114	50	95	47
345.1	56	*19	*37	*	*	*	*	*	*	*	*
345.10	47	*	*33	*	*	*	*	*	*	*	*
345.3	106	*68	*38	*	*	*	*58	*	*	*33	*
345.4	*37	*	*	*	*	*	*	*	*	*	*
345.5	*26	*	*	*	*	*	*	*	*	*	*
345.9	59	*32	*27	*14	*	*	*	*	*	*	*
345.90	47	*28	*19	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
346	112	*15	97	*	61	40	*	*22	*26	48	*
346.8	*19	*	*	*	*	*	*	*	*	*	*
346.9	69	*	59	*	34	*29	*	*	*	37	*
346.90	35	*	26	*	24	*	*	*	*	*16	*
346.91	*34	*	*33	—	*	*	—	*	*	*	*
348	238	100	138	*	*67	*53	100	*45	*57	89	*47
348.1	*96	*	*	*	*	*	*	*	*	*	*
348.3	74	*38	*	*	*	*	*57	*	*	*	*
348.8	*25	*	*	*	*	*	*	*	*	*	*
349	58	*22	*36	*	*31	*	*	*	*	*37	*
349.0	*18	*	*	*	*14	*	*	*	*	*	*
353	*22	*	*	*	*	*	*	*	*	*	*
355	*34	*	*	*	*	*	*	*	*	*	*
357	*63	*47	*	*	*	*	*	*	*	*	*
357.0	*57	*	*	*	*	*	*	*	*	*	*
359	*31	*	*	*	*	*	*	*	*	*	*
361	20	*8	*13	*	*	*	*13	*	*	*13	*
361.0	16	*	*11	*	*	*	*12	*	*	*11	*
362	*20	*	*11	*	*	*	*	*	*	*	*
366	26	10	16	—	*	*	21	*6	*	*11	*8
366.1	*7	*	*	—	—	*	*6	*	*	*	*
366.9	13	*7	*	—	*	*	11	*	—	*9	*
368	*26	*	*	*	*	*	*	*	*	*	*
376	37	*23	*	*21	*	*	*	*	*	*	*
376.0	*34	*22	*	*21	*	*	*	*	*	*	*
376.01	*34	*22	*	*21	*	*	*	*	*	*	*
381	21	*14	*	*19	*	*	*	*	*	*	*
382	49	28	*21	44	*	*	*	*	*	*24	*
382.9	44	26	*19	41	*	*	*	*	*	*23	*
386	81	28	53	*	*	*18	57	*18	*22	28	*
386.1	*16	*	*	*	*	*	*	*	*	*	*
386.3	51	*21	30	*	*	*	39	*	*	*18	*
386.30	44	*18	26	*	*	*	33	*	*	*15	*
390-459	33,730	16,440	17,290	154	1,792	8,292	23,492	8,937	7,621	12,196	4,975
396	120	*43	*76	*	*	*	88	*	*	*	*
398	*56	*	*	*	*	*	*51	*	*	*	*
398.9	*56	*	*	*	*	*	*51	*	*	*	*
398.91	*54	*	*	—	—	*	*50	*	*	*	*
401	548	207	342	*	72	198	277	125	111	239	73
401.0	98	33	65	—	*24	34	40	*19	*	50	*12
401.9	438	166	272	*	48	157	232	99	94	185	60
402	553	168	385	*	*	93	444	175	112	191	74
402.0	*35	*	*	—	*	*	*	*	*	*	*
402.9	506	154	351	*	*	83	412	163	106	166	70
402.90	47	*	34	—	*	*16	*30	*	*	*19	*
402.91	458	141	317	*	*	67	382	148	99	147	64
403	517	273	243	*	83	188	241	103	92	252	70
403.0	*34	*	*	—	*	*	*	*	*	*	*
403.9	479	259	220	*	74	173	228	98	87	225	69
403.91	459	256	203	*	73	171	212	93	83	214	69

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
404	250	97	153	—	*	80	159	*92	*52	84	*22
404.9	229	90	139	—	*	*76	144	*87	*47	77	*19
404.93	185	73	112	—	*	*59	121	*80	*	*57	*
410	5,166	2,978	2,188	*	185	1,574	3,402	1,368	1,210	1,885	703
410.0	220	135	84	—	*	85	128	*41	74	69	*36
410.01	213	133	80	—	*	83	123	*40	*71	66	*35
410.1	1,010	605	405	—	50	330	630	291	198	377	145
410.11	934	558	376	—	*47	297	590	282	197	338	118
410.12	*45	*	*	—	*	*	*	*	*	*	*
410.2	106	51	55	—	*	*34	68	*31	*	*40	*
410.21	106	51	55	—	*	*33	68	*31	*	*39	*
410.3	139	91	*48	—	*	*41	90	*	*	*43	*
410.31	137	90	*47	—	*	*41	88	*	*	*42	*
410.4	1,011	646	365	*	51	416	541	253	259	387	112
410.41	987	631	356	*	50	409	526	248	257	376	107
410.42	*15	*	*	—	*	*	*	*	*	*	*
410.5	119	65	*54	—	*	*51	67	*43	*	*	*29
410.51	114	60	*54	—	*	*50	63	*41	*	*	*29
410.7	1,808	966	842	*	52	438	1,317	474	435	651	248
410.70	*36	*	*	—	—	*	*	*	*	*	*
410.71	1,717	920	797	*	51	400	1,264	459	424	600	234
410.72	55	*32	*	—	*	*	*29	*	*	*29	*
410.8	163	83	*80	—	*	*	128	*62	*	58	*
410.81	161	82	*79	—	*	*	127	*62	*	58	*
410.9	552	308	244	*	*	132	412	111	131	227	83
410.90	97	61	*36	—	*	*49	48	*	*	*39	*
410.91	436	234	201	*	*	81	349	83	122	172	58
411	926	450	475	*	37	268	618	362	203	269	92
411.1	783	373	410	—	32	226	525	307	170	226	79
411.8	140	76	64	*	*	42	91	54	*32	41	*
411.81	59	*45	*	—	*	*29	*	*	*	*	*
411.89	80	*31	50	*	*	*	64	*32	*	32	*
412	27	*17	*	—	*	*	*	*	*	*	*
413	284	134	149	—	22	109	153	71	71	116	26
413.9	275	129	145	—	*21	107	147	67	70	114	25
414	5,004	2,911	2,092	*	147	1,716	3,134	1,282	1,260	1,714	747
414.0	4,881	2,827	2,054	*	143	1,678	3,054	1,254	1,244	1,649	734
414.00	763	327	436	*	*11	215	535	152	194	322	97
414.01	3,862	2,320	1,542	*	126	1,368	2,364	1,071	947	1,242	603
414.02	255	179	76	—	*	95	154	*31	103	85	*35
414.8	79	*50	*29	*	*	*	*46	*	*	*	*
414.9	29	*25	*	*	*	*	*22	*	*	*17	*
415	496	196	300	*	74	109	313	123	125	138	109
415.1	480	193	287	*	74	106	301	113	122	137	108
415.11	*56	*	*	—	*	*	*	*	*	*	*
415.19	425	173	251	*	58	96	270	104	119	121	81
416	*58	*	*43	*	*	*	*	*	*	*	*
420	75	40	*35	*	*	*	*32	*	*	*	*
420.9	75	40	*35	*	*	*	*32	*	*	*	*
420.90	48	*26	*	*	*	*	*	*	*	*	*
420.91	*20	*	*	*	*	*	*	*	*	*	*
421	242	*129	*113	*	*	*	*115	*114	*	*55	*
421.0	236	*129	*107	*	*	*	*113	*	*	*	*
423	70	*41	*	*	*	*33	*	*	*	*32	*
423.9	47	*21	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD–9–CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
424	606	322	284	*	41	131	425	166	143	198	99
424.0	186	72	114	*	*27	*33	124	*42	*31	65	*47
424.1	405	243	162	*	*	92	298	119	109	131	*46
425	162	99	63	*	*	58	64	*51	*34	65	*
425.4	133	77	56	*	*	51	57	*32	*30	60	*
426	221	110	110	*	*10	26	179	60	53	76	33
426.0	111	64	*47	*	*	*	100	*26	*28	*39	*
426.1	65	*28	*37	*	*	*	53	*	*	*27	*
426.13	*42	*	*	*	*	*	*33	*	*	*	*
427	2,430	1,055	1,374	*17	151	495	1,767	717	517	841	355
427.0	105	41	65	*	*	23	76	*29	*24	42	*
427.1	254	170	84	*	*	81	152	91	74	49	39
427.3	1,174	496	679	*	42	209	922	345	264	420	146
427.31	1,048	436	611	*	30	170	847	310	233	373	131
427.32	127	59	67	*	*	39	75	35	30	46	*16
427.4	96	*44	*	*	*	*58	*	*	*	*	*
427.41	94	*41	*	*	*	*	*	*	*	*	*
427.5	144	*70	*74	*	*	*	71	*	*	*84	*
427.6	56	28	28	*	*	*13	38	*20	*	*13	*14
427.69	46	26	*20	*	*	*	31	*	*	*12	*13
427.8	563	191	371	*	27	79	447	150	124	202	87
427.81	272	82	190	*	*	*13	255	63	51	114	*44
427.89	290	109	181	*	23	66	192	87	73	88	43
427.9	35	*	*20	*	*	*	*26	*	*	*	*
428	5,295	2,157	3,138	*	131	881	4,264	1,391	1,273	1,907	724
428.0	5,256	2,142	3,114	*	131	879	4,231	1,371	1,264	1,901	721
428.1	*31	*	*	*	*	*	*	*	*	*	*
429	85	44	41	*	*	*	64	*	*	32	*
429.2	35	*	*	—	*	*	*26	*	*	*	*
429.4	*24	*	*	*	*	*	*	*	*	*	*
430	302	*61	241	*	*66	183	*52	*	*51	*114	*81
431	605	292	312	*	*	148	411	143	111	241	109
432	156	113	*42	*	*	*	103	*	*	*69	*41
432.1	133	96	*37	*	*	*	91	*	*	*57	*
433	681	356	325	*	*	132	527	184	135	289	73
433.1	447	224	224	*	*	90	340	120	86	194	47
433.10	336	175	161	*	*	66	255	84	70	148	34
433.11	112	*49	63	—	*	*	86	*36	*	*46	*
433.3	177	91	85	—	*	*29	147	57	*39	57	*23
433.30	106	65	41	—	*	*18	87	*29	*21	40	*
433.31	*71	*	*	—	*	*	*60	*	*	*	*
434	2,565	1,174	1,390	*	63	412	2,087	761	494	844	466
434.0	118	*34	85	—	*	*	82	*52	*	*37	*
434.01	105	*31	73	—	*	*	74	*51	*	*	*
434.1	407	194	213	—	*	*75	329	118	68	111	110
434.10	*44	*	*	—	*	*	*41	*	*	*	*
434.11	363	174	189	—	*	*72	288	107	*52	*95	109
434.9	2,039	947	1,092	*	*51	310	1,676	591	401	696	351
434.90	*61	*	*	*	*	*	*	*	*	*	*
434.91	1,979	916	1,063	*	*51	297	1,629	561	395	673	350
435	671	249	422	*	*23	109	538	205	155	236	74
435.3	*31	*15	*	—	*	*	*26	*	*	*	*
435.8	*30	*	*	—	*	*	*	*	*	*	*
435.9	594	220	374	*	*22	96	476	182	143	201	68

* Figure does not meet standard of reliability or precision.
— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
436	1,029	424	605	—	*	172	844	140	191	482	217
437	139	41	98	*	*	*40	85	*	*35	44	*27
437.1	*36	*	*28	*	—	*	*35	*	*	*	*
437.3	*45	*	*	*	*	*	*	*	*	*	*
438	220	*94	*126	—	*	*	183	*	*	*86	*
440	1,089	566	522	*	*	254	809	310	231	437	111
440.2	986	521	465	—	*	221	746	289	208	397	92
440.20	81	53	*28	—	*	*	61	*	*	*46	*
440.21	127	82	*45	—	*	*47	78	*	*	*56	*14
440.22	107	*57	*50	—	*	*	80	*	*	*	*
440.23	142	*67	75	—	*	*39	101	*51	*	*46	*
440.24	517	251	265	—	*	90	419	159	105	207	*
440.3	*48	*	*	—	*	*	*	*	*	*	*
441	568	439	129	*	*	89	467	130	125	233	80
441.0	*109	*	*	—	*	*	*74	*	*	*	*
441.01	*51	*	*	—	*	*	*	*	*	*	*
441.3	97	*82	*	—	*	*	*80	*	*	*	*
441.4	321	258	*63	*	—	*	277	*76	*75	140	*
442	94	55	*39	—	*	*	68	*	*	*34	*
442.3	*56	*31	*	—	*	*	*45	*	*	*	*
443	127	60	68	—	*	*	99	*31	*	66	*
443.9	110	*52	58	—	*	*	91	*	*	61	*
444	327	162	165	*	*44	69	214	99	73	113	41
444.2	262	127	136	—	*	*47	185	78	61	93	*30
444.21	*35	*	*25	—	*	*	*	*	*	*	*
444.22	227	116	111	—	*	*40	159	*70	*51	82	*
444.8	*29	*	*	*	*	*	*	*	*	*	*
444.81	*26	*	*	*	*	*	*	*	*	*	*
446	67	*	*39	*	*	*	*	*	*	*	*
447	84	*33	*51	*	*	*30	*42	*23	*	*37	*
447.1	38	*15	*	*	*	*	*25	*	*	*	*
451	280	115	165	*	*47	72	161	74	84	93	*29
451.0	54	*	*33	—	*	*	*	*	*34	*	*
451.1	163	69	94	*	*	*39	93	*50	*39	*56	*
451.11	*52	*	*35	*	*	*	*	*	*	*	*
451.19	111	*52	*59	—	*	*	69	*	*	*37	*
451.2	*41	*	*	—	*	*	*	*	*	*	*
453	937	414	523	*	151	247	535	187	262	353	135
453.8	871	379	492	*	122	221	524	177	246	317	131
454	109	*28	81	*	*	*	76	*	*36	*	*
454.2	*81	*	*61	—	*	*	*64	*	*	*	*
454.9	*12	*	*	—	*	*	*	*	*	*	*
455	74	41	34	—	24	*23	*27	*14	*18	37	*
455.2	40	*19	*20	—	*	*	*22	*	*	*	*
456	*22	*	*	—	*	*	*	*	*	*	*
458	220	107	113	*	*	44	167	55	53	64	49
458.0	115	55	60	—	*	*20	91	*28	29	36	*21
458.9	99	48	51	*	*	*	71	*24	*22	*26	*27
459	71	*29	*42	*	*	*	*48	*	*	*	*
460-519	19,153	9,125	10,029	2,208	2,156	3,545	11,244	4,412	4,528	7,365	2,848
461	28	*	*	*	*	*	*	*	*	*	*
462	26	*	*13	*	*15	*	*	*	*	*16	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD–9–CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
463	34	*	*20	*	*18	*	*	*	*	*16	*
464	53	36	*16	34	*	*	*	*	*11	*16	*10
464.4	29	19	*10	29	*	—	—	*	*8	*12	*
465	93	42	51	59	*	*	*20	*27	*16	32	*19
465.9	88	41	48	55	*	*	*20	*23	*15	32	*19
466	824	384	440	476	53	83	211	192	190	293	149
466.0	387	135	252	47	53	79	207	97	88	167	35
466.1	437	249	188	428	*	*	*	95	102	126	114
470	*9	*	*	—	*	*	*	*	*	*	*
473	43	*19	*24	*	*17	*	*	*	*	*15	*
474	43	21	22	35	*6	*	—	24	*	*8	*
474.0	14	*6	*9	*9	*	*	—	*7	*	*	*
474.1	28	*15	*	26	*	*	—	17	*	*	*
474.10	24	*11	*	*23	*	—	—	*13	*	*	*
475	46	*	*38	*	*36	*	*	*	*	*	*
478	114	59	55	*36	49	*17	*	*29	*	55	*15
478.2	35	*22	*	*	*	*	*	*	*	*	*
478.29	*23	*17	*	*	*	*	*	*	*	*	*
478.7	*33	*	*	*	*	*	*	*	*	*	*
480	218	124	95	176	*	*	*	*55	47	54	63
480.1	110	55	55	110	*	—	—	*	*29	*36	*30
480.9	79	41	*38	38	*	*	*	*	*	*18	*29
481	362	201	161	*	76	96	176	77	93	125	67
482	2,459	1,319	1,140	76	171	329	1,882	348	654	1,147	310
482.0	210	113	97	*	*	*	166	*	*	112	*
482.1	523	289	235	*	*56	*80	351	*76	126	252	*70
482.2	176	71	105	*	*	*	124	*	69	*37	*39
482.3	104	*51	*52	*	*	*	84	*	*	*45	*
482.30	*41	*	*	*	*	*	*	*	*	*	*
482.39	*39	*	*	*	*	*	*	*	*	*	*
482.4	428	224	204	*	*	*	344	*68	*63	231	*66
482.8	907	507	400	*	*37	110	743	96	296	431	84
482.83	580	346	234	*	*	*49	493	*45	248	244	*
482.89	270	141	129	*	*	*	207	*	*41	157	*
482.9	111	*63	*48	*	*	*	69	*35	*	*39	*
483	*46	*	*	*	*	*	*	*	*	*	*
483.0	*38	*	*	*	*	*	*	*	*	*	*
485	222	72	150	60	*	*	105	*	81	91	*
486	4,452	2,045	2,407	388	497	676	2,890	1,161	1,039	1,629	622
487	73	26	47	*	*11	*	*37	*	*	29	*
487.0	*25	*	*	*	*	*	*	*	*	*	*
487.1	45	*18	*27	*	*	*	*27	*	*	*21	*
490	56	*25	31	*17	*	*	*22	*	*	*23	*
491	2,262	945	1,317	*	84	549	1,612	531	589	896	246
491.2	2,248	937	1,311	*	83	548	1,606	527	587	890	244
491.20	87	*30	58	*	*	*21	*55	*	*	*38	*
491.21	2,161	907	1,254	*	73	527	1,551	506	567	852	236
492	191	101	90	—	*	62	127	*56	*47	45	*44
492.8	146	68	78	—	*	53	93	*48	*38	*38	*
493	1,710	618	1,093	521	471	386	332	486	327	606	291
493.0	85	44	41	69	*13	*	*	24	*	*	43

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD–9–CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
493.00	31	*14	*17	24	*	*	*	*	*	*	*16
493.01	55	30	25	44	*	*	—	*19	*	*	26
493.2	401	93	308	*	66	142	192	122	65	141	73
493.20	261	62	199	*	*35	97	127	57	44	113	*47
493.21	140	*30	110	—	*30	*45	64	*65	*21	*28	*
493.9	1,217	480	738	451	389	239	139	340	249	455	173
493.90	663	231	431	173	242	130	117	161	128	275	99
493.91	555	248	306	277	146	110	*	178	121	180	75
494	*50	*	*	*	*	*	*	*	*	*	*
496	730	344	386	*	*	219	502	186	117	329	98
507	1,493	854	638	*29	115	142	1,207	319	261	624	289
507.0	1,481	852	629	*	115	141	1,201	318	255	619	289
510	171	*114	*	*	*	*	*79	*	*	*	*
510.9	143	*106	*	*	*	*	*	*	*	*	*
511	509	243	266	*	*44	130	331	98	124	163	125
511.0	35	*	*20	—	*	*	*18	*	*	*	*
511.8	118	*	*	*	*	*	*62	*	*	*	*
511.9	335	146	189	*	*35	61	239	73	95	112	55
512	263	165	98	*	78	95	88	*52	*62	116	*33
512.1	*30	*	*	*	*	*	*	*	*	*	*
512.8	207	128	79	*	73	*71	*62	*41	*35	102	*
513	*64	*54	*	*	*	*	*	*	*	*	*
513.0	*51	*	*	*	*	*	*	*	*	*	*
515	99	40	59	*	*	*	59	*	*23	*48	*
516	97	*	*63	*	*	*	*65	*	*	*	*
516.3	*33	*	*	*	*	*	*	*	*	*	*
516.8	*63	*	*	*	*	*	*	*	*	*	*
518	2,141	1,063	1,077	*119	185	569	1,267	434	656	732	319
518.0	*40	*	*	*	*	*	*	*	*	*	*
518.4	45	*	*30	*	*	*	*35	*	*	*	*
518.8	1,953	1,003	950	*101	164	519	1,169	382	637	643	290
518.81	1,836	944	891	*	147	483	1,132	355	599	622	260
518.82	*86	*	*	*	*	*	*	*	*	*	*
518.89	*32	*	*	*	*	*	*	*	*	*	*
519	97	*48	*49	*53	*	*	*	*	*	*	*
519.1	*48	*	*	*	*	*	*	*	*	*	*
520–579	14,745	6,716	8,029	707	3,050	3,603	7,386	3,737	3,407	5,265	2,336
524	35	*16	19	*	29	*	*	*	*9	*	*
524.1	*16	*	*	*	*14	*	*	*	*	*	*
524.10	*13	*	*	*	*10	*	*	*	*	*	*
527	58	*	*43	*	*	*	*45	*	*	*	*
527.2	*42	*	*29	*	*	*	*	*	*	*	*
528	80	*50	*30	*	*	*	*	*	*	*	*
530	668	322	346	90	99	196	283	155	137	289	87
530.1	143	68	75	*	36	42	60	45	*29	48	*22
530.10	51	*25	26	*	*	*20	*	*	*	*20	*
530.11	63	30	33	*	*14	*19	*27	*	*	*23	*
530.19	*30	*	*	*	*	*	*	*	*	*	*
530.2	80	51	*29	*	*	*	48	*	*	*22	*
530.3	36	*	*	*	*	*	*29	*	*	*	*
530.5	31	*	*21	*	*	*18	*	*	*	*17	*
530.7	58	38	*	*	*	*19	*30	*	*	*24	*
530.8	268	115	153	82	35	91	61	48	52	134	34
530.81	223	98	125	81	28	63	50	41	45	103	34

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
531	578	273	305	*	71	153	349	114	148	167	149
531.0	140	61	79	*	*	*29	100	*	45	*37	*35
531.00	129	60	69	*	*	*28	91	*	*40	*37	*34
531.4	232	115	117	*	*	44	150	*41	54	92	46
531.40	230	114	116	*	*	44	148	*39	53	92	45
531.5	*86	*	*	—	*	*	*	*	*	*	*
531.50	*77	*	*	—	*	*	*	*	*	*	*
531.9	*66	*	*	*	*	*	*	*	*	*	*
531.90	*62	*	*	*	*	*	*	*	*	*	*
532	451	260	191	*	67	91	293	123	94	161	73
532.0	93	58	*35	—	*	*30	53	*	*20	*41	*
532.00	91	57	*34	—	*	*30	51	*	*20	*41	*
532.4	180	106	74	*	*21	*30	128	*29	*33	74	*43
532.40	177	103	74	*	*21	*28	128	*29	*33	74	*41
532.9	*24	*	*	—	*	*	*	*	*	*	*
532.90	*20	*	*	—	*	*	*	*	*	*	*
533	94	30	65	*	*	*22	61	*	*27	42	*
533.4	*30	*	*22	*	*	*	*	*	*	*	*
533.40	*29	*	*21	*	*	*	*	*	*	*	*
533.9	*21	*	*	—	*	*	*	*	*	*	*
533.90	*18	*	*	—	*	*	*	*	*	*	*
535	712	334	378	*	117	151	434	193	122	319	78
535.0	158	67	91	*	25	30	100	*24	40	79	*15
535.00	84	33	51	*	*18	*18	46	*16	*19	42	*
535.01	74	*34	*40	*	*	*	54	*	*	*37	*
535.1	55	*	*39	*	*	*	*34	*	*	*33	*
535.10	*23	*	*	*	*	*	*	*	*	*	*
535.11	*32	*	*	*	*	*	*	*	*	*	*
535.3	31	*24	*	*	*	*	*	*	*	*	*
535.31	*18	*	*	*	*	*	*	*	*	*	*
535.4	243	116	127	*	33	*30	176	*90	*32	90	30
535.40	94	*20	74	*	*22	*20	49	*	*	48	*
535.41	149	96	52	*	*	*	127	*68	*	43	*20
535.5	194	96	98	*	35	64	93	*42	32	97	*23
535.50	101	37	65	*	*24	35	42	*	*21	52	*
535.51	92	59	*33	*	*	*	*52	*	*	*45	*
535.6	*31	*	*	*	*	*	*	*	*	*	*
536	95	32	63	*	*36	*23	*27	*23	*	49	*
536.2	*29	*	*23	*	*	*	*	*	*	*	*
536.3	*37	*	*	*	*	*	*	*	*	*	*
536.8	22	*	*	*	*	*	*	*	*	*	*
537	149	*92	*57	*	*	*	*76	*	*	*51	*
537.0	88	*63	*	*	*	*	*53	*	*	*	*
540	784	473	310	201	350	125	109	147	164	263	209
540.0	250	141	109	79	84	*40	*47	48	55	72	76
540.1	180	112	68	*	62	*43	*39	*	*36	80	*40
540.9	353	220	133	87	203	41	*	74	74	112	93
541	*17	*	*10	*	*15	*	*	*	*	*	*
550	151	123	*29	16	*17	26	93	55	29	47	*20
550.1	73	60	*	*	*	*	56	*24	*18	*22	*
550.10	55	46	*	*	*	*	*40	*21	*	*	*
550.9	69	58	*	*9	*12	*18	30	25	*	23	*11
550.90	39	32	*	*	*	*12	*16	16	*	*13	*
550.91	*10	*	*	*	*	*	*	*	*	*	*
550.92	*17	*16	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
552	209	69	140	*	*29	48	132	46	64	66	*33
552.1	32	*	*20	*	*	*	*	*	*	*	*
552.2	105	*30	75	*	*	*23	66	*	*35	*37	*
552.20	*29	*	*	—	*	*	*	*	*	*	*
552.21	69	*	*50	*	*	*	*48	*	*	*	*
553	275	84	191	*	38	76	155	99	52	106	*19
553.1	*12	*	*	*	*	*	*	*	*	*	*
553.2	156	54	102	*	*21	51	83	47	35	64	*11
553.20	48	*	*36	*	*	*	*	*	*	*	*
553.21	100	42	58	*	*	41	47	27	*29	38	*
553.3	104	*25	79	*	*13	*20	67	*	*17	38	*
555	254	87	167	*	133	75	*	93	*45	78	*37
555.0	69	*	*55	—	*	*	*	*	*	*	*
555.1	*64	*	*37	*	*	*	*	*	*	*	*
555.2	*58	*	*	*	*	*	*	*	*	*	*
555.9	62	*	*27	*	*46	*	*	*	*	*	*
556	238	138	99	*	138	*53	*41	*88	*54	*62	*34
556.9	109	*57	*52	*	*57	*	*	*	*33	*49	*
557	355	140	215	*	*	*64	268	*88	107	80	*80
557.0	233	*88	146	*	*	*	167	*49	*61	*58	*
557.1	*70	*	*	—	*	*	*64	*	*	*	*
557.9	*52	*	*30	—	*	*	*37	*	*	*	*
558	571	253	318	130	122	84	236	133	167	223	48
558.9	549	244	305	130	120	77	223	127	162	217	43
560	1,807	737	1,070	*40	184	392	1,191	423	429	630	326
560.1	263	102	160	*	*	41	192	*44	*71	93	56
560.2	187	*99	*88	*	*	*	152	*	*	*106	*
560.3	144	49	95	*	*	*	116	*53	*25	*42	*
560.39	132	48	84	*	*	*	106	*44	*25	*38	*
560.8	756	284	472	*	90	203	455	164	215	227	151
560.81	588	214	374	*	76	169	335	131	151	171	136
560.89	168	*70	98	*	*	*33	120	*33	*64	*56	*
560.9	447	197	251	*	54	109	276	114	89	161	82
562	1,283	541	742	*	135	300	844	379	278	464	162
562.1	1,273	535	738	*	135	297	837	378	274	460	161
562.10	116	51	66	—	*	*39	70	*34	*28	44	*
562.11	792	353	439	*	124	231	433	244	162	277	109
562.12	234	94	140	—	*	*	216	*53	53	99	*29
562.13	131	*38	93	—	*	*	119	*	*	*39	*
564	228	101	127	*31	45	*23	129	45	53	105	25
564.0	83	38	45	*	*	*	45	*	*28	*23	*12
564.1	55	*	51	*	*27	*	*	*	*	*29	*
565	*23	*	*	*	*	*	*	*	*	*	*
566	80	49	*	*	*24	*25	*	*	*	*28	*
567	127	*58	69	*	*	*54	*36	*	*45	*	*
567.2	93	*	*52	*	*	*	*	*	*	*	*
568	68	*	50	*	*32	*	*	*	*	*	*
568.0	60	*	43	*	*29	*	*	*	*	*	*
569	515	187	327	*	66	116	322	142	139	174	59
569.3	60	*26	34	*	*	*	39	*	*19	*	*
569.4	*33	*	*	*	*	*	*	*	*	*	*
569.6	79	*34	*45	*	*	*	*45	*	*	*	*
569.69	61	*	*39	*	*	*	*39	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
569.8	282	88	194	*	*	*	196	*	*79	103	*31
569.83	*106	*	*	—	*	*	*77	*	*	*	*
569.85	59	*	*35	—	*	*	*40	*	*	*	*
570	*76	*	*	*	*	*	*	*	*	*	*
571	515	321	194	*	133	221	150	125	79	222	89
571.1	66	*55	*	—	*31	*33	*	*	*	*	*
571.2	242	174	*68	—	78	117	*46	*66	*37	90	49
571.3	*20	*	*	—	*	*	*	*	*	*	*
571.5	164	63	101	*	*	*54	87	*	*	87	*
572	200	96	104	*	*	95	65	*57	*	72	*38
572.2	100	47	52	*	*	*37	*45	*	*	*39	*
573	49	*	*28	*	*	*	*	*	*	*	*
574	1,546	602	945	*	326	413	799	390	301	612	244
574.0	704	335	369	*	115	183	404	183	112	297	111
574.00	614	295	319	*	109	161	342	154	99	266	95
574.01	90	*40	*50	—	*	*22	*62	*	*	*31	*
574.1	537	160	377	*	138	173	224	137	124	203	73
574.10	514	154	360	*	132	170	211	132	119	192	71
574.11	*23	*	*	*	*	*	*	*	*	*	*
574.2	106	31	75	*	24	25	51	*23	23	33	*28
574.20	99	*27	72	*	22	25	47	*19	20	32	*28
574.3	*53	*	*29	—	*	*	*	*	*	*	*
574.4	55	*	*45	*	*	*	*	*	*	*	*
574.40	*22	*	*	*	*	*	*	*	*	*	*
574.5	91	*40	51	—	*	*	64	*	*	48	*
574.50	37	*	*24	—	*	*	*25	*	*	*	*
574.51	54	*	*	—	*	*	*38	*	*	*30	*
575	265	129	137	*	71	71	122	49	86	100	*30
575.0	156	89	67	—	33	*44	80	*	53	58	*20
575.1	81	*36	45	*	27	*24	*30	*	*22	37	*
576	159	76	83	*	*20	*45	75	*	*30	67	*27
576.1	61	*35	*26	*	*	*25	*	*	*	*	*
576.2	59	*	*39	*	*	*	*31	*	*	*	*
577	1,220	577	643	*	446	377	381	276	388	372	184
577.0	920	424	497	*	343	257	310	196	288	296	140
577.1	207	112	94	*	88	75	*38	*45	85	*47	*29
577.2	*66	*	*	*	*	*	*	*	*	*	*
578	636	253	383	*	39	124	470	179	163	188	106
578.0	66	*24	*42	*	*	*	*58	*	*	*	*
578.1	97	46	51	*	*	*	75	*	*	*30	*22
578.9	473	183	291	*	*31	104	337	129	123	150	71
579	*40	*	*	*	*	*	*	*	*	*	*
580-629	6,866	2,330	4,536	285	1,755	1,489	3,337	1,575	1,451	2,789	1,050
581	*43	*	*	*	*	*	*	*	*	*	*
581.9	*32	*	*	*	*	*	*	*	*	*	*
584	793	337	456	*	75	152	553	192	146	375	80
584.9	725	300	425	*	69	139	513	168	131	355	70
585	267	132	135	*	*48	58	156	86	*44	84	*54
586	69	*	*36	*	*	*	*43	*	*	*	*
590	506	123	383	63	190	110	143	110	124	193	80
590.1	324	67	256	40	130	52	103	77	83	107	56
590.10	323	67	256	40	129	52	103	77	83	107	56
590.8	159	48	111	*21	54	50	*34	*30	*35	74	*19
590.80	159	48	111	*21	54	50	*34	*30	*35	74	*19

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD–9–CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
591	49	*19	*30	*	*16	*	*	*	*	*	*
592	406	233	172	*	180	142	77	82	96	172	56
592.0	132	58	74	*	58	46	*21	29	23	60	*19
592.1	270	173	97	*	119	94	55	52	71	111	36
593	158	73	86	*25	*29	46	58	42	51	41	*24
593.4	56	*20	*35	*	*	*	*	*	*	*	*
593.7	*27	*	*	*	*	*	—	*	*	*	*
593.70	*24	*	*	*	*	*	—	*	*	*	*
593.8	*24	*	*	*	*	*	*	*	*	*	*
593.9	*33	*20	*	*	*	*	*21	*	*	*	*
594	*23	*20	*	*	*	*	*	*	*	*	*
594.1	*20	*	*	—	*	*	*	*	*	*	*
595	89	40	48	*	*	*19	60	*23	*24	*33	*
595.0	37	*	*23	*	*	*	*26	*	*14	*	*
596	114	93	*	*	*	*	92	*	*	*	*
596.0	*28	*28	*	*	*	*	*25	*	*	*	*
598	*20	*17	*	*	*	*	*	*	*	*	*
598.9	*14	*	*	*	*	*	*	*	*	*	*
599	1,918	646	1,272	109	133	253	1,423	473	353	831	261
599.0	1,788	550	1,238	107	116	241	1,324	454	325	774	235
599.7	90	62	28	*	*15	*	67	*15	*22	40	*
600	372	372	—	—	*	60	311	90	116	118	49
601	*22	*22	—	—	*	*	*	*	*	*	*
604	42	42	—	*	*15	*	*	*	*	*	*
604.9	35	35	—	*	*	*	*	*	*	*	*
604.90	*18	*18	—	*	*	*	*	*	*	*	*
607	*23	*23	—	*	*	*	*	*	*	*	*
607.8	*12	*12	—	*	*	*	*	*	*	*	*
607.84	*9	*9	—	—	*	*	*	*	*	*	*
608	*30	*30	—	*	*	*	*	*	*	*	*
611	65	*	64	*	36	*14	*	16	*	*15	*19
611.1	22	*	21	*	16	*	*	*10	*	*	*
614	251	—	251	*	215	*27	*	55	36	125	35
614.0	*29	—	*29	*	*27	*	*	*	*	*	*
614.1	*17	—	*17	*	*11	*	*	*	*	*	*
614.2	*25	—	*25	—	*	*	—	*	*	*	*
614.3	*32	—	*32	*	*29	*	—	*	*	*	*
614.6	88	—	88	—	80	*	*	*	*	*58	*
614.9	44	—	44	*	40	*	—	*	*	*20	*
615	*18	—	*18	—	*15	*	—	*	*	*	*
616	*24	—	*24	*	*11	*	*	*	*	*	*
617	244	—	244	*	187	55	*	39	58	109	37
617.0	136	—	136	—	92	43	*	*25	33	63	*16
617.1	40	—	40	*	35	*	—	*	*	*	*
617.3	46	—	46	—	39	*	*	*	*15	*20	*
618	419	—	419	*	89	155	173	69	95	171	83
618.0	102	—	102	*	*10	42	49	*16	*23	41	22
618.1	28	—	28	—	*12	*	*	*	*	*	*
618.2	53	—	53	—	*19	*24	*	*	*15	*15	*
618.3	65	—	65	—	*	*17	40	*18	*18	*17	*
618.4	51	—	51	—	*20	*15	*	*	*	*25	*
618.5	62	—	62	—	*	27	33	*	*	33	*13
618.8	43	—	43	—	*17	*14	*	*	*	*28	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
620	218	—	218	*	141	54	*	52	33	91	42
620.0	*13	—	*13	*	*	*	—	*	*	*	*
620.1	61	—	61	*	42	*	*	*	*	30	*
620.2	121	—	121	*	77	*31	*	33	*14	46	*28
621	38	—	38	—	*	*20	*	*14	*	*13	*
621.0	*12	—	*12	—	*	*	*	*	*	*	*
621.3	*18	—	*18	—	*	*12	*	*	*	*	*
622	31	—	31	—	*23	*	*	*	*	*18	*
622.1	30	—	30	—	*22	*	*	*	*	*17	*
625	236	—	236	*	125	76	35	32	56	96	53
625.3	26	—	26	*	*19	*	—	*	*	*	*
625.6	132	—	132	—	35	63	34	*17	33	44	38
625.9	60	—	60	*	55	*	*	*	*	38	*
626	206	—	206	*	123	79	*	*22	52	97	36
626.2	143	—	143	—	91	53	—	*18	31	71	*23
626.8	54	—	54	*	28	*22	*	*	*16	22	*
627	29	—	29	—	*	*14	*	*	*	*14	*
627.1	*24	—	*24	—	*	*	*	*	*	*	*
630-677	1,338	—	1,338	*	1,323	*	—	310	262	519	247
632	25	—	25	*	25	—	—	*	*	*12	*
633	86	—	86	—	86	—	—	*16	*19	38	*13
633.1	79	—	79	—	79	—	—	*15	*17	35	*12
634	39	—	39	*	39	—	—	16	*	12	*
634.9	28	—	28	*	28	—	—	14	*	*8	*
634.91	20	—	20	*	20	—	—	*9	*	*	*
635	21	—	21	*	20	*	—	*	*	*	*
635.9	*15	—	*15	—	*14	*	—	*	*	*	*
637	*11	—	*11	—	*11	*	—	*	*	*	*
640	*16	—	*16	—	*16	—	—	*	*	*	*
640.0	*15	—	*15	—	*15	—	—	*	*	*	*
640.03	*15	—	*15	—	*15	—	—	*	*	*	*
641	51	—	51	—	51	—	—	*15	*14	*	*
641.1	*20	—	*20	—	*20	—	—	*	*	*	*
641.13	*20	—	*20	—	*20	—	—	*	*	*	*
641.2	*16	—	*16	—	*16	—	—	*	*	*	*
641.23	*16	—	*16	—	*16	—	—	*	*	*	*
642	46	—	46	*	45	*	—	*12	*9	*17	*
642.3	*15	—	*15	—	*15	—	—	*	*	*	*
642.33	*15	—	*15	—	*15	—	—	*	*	*	*
642.4	*11	—	*11	—	*11	*	—	*	*	*	*
642.43	*10	—	*10	—	*9	*	—	*	*	*	*
643	88	—	88	—	88	—	—	*23	*18	36	*
643.0	26	—	26	—	26	—	—	*	*	*12	*
643.03	26	—	26	—	26	—	—	*	*	*12	*
643.1	56	—	56	—	56	—	—	*	*	*21	*
643.13	56	—	56	—	56	—	—	*	*	*21	*
644	315	—	315	*	315	*	—	57	65	121	72
644.0	267	—	267	*	267	*	—	51	59	92	65
644.03	267	—	267	*	267	*	—	51	59	92	65
644.1	48	—	48	—	48	—	—	*6	*6	29	*
644.13	48	—	48	—	48	—	—	*6	*6	29	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
646	147	—	147	*	146	—	—	22	23	79	22
646.6	111	—	111	*	111	—	—	*17	*15	59	*21
646.63	105	—	105	*	105	—	—	*16	*14	56	*21
646.8	28	—	28	—	28	—	—	*	*	*17	*
646.83	26	—	26	—	26	—	—	*	*	*17	*
647	*23	—	*23	—	*23	*	—	*	*	*	*
648	253	—	253	*	249	*	—	73	48	101	31
648.0	33	—	33	—	33	—	—	*	*	*	*
648.03	33	—	33	—	33	—	—	*	*	*	*
648.4	*35	—	*35	*	*33	—	—	*	*	*	*
648.43	*28	—	*28	*	*27	—	—	*	*	*	*
648.8	37	—	37	—	37	—	—	*	*	*18	*
648.83	36	—	36	—	36	—	—	*	*	*17	*
648.9	94	—	94	*	92	*	—	21	*18	41	*15
648.93	90	—	90	*	90	*	—	21	*18	38	*13
654	*27	—	*27	—	*17	*	—	*	*	*	*
658	52	—	52	—	52	—	—	*	*	*	*
659	10	—	10	—	10	—	—	*	*	*	*
659.8	*6	—	*6	—	*6	—	—	*	*	*	*
659.83	*6	—	*6	—	*6	—	—	*	*	*	*
670	*20	—	*20	—	*20	—	—	*	*	*	*
670.0	*20	—	*20	—	*20	—	—	*	*	*	*
670.04	*20	—	*20	—	*20	—	—	*	*	*	*
674	*17	—	*17	—	*17	—	—	*	*	*	*
680-709	2,844	1,432	1,412	170	640	782	1,252	760	629	969	486
681	93	61	*33	*	*21	*30	*34	*	*31	*34	*
681.0	45	*30	*	*	*19	*	*	*	*	*	*
681.00	41	*26	*	*	*	*	*	*	*	*	*
681.1	48	*31	*	*	*	*	*	*	*	*	*
681.10	47	*30	*	*	*	*	*	*	*	*	*
682	1,731	855	876	85	410	530	706	481	398	587	266
682.0	76	40	*36	*21	*23	*23	*	*	*	*29	*15
682.1	*33	*	*	*	*	*	*	*	*	*	*
682.2	116	59	58	*	*31	*27	*53	*	*21	*46	*31
682.3	177	94	83	*	76	52	*39	64	31	49	*34
682.4	77	50	*27	*	*24	*	*29	*28	*	*26	*
682.5	*32	*	*	*	*	*	*	*	*	*	*
682.6	856	361	494	*20	161	260	415	242	224	298	91
682.7	317	185	132	*	*55	112	144	87	75	*101	*53
682.9	*36	*	*	*	*	*	*	*	*	*	*
686	*24	*	*	*	*	*	*	*	*	*	*
693	*26	*	*	*	*	*	*	*	*	*	*
693.0	*24	*	*	*	*	*	*	*	*	*	*
695	*56	*	*	*	*	*	*	*	*	*	*
695.1	*45	*	*	*	*	*	*	*	*	*	*
707	762	378	385	*	133	181	445	194	151	266	151
707.0	510	280	229	*	*105	*112	292	*123	*85	201	*
707.1	237	93	144	*	*	*68	141	*68	66	60	*43
708	*17	*	*	*	*	*	*	*	*	*	*
709	*33	*	*	*	*	*	*	*	*	*	*
710-739	7,162	2,781	4,382	136	1,343	1,777	3,907	1,593	1,518	2,651	1,401
710	149	*	129	*	84	*30	*	*	*	67	*
710.0	114	*	103	*	81	*	*	*	*	*54	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
711	123	*77	*47	*	*39	*	*	*	*	*65	*
711.0	122	*76	*46	*	*38	*	*	*	*	*65	*
711.06	*55	*	*	*	*	*	*	*	*	*	*
714	212	44	169	*	*21	69	115	*63	*19	57	73
714.0	193	36	157	*	*	69	108	*58	*	51	*67
715	2,001	703	1,299	—	49	411	1,541	371	540	704	388
715.1	165	57	108	—	*	*37	113	*32	42	63	*28
715.15	48	*	*32	—	*	*	*38	*	*	*	*
715.16	107	*34	73	—	*	*	74	*	*27	44	*
715.3	729	276	453	—	*	160	562	150	213	234	132
715.35	213	92	121	—	*	55	156	*52	67	58	*36
715.36	503	180	322	—	*	102	398	96	142	173	91
715.8	*25	*	*	—	*	*	*	*	*	*	*
715.89	*25	*	*	—	*	*	*	*	*	*	*
715.9	1,048	346	702	—	*25	197	827	181	269	386	212
715.95	298	120	178	—	*	63	222	68	82	92	56
715.96	699	213	486	—	*	129	560	107	171	274	146
716	74	34	40	*	*	*27	*32	*	*	*33	*
716.1	*24	*	*	—	*	*	*	*	*	*	*
716.9	32	*15	*	*	*	*13	*	*	*	*	*
716.96	*18	*	*	*	*	*	*	*	*	*	*
717	31	18	*12	*	25	*	*	16	*	*	*
717.8	21	14	*	*	20	*	—	*12	*	*	*
717.83	17	*11	*	*	16	*	—	*10	*	*	*
718	47	20	26	*	24	*	*	20	*10	*12	*
718.3	15	*	*	*	*9	*	*	*	*	*	*
718.31	*7	*	*	*	*	*	*	*	*	*	*
719	178	31	147	*	*55	*	101	*34	*	101	*
719.4	92	*19	73	*	*50	*	*29	*	*	*56	*
719.45	39	*	*32	*	*	*	*	*	*	*22	*
719.7	*61	*	*59	—	—	*	*59	*	*	*	*
721	276	121	155	*	*44	83	141	*47	55	109	65
721.0	48	25	*23	*	*	22	*18	*	*	19	*
721.1	72	*29	*43	—	*	*36	*	*	*	*44	*
721.3	114	*38	77	*	*	*18	83	*	*34	*30	*
722	1,040	534	506	*	444	360	236	237	196	422	186
722.0	104	54	50	—	51	47	*	*17	20	54	*13
722.1	657	338	318	*	296	222	139	160	139	250	108
722.10	648	333	315	*	289	220	138	159	134	248	107
722.4	*20	*	*14	—	*	*	*	*	*	*	*
722.5	88	41	47	—	*	*31	34	*	*	*35	*25
722.52	88	41	47	—	*	*31	34	*	*	*35	*25
722.7	64	40	*24	—	*12	*19	*	*	*	*31	*
722.71	46	*30	*	—	*	*	*	*	*	*	*
722.73	*15	*	*	—	*	*	*	*	*	*	*
722.8	42	*	*28	—	*	*	*	*	*	*25	*
722.83	39	*	*24	—	*	*	*	*	*	*22	*
722.9	*42	*	*	—	*	*	*	*	*	*	*
722.93	*37	*	*	—	*	*	*	*	*	*	*
723	75	42	33	*	*33	*29	*	*	*	*30	*
723.0	*40	*28	*	—	*	*	*	*	*	*	*
723.4	*15	*	*	—	*	*	*	*	*	*	*
724	540	201	339	*	91	127	321	99	116	189	136
724.0	250	107	143	*	*	57	180	62	51	89	48

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
724.02	244	105	139	*	*	56	175	62	49	88	45
724.2	112	*20	93	*	*22	31	*59	*	*23	34	*43
724.4	60	*31	*29	—	*18	*	*	*	*	*25	*
724.5	68	*	55	*	*	*	*31	*	*	*	*
726	109	55	54	*	28	35	42	37	21	24	27
726.1	23	*9	*14	—	*	*	*15	*	*	*	*7
726.10	16	*9	*7	—	*	*	*9	*	*	*	*7
726.2	14	*6	*8	—	*	*7	*	*6	*	*	*
726.5	*20	*	*19	—	*	*	*18	*	*	*	*
726.6	*26	*	*	*	*	*	*	*	*	*	*
727	52	23	28	*	*20	*11	*	*12	*14	*17	*
727.0	27	*14	*13	*	*15	*	*	*	*	*	*
727.05	*18	*	*	*	*	*	*	*	*	*	*
728	258	*132	126	*	*84	*	*90	*	*	*92	*
728.8	174	*97	*77	*	*	*	*	*	*	*	*
728.89	*40	*	*	*	*	*	*	*	*	*	*
729	145	51	94	*	37	44	62	*	28	76	*25
729.1	*33	*	*17	*	*	*	*	*	*	*	*
729.5	39	*18	*21	*	*	*13	*	*	*	*	*
729.8	*42	*	*	*	*	*	*	*	*	*	*
730	630	362	268	*	*80	207	321	260	159	146	*65
730.0	321	222	*98	*	*	*118	141	*176	*	*70	*
730.07	188	*143	*	*	*	*	*	*	*	*	*
730.1	*114	*	*	*	*	*	*	*	*	*	*
730.2	188	93	96	*	*	*58	102	*67	*	*58	*
730.27	111	*61	*49	*	*	*	*53	*	*	*	*
732	*19	*	*	*12	*	*	—	*	*	*	*
733	995	239	755	*	110	171	712	182	185	431	198
733.1	642	103	539	*	*	51	577	113	128	260	142
733.13	339	*57	281	*	*	*	321	71	85	130	*53
733.14	107	*	79	*	*	*	86	*	*	*38	*
733.15	*80	*	*67	—	*	*	*65	*	*	*	*
733.19	*83	*	*80	—	*	*	*81	*	*	*	*
733.4	126	65	61	*	*29	53	*42	*28	*	60	*
733.42	101	51	50	*	*	*46	*35	*	*	53	*
733.6	42	*12	29	*	*	*22	*	*	*15	*14	*
733.8	143	51	92	*	46	38	59	*23	*16	73	*31
733.81	*27	*	*	*	*	*	*	*	*	*	*
733.82	116	42	74	—	38	*33	*45	*	*	57	*29
733.9	*14	*	*	*	*	*	*	*	*	*	*
735	*24	*	*19	—	*	*	*	*	*	*	*
736	*30	*	*20	*	*	*	*	*	*	*	*
737	*54	*	*34	*	*	*	*	*	*	*	*
737.3	*46	*	*33	*	*	*	*	*	*	*	*
737.30	*31	*	*24	*	*	*	*	*	*	*	*
738	61	*23	*38	*	*	*	*	*	*	*	*
738.4	50	*	*31	*	*	*	*	*	*	*	*
740-759	1,091	618	473	854	121	62	*55	433	191	300	167
742	*43	*	*	*	*	*	*	*	*	*	*
745	236	127	110	217	*	*	*	120	*	*70	*
745.2	*63	*	*	*62	*	*	—	*	*	*	*
745.4	*46	*	*	*45	*	—	—	*	*	*	*
745.5	*26	*	*	*	*	*	*	*	*	*	*
746	115	*65	*50	*83	*	*	*	*38	*	*	*
746.8	*35	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
747	134	*61	73	90	*	*	*	*51	*	*	*
749	17	*11	*	17	*	—	—	*	*	*	*
749.2	*9	*	*	*	*	—	—	*	*	*	*
750	45	35	*	44	—	*	*	*	*21	*	*
750.5	30	*24	*	30	—	—	—	*	*	*	*
751	104	*54	*50	89	*	*	*	*	*	*	*
753	48	*	*	*	*	*	*	*	*	*	*
754	*29	*	*	*24	*	*	—	*	*	*	*
755	*16	*	*	*	*	*	*	*	*	*	*
756	177	129	*48	*141	*	*15	*	*	*	*83	*
756.1	38	*20	*	*	*	*15	*	*	*	*17	*
756.12	*27	*	*	*	*	*	*	*	*	*	*
759	*45	*	*	*	*	*	*	*	*	*	*
760-779	1,542	876	666	1,542	—	—	—	237	362	483	460
765	528	300	*228	528	—	—	—	*	*	*256	*124
765.1	363	220	*143	363	—	—	—	*	*	*164	*69
769	358	*219	*139	358	—	—	—	*	*	*	*146
770	298	143	155	298	—	—	—	*37	76	*72	113
770.1	*60	*	*	*60	—	—	—	*	*	*	*
770.6	*59	*	*	*59	—	—	—	*	*	*	*
770.8	73	38	*35	73	—	—	—	*	*	*	*
771	68	*46	*21	68	—	—	—	*	*	*	*
771.8	54	*36	*	54	—	—	—	*	*	*	*
773	*15	*	*	*15	—	—	—	*	*	*	*
773.1	*12	*	*	*12	—	—	—	*	*	*	*
774	75	43	32	75	—	—	—	*10	*18	25	22
774.6	65	39	26	65	—	—	—	*	*15	22	21
778	*20	*14	*	*20	—	—	—	*	*	*	*
778.4	*15	*	*	*15	—	—	—	*	*	*	*
779	*30	*	*	*30	—	—	—	*	*	*	*
780-799	993	531	462	117	366	218	293	201	104	501	188
780	130	56	74	49	40	*15	27	40	22	52	*17
780.2	21	*9	*13	*	*	*	*	*	*	*8	*
780.3	63	23	40	26	25	*	*	24	*	21	*
780.6	29	*	*13	*19	*	*	*	*	*	*	*
784	*19	*	*	*	*	*	*	*	*	*	*
786	141	72	70	*10	51	61	19	25	33	61	21
786.0	*11	*	*	*	*	*	*	*	*	*	*
786.5	127	62	65	*	50	59	18	22	30	58	17
786.50	105	50	56	*	38	53	*15	17	27	46	15
786.52	*11	*	*	*	*	*	*	*	*	*	*
786.59	*7	*	*	—	*	*	*	*	*	*	*
788	*10	*	*	*	*	*	*	*	*	*	*
789	80	26	54	15	51	*11	*	21	17	29	*13
789.0	78	25	54	*14	51	*10	*	21	17	29	*11
789.00	19	*	*13	*	*11	*	*	*	*	*	*
789.03	31	*9	22	*	22	*	*	*12	*	*10	*
799	551	329	222	*20	206	112	213	94	*17	322	118
799.9	540	319	220	*20	197	111	212	93	*	312	118

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD–9–CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
800–999	13,834	6,753	7,082	852	3,799	2,559	6,623	3,366	2,777	5,156	2,536
800	56	*49	*	*	*	*	*	*	*	*	*
801	186	144	42	*43	114	*	*	*56	*	*75	*32
801.0	*26	*	*	*	*	*	*	*	*	*	*
801.2	*63	*47	*	*	*53	*	*	*	*	*	*
802	168	134	35	*	118	*	*	43	*33	47	45
802.2	62	49	*	*	52	*	*	*	*	*	*17
802.4	*31	*28	*	*	*24	*	*	*	*	*	*
802.6	*19	*	*	*	*	*	*	*	*	*	*
803	*32	*26	*	*	*	*	*	*	*	*	*
805	392	189	203	*	105	*49	236	77	114	114	87
805.0	90	*44	*	*	*33	*	*	*	*	*	*47
805.02	*47	*	*	*	*	*	*	*	*	*	*
805.2	103	*58	*46	*	*24	*	*62	*	*26	*37	*
805.4	175	80	95	*	*40	*	115	*29	67	59	*20
806	*134	*	*	*	*	*	*	*	*	*	*
807	200	111	90	*	*24	67	109	*38	41	79	*42
807.0	148	83	65	*	*21	44	83	*37	*36	53	*22
807.01	*14	*	*	*	*	*	*	*	*	*	*
807.02	*19	*	*	—	*	*	*	*	*	*	*
807.03	*38	*24	*	—	*	*	*	*	*	*	*
807.09	*32	*	*	—	*	*	*	*	*	*	*
808	368	128	240	*	81	*64	214	88	95	151	*34
808.0	107	*51	*57	*	*	*	*	*	*	*	*
808.2	149	*	127	*	*	*	103	*	*34	*52	*
808.4	*63	*47	*	*	*	*	*	*	*	*	*
808.8	*39	*	*33	*	*	*	*34	*	*	*	*
812	279	67	212	23	*29	87	141	63	55	102	60
812.0	161	*29	132	*	*	*66	87	*43	*23	70	*25
812.00	57	*	*53	*	*	*	*	*	*	*36	*
812.01	61	*	*49	*	*	*	*43	*	*	*	*
812.09	*33	*	*	—	*	*	*	*	*	*	*
812.2	42	*	*33	*	*	*	*26	*	*	*	*
812.21	*27	*	*	*	*	*	*	*	*	*	*
812.4	62	18	45	20	*	*	*	*14	*15	*	*22
812.41	39	*11	*27	15	*	*	*	*	*	*	*18
813	169	73	96	21	61	30	56	48	28	55	38
813.0	31	*20	*	*	*14	*	*	*	*	*	*
813.2	*16	*	*	*	*	*	*	*	*	*	*
813.23	*12	*	*	*	*	*	*	*	*	*	*
813.4	67	*16	51	*	*20	*	*35	*	*	*21	*
813.42	38	*	*32	*	*	*	*	*	*	*	*
813.44	*17	*	*	*	*	*	*	*	*	*	*
813.5	*19	*	*	*	*	*	*	*	*	*	*
815	*19	*	*	—	*	*	*	*	*	*	*
816	33	29	*	*	*17	*	*	*15	*	*	*
816.1	26	25	*	*	*	*	*	*	*	*	*
816.11	*13	*13	*	*	*	*	*	*	*	*	*
820	2,594	639	1,954	*	*61	141	2,383	631	575	1,006	380
820.0	775	193	582	*	*	*47	720	176	218	258	123
820.02	*42	*	*	—	—	*	*36	*	*	*	*
820.03	54	*	*40	*	*	*	*49	*	*	*	*
820.09	642	147	495	—	*	*34	604	161	180	216	85
820.2	1,224	317	907	*	*	62	1,124	330	266	473	155
820.20	75	*	*56	*	*	*	*66	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD–9–CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
820.21	1,014	244	771	*	*	*54	935	244	228	405	138
820.22	134	*	*80	*	*	*	123	*	*	*56	*
820.8	561	113	448	*	*	*	522	123	83	260	95
821	402	213	189	81	84	*64	173	77	89	173	*62
821.0	230	143	87	68	51	*	74	*38	*53	99	*
821.00	*53	*	*	*	*	*	*	*	*	*	*
821.01	177	121	*57	*60	*40	*	*40	*	*44	73	*
821.2	151	*52	98	*	*	*	98	*	*	*66	*
821.23	*70	*	*50	*	*	*	*40	*	*	*	*
822	63	*17	45	*	*12	*	*35	*	*	*25	*
822.0	55	*	41	*	*	*	*34	*	*	*20	*
823	327	190	137	*22	136	78	90	75	62	146	44
823.0	108	56	52	*	*32	*33	*38	*	*26	*40	*
823.00	85	48	37	*	*27	*31	*23	*	*	*32	*
823.2	53	35	*	*	29	*	*	*	*	*20	*
823.20	*16	*	*	*	*	*	*	*	*	*	*
823.22	34	*24	*	*	*17	*	*	*	*	*	*
823.8	88	*31	56	*	*26	*	*36	*	*	*47	*
823.80	*27	*	*	*	*	*	*	*	*	*	*
823.82	56	*25	*30	*	*	*	*	*	*	*	*
823.9	*33	*	*	*	*	*	*	*	*	*	*
824	391	169	223	*	159	91	132	92	83	164	51
824.0	*22	*	*	*	*	*	*	*	*	*	*
824.2	*19	*	*	*	*	*	*	*	*	*	*
824.4	100	*25	75	*	33	*22	42	*30	*15	38	*17
824.6	98	29	69	*	29	29	*40	*23	*25	38	*13
824.8	81	42	39	*	25	*	*36	*26	*17	*29	*10
825	80	46	*34	*	40	*21	*	*23	*	*26	*
825.0	44	*23	*	—	*24	*	*	*	*	*	*
825.2	*21	*	*	*	*	*	*	*	*	*	*
836	23	*14	*	*	*14	*	*	*	*	*	*
840	77	47	30	—	*8	42	27	*25	20	20	*12
840.4	74	45	29	—	*	42	26	*24	19	20	*12
844	61	29	31	*	30	*	*	*23	*16	*17	*
844.2	28	14	*14	*	23	*	*	*	*	*11	*
845	*39	*	*	—	*	*	*	*	*	*	*
845.0	*39	*	*	—	*	*	*	*	*	*	*
846	*36	*	*	—	*	*	*	*	*	*	*
846.0	*26	*	*	—	*	*	*	*	*	*	*
847	47	*22	*25	*	*14	*	*23	*	*	*22	*
847.2	*27	*	*	—	*	*	*	*	*	*	*
850	114	60	54	*9	57	*17	*31	44	*14	46	*11
850.0	27	*17	*	*	*	*	*	*	*	*	*
850.1	59	*24	*34	*	*35	*	*	*16	*	*	*
850.5	*20	*15	*	*	*10	*	*	*	*	*	*
851	155	123	*	*	*76	*	*	*	*	*	*83
851.8	*83	*59	*	*	*	*	*	*	*	*	*
852	418	254	164	*	*92	*	257	*66	*70	202	*81
852.2	374	230	144	*	*76	*	239	*	*57	187	*74
852.20	162	*85	*	*	*	*	145	*	*	*96	*
852.21	*76	*	*49	*	*	*	*59	*	*	*	*
853	*57	*	*	*	*	*	*	*	*	*	*
853.0	*54	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
854	196	118	79	*44	70	*19	64	*42	43	36	75
854.0	194	116	78	*44	68	*18	63	*41	43	34	75
854.00	56	*36	*	*	*	*	*22	*	*	*	*
854.01	26	*19	*	*	*	*	*	*	*	*	*
854.02	48	39	*	*	*18	*	*	*	*	*	*
854.06	*48	*	*	*	*	*	*	*	*	*	*
860	148	105	*44	*	76	*46	*	*	*33	77	*
860.0	62	*43	*	*	*25	*	*	*	*	*27	*
861	71	*32	*39	*	*38	*	*	*	*	*24	*
861.2	*54	*	*	*	*	*	*	*	*	*	*
861.21	*54	*	*	*	*	*	*	*	*	*	*
863	175	166	*	*	*122	*	*	*	*	*	*
864	*66	*50	*	*	*48	*	*	*	*	*	*
864.0	*34	*	*	*	*	*	*	*	*	*	*
865	91	*49	*42	*	*54	*	*	*	*	*	*
865.0	78	*40	*	*	*45	*	*	*	*	*	*
871	*23	*21	*	*	*	*	*	*	*	*	*
873	62	34	*27	*	29	*	*18	*	*15	24	*
873.0	*16	*	*	*	*	*	*	*	*	*	*
873.4	28	*14	*	*	*9	*	*	*	*	*14	*
873.42	*16	*	*	*	*	*	*	*	*	*	*
879	31	*23	*	*	*27	*	*	*	*	*	*
879.2	*21	*	*	—	*	*	*	*	*	*	*
881	*32	*21	*	*	*22	*	*	*	*	*	*
882	*17	*	*	*	*	*	*	*	*	*	*
883	*18	*	*	*	*13	*	*	*	*	*	*
883.2	*11	*	*	*	*	*	*	*	*	*	*
886	23	21	*	*	*	*12	*	*	*	*	*
886.0	18	16	*	*	*	*	*	*	*	*	*
891	56	37	*	*	*28	*	*	*	*	*27	*
891.0	*19	*	*	*	*	*	*	*	*	*	*
891.1	*35	*23	*	*	*	*	*	*	*	*	*
892	*21	*	*	*	*	*	*	*	*	*	*
905	*91	*	*	—	*	*	*	*	*	*	*
920	*25	*	*	*	*	*	*	*	*	*	*
921	*13	*	*	*	*	—	*	*	*	*	*
922	38	18	*20	*	*9	*	*22	*	*	*16	*
922.1	*14	*	*	—	*	*	*	*	*	*	*
924	94	24	71	*	*16	*	72	*	*15	*31	*15
924.0	54	*17	*37	—	*	*	49	*	*	*	*
924.01	49	*13	*36	—	*	*	46	*	*	*	*
924.8	*23	*	*	*	*	*	*	*	*	*	*
935	19	*7	*12	*7	*	*	*	*	*	*	*
935.1	15	*	*	*6	*	*	*	*	*	*	*
942	*59	*	*	*	*	*	*	*	*	*	*
944	*40	*	*	*	*	*	*	*	*	*	*
945	*53	*	*	*	*	*	*	*	*	*	*
946	69	*54	*	*	*	*	*	*	*	*42	*
946.2	*30	*	*	*	*	*	*	*	*	*	*
952	*33	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
958	*32	*25	*	*	*	*	*	*	*	*	*
959	51	*29	*23	*	*20	*	*	*	*	*26	*
965	162	64	98	*	116	*23	*	64	27	45	26
965.0	57	*39	*19	*	*44	*	*	*	*	*	*
965.1	20	*	*12	*	*14	*	*	*	*	*	*
965.4	73	*	61	*	51	*	*	*24	*12	*28	*
966	40	*18	*21	*	*20	*	*	*	*	*	*
966.1	*26	*	*	*	*	*	*	*	*	*	*
967	*18	*	*12	*	*	*	*	*	*	*	*
968	*25	*19	*	*	*15	*	*	*	*	*	*
969	158	73	85	*	100	22	*	*31	55	47	26
969.0	40	*18	23	*	24	*	*	*	*12	*13	*
969.4	86	46	39	*	57	*	*	*	39	*20	*
972	28	*	*20	*	*	*	*20	*	*	*	*
972.1	*20	*	*	*	*	*	*	*	*	*	*
977	47	*36	*11	*	19	*	*	*	*	*16	*
977.9	*38	*	*	*	*13	*	*	*	*	*	*
980	*18	*14	*	*	*13	*	*	*	*	*	*
980.0	*14	*12	*	*	*11	*	—	*	*	*	*
987	*21	*	*	*	*	*	*	*	*	*	*
989	*11	*8	*	*	*	*	*	*	*	*	*
989.5	*9	*	*	*	*	*	*	*	*	*	*
991	*44	*	*	*	*	*	*	*	*	*	*
994	*20	*	*	*	*	*	*	*	*	*	*
995	70	37	33	*	*	*19	*27	*15	*20	*23	*
995.1	*17	*	*	*	*	*	*	*	*	*	*
995.2	*29	*	*	*	*	*	*	*	*	*	*
996	2,234	1,112	1,122	161	448	515	1,110	606	384	897	347
996.0	99	44	55	*	*	*20	72	*36	*21	*28	*13
996.01	61	*20	41	*	*	*	52	*26	*	*21	*
996.1	59	*	*42	*	*	*	*	*	*	*	*
996.2	101	47	*54	59	*	*	*	*18	*27	*	*
996.3	*20	*15	*	*	*	*	*	*	*	*	*
996.39	*12	*	*	*	*	*	*	*	*	*	*
996.4	318	142	175	*	44	76	196	94	57	115	51
996.5	61	*	50	*	*	*	*	*	*	*	*
996.59	50	*	*42	*	*	*	*	*	*	*	*
996.6	828	489	338	*53	185	159	430	217	155	344	111
996.62	354	223	131	*	81	67	175	95	*59	151	*48
996.64	*52	*	*	—	*	*	*44	*	*	*	*
996.66	106	*69	*	*	*	*	*78	*	*	*	*
996.67	78	*	*47	*	*	*	*	*	*	*	*
996.69	125	*71	*54	*	*	*	*	*	*	*	*
996.7	605	262	343	*	92	158	340	142	94	266	103
996.73	168	85	82	*	42	59	62	42	*27	55	43
996.74	296	110	186	*	*	*53	224	*50	*	170	*35
996.77	*36	*	*	—	*	*	*	*	*	*	*
996.78	*34	*	*	*	*	*	*	*	*	*	*
996.8	145	86	*59	*	74	*44	*	*59	*	*60	*
996.81	78	*40	*	*	*40	*	*	*	*	*44	*
997	608	358	250	*	129	200	263	113	96	275	124
997.1	39	*17	*22	—	*	*	*24	*	*	*	*
997.2	55	*33	*	—	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
997.3	46	*	*23	*	*	*	*	*	*	*	*
997.4	242	131	111	*	78	45	107	59	43	94	45
997.5	*38	*15	*	*	*	*	*	*	*	*	*
997.6	162	*130	*	*	*	*	*57	*	*	*89	*
997.62	*140	*113	*	*	*	*	*45	*	*	*	*
998	1,079	486	593	54	242	347	436	260	220	368	230
998.1	120	60	60	*	31	*30	54	35	*29	33	*23
998.2	*41	*	*	*	*	*	*	*	*	*	*
998.3	63	*23	*40	*	*	*	*	*	*	*27	*
998.5	768	340	428	*	149	258	324	172	163	257	175
998.8	55	*27	28	*	*22	*	*	*	*	*18	*
998.89	55	*27	28	*	*21	*	*	*	*	*18	*
999	*35	*	*	*	*	*	*	*	*	*	*
V01-V82	11,659	1,410	10,249	259	8,798	603	1,999	2,474	3,222	3,546	2,417
V10	*16	*	*	—	*	*	*	*	*	*	*
V22	18	—	18	—	18	—	—	*	*	*	*
V22.2	*13	—	*13	—	*13	—	—	*	*	*	*
V24	*8	—	*8	—	*8	—	—	*	*	*	*
V25	*7	—	*7	—	*7	—	—	*	*	*	*
V25.2	*7	—	*7	—	*7	—	—	*	*	*	*
V27	8,367	—	8,367	27	8,331	*	—	1,618	1,692	3,073	1,983
V27.0	6,712	—	6,712	*22	6,680	*	—	1,355	1,279	2,476	1,601
V27.1	38	—	38	*	38	—	—	*	*	*16	*
V27.2	154	—	154	*	154	*	—	33	*25	50	46
V27.9	1,443	—	1,443	*	1,438	*	—	219	371	526	326
V30	*73	*	*	*73	—	—	—	*	*	*	*
V30.1	*73	*	*	*73	—	—	—	*	*	*	*
V45	*31	*	*	*	*	*	*	*	*	*	*
V45.8	*27	*	*	*	*	*	*	*	*	*	*
V54	*24	*	*	*	*	*	*	*	*	*	*
V55	102	61	*41	*	*	*	*40	*31	*30	*	*
V55.3	80	*47	*33	*	*	*	*	*	*	*	*
V57	2,331	929	1,402	*	220	374	1,708	539	1,271	259	262
V57.1	614	218	396	*	*	*	477	289	197	*	*
V57.8	1,709	711	998	*	178	293	1,225	245	1,073	178	212
V57.89	1,709	711	998	*	178	293	1,225	245	1,073	178	212
V58	452	230	221	73	94	127	157	160	162	71	59
V58.0	*41	*	*	*	*	*	*	*	*	*	*
V58.1	354	188	166	70	83	104	98	137	114	57	46
V58.4	*31	*	*	*	*	*	*	*	*	*	*
V58.49	*30	*	*	*	*	*	*	*	*	*	*
V70	33	*25	*	*	*25	*	*	*	*	*	*
V70.7	*33	*25	*	*	*25	*	*	*	*	*	*
V71	35	20	*15	*	*10	*	*	*14	*	*	*9
V71.4	*12	*	*	*	*	*	*	*	*	*	*
V71.8	*14	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimates in table 1; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
All codes	121,712	49,343	72,369	5,841	32,792	25,464	57,615	26,396	29,990	42,679	22,647
001–139	3,527	1,586	1,942	480	978	657	1,413	798	804	1,248	678
003	10	*5	*	*	*	*	*	*	*	*	*
003.0	*7	*	*	*	*	*	*	*	*	*	*
008	236	100	137	84	31	36	85	62	60	77	38
008.4	94	38	56	*6	12	20	57	29	26	23	16
008.45	82	33	49	*	*9	18	52	27	22	21	13
008.6	39	19	20	36	*	*	*	*7	15	10	*7
008.61	35	18	17	35	*	*	—	*6	15	9	*
008.8	97	39	59	41	16	16	24	25	17	42	14
009	20	*8	12	*	*7	*	*8	*	*	10	*
009.0	10	*	*8	*	*	*	*6	*	*	*5	*
011	22	14	*9	*	11	*	*7	*5	*	*9	*5
011.9	18	11	*7	*	10	*	*5	*	*	*8	*
011.90	12	*7	*5	*	*6	*	*	*	*	*6	*
031	24	13	11	*	13	*	*7	*	*	10	*8
031.0	18	*9	*9	*	*8	*	*7	*	*	*9	*
034	27	13	13	14	11	*	*	*	*8	12	*
034.0	25	12	13	13	11	*	*	*	*8	10	*
038	661	320	342	45	97	125	394	148	141	247	126
038.0	47	22	26	*6	9	11	21	10	11	15	12
038.1	136	74	61	*8	23	34	71	35	30	47	23
038.2	20	11	9	*	*6	*	*6	*	*	*5	*6
038.3	*8	*	*	*	*	*	*	*	*	*	*
038.4	160	76	84	*6	20	37	97	38	31	57	33
038.40	17	*8	9	*	*	*	10	*	*	*7	*
038.42	76	30	46	*	*9	19	47	19	18	22	16
038.43	18	12	*6	*	*	*	11	*	*	*9	*
038.49	44	24	20	*	*5	*8	27	*9	*7	19	*9
038.8	29	13	16	*	*	*	18	10	*	*9	*6
038.9	262	119	143	19	34	33	176	47	58	113	44
041	1,259	468	792	97	283	245	634	242	331	451	234
041.0	224	76	148	12	90	39	83	39	63	80	41
041.00	56	25	32	*	13	12	28	14	15	20	*8
041.01	12	*6	*6	*	*	*	*	*	*	*	*
041.02	70	*7	63	*	56	*5	*8	12	17	25	16
041.04	42	20	23	*	*6	*8	27	*6	15	15	*6
041.09	40	18	22	*	9	10	16	*6	11	15	*7
041.1	216	114	102	22	49	58	87	44	52	80	40
041.10	29	15	15	*	*7	*8	11	*8	*6	9	*7
041.11	142	77	65	15	31	41	55	26	35	55	26
041.19	44	22	22	*	10	10	21	10	12	15	*7
041.3	83	26	57	*	11	16	53	19	20	29	15
041.4	355	83	272	31	73	58	193	69	89	129	67
041.5	19	10	*9	*	*	*	*9	*	*	*9	*
041.6	53	18	35	*	*6	*7	38	*8	17	22	*6
041.7	90	47	43	9	11	18	52	19	24	30	17
041.8	205	88	117	12	38	42	112	36	58	68	42
041.84	20	*9	11	*	*	*	11	*	*	*5	*6
041.85	78	32	46	*7	13	15	44	*9	25	28	16
041.86	61	30	32	*	11	17	33	13	16	18	14
041.89	42	15	27	*	10	*6	24	10	11	15	*7
041.9	9	*	*6	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

– Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
042	190	138	53	*	141	43	*	66	22	66	36
047	24	11	13	11	11	*	*	*	*	*9	*7
047.9	22	10	12	9	11	*	*	*	*	*8	*6
052	11	*6	*6	*7	*	*	*	*	*	*	*
052.9	*7	*	*	*	*	*	*	*	*	*	*
053	59	19	40	*	*9	11	39	13	16	15	15
053.1	17	*6	11	—	*	*	14	*	*	*	*
053.19	14	*5	9	—	*	*	12	*	*	*	*
053.9	35	11	24	*	*6	*8	20	*7	10	11	*8
054	58	20	38	*5	34	10	*9	15	*9	23	12
054.1	21	*	17	*	17	*	*	*5	*	*8	*
054.10	13	*	11	*	11	*	*	*	*	*	*
054.2	*7	*	*	*	*	*	*	*	*	*	*
054.9	21	*8	13	*	11	*	*5	*6	*	*9	*
070	101	60	41	*	60	28	11	28	13	38	22
070.1	9	*	*6	*	*5	*	*	*	*	*	*
070.3	25	17	*8	*	18	*	*	*8	*	10	*
070.30	15	10	*5	*	11	*	*	*5	*	*6	*
070.32	10	*7	*	*	*7	*	*	*	*	*	*
070.5	60	36	23	*	34	20	*5	17	*7	22	14
070.51	28	18	10	*	17	10	*	*8	*	10	*5
070.54	31	18	12	*	16	10	*	*8	*	12	*8
075	15	*8	*8	*6	*9	*	*	*	*	*5	*
078	48	28	20	*5	30	*9	*	11	*8	15	14
078.1	14	*	9	*	12	*	*	*	*	*5	*
078.10	*7	*	*5	*	*7	*	—	*	*	*	*
078.5	26	19	*7	*	15	*7	*	*5	*	*8	10
078.8	*8	*	*	*	*	*	*	*	*	*	*
078.89	*6	*	*	*	*	*	*	*	*	*	*
079	200	95	105	128	38	17	18	46	48	64	43
079.8	73	43	30	68	*	*	*	13	23	21	16
079.89	71	43	28	68	*	*	*	13	23	20	16
079.9	120	50	70	56	32	15	18	31	24	40	24
079.99	115	49	66	56	28	15	18	30	23	38	24
088	*6	*	*	*	*	*	*	*	*	*	*
088.8	*6	*	*	*	*	*	*	*	*	*	*
088.81	*6	*	*	*	*	*	*	*	*	*	*
094	*7	*	*	—	*	*	*	*	*	*	*
097	10	*	*6	*	*6	*	*	*	*	*	*
097.1	*6	*	*	—	*	*	*	*	*	*	*
098	*9	*	*8	*	*8	*	*	*	*	*	*
110	45	23	23	*	10	*9	25	13	18	10	*
110.1	20	*9	11	—	*	*	15	*7	*8	*	*
110.3	*6	*	*	—	*	*	*	*	*	*	*
110.4	9	*6	*	*	*	*	*	*	*	*	*
110.5	*6	*	*	*	*	*	*	*	*	*	*
111	*6	*	*	*	*	*	*	*	*	*	*
112	247	109	138	38	70	47	93	59	58	83	47
112.0	133	67	66	23	39	25	47	31	33	47	22
112.1	18	—	18	*	10	*	*	*	*	*	*
112.2	21	*6	15	*	*	*	12	*8	*	*8	*
112.3	14	*5	9	*7	*	*	*5	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
112.4	13	*8	*5	*	*	*	10	*	*	*5	*
112.5	*7	*	*	*	*	*	*	*	*	*	*
112.8	35	17	18	*	11	*8	13	*7	*7	11	9
112.84	25	13	11	*	*9	*6	9	*6	*	*8	*6
112.89	*8	*	*5	*	*	*	*	*	*	*	*
112.9	*6	*	*	*	*	*	*	*	*	*	*
117	23	14	*9	*	10	*	*7	*6	*	*8	*6
117.3	9	*	*	*	*	*	*	*	*	*	*
117.5	*8	*7	*	*	*5	*	*	*	*	*	*
117.9	*6	*	*	*	*	*	*	*	*	*	*
130	*8	*6	*	*	*5	*	—	*	*	*	*
131	18	*	16	*	14	*	*	*	*	9	*
131.0	17	*	14	*	13	*	*	*	*	*9	*
131.00	*7	*	*6	—	*6	*	*	*	*	*	*
131.01	*7	—	*7	*	*6	*	*	*	*	*	*
135	36	*8	28	*	13	13	10	10	*7	14	*5
136	39	25	13	*	18	9	10	*8	*8	16	*7
136.3	27	20	*8	*	17	*8	*	*7	*	9	*6
136.9	10	*5	*	*	*	*	*7	*	*	*6	*
138	19	11	*8	—	*	*	14	*	*	*6	*6
140–239	4,488	1,876	2,612	107	694	1,382	2,305	1,128	1,022	1,487	851
141	10	*8	*	*	*	*5	*	*	*	*	*
150	23	16	*8	—	*	*6	16	9	*6	*	*
150.5	*6	*	*	—	*	*	*	*	*	*	*
150.9	11	*7	*	—	—	*	*7	*	*	*	*
151	43	28	15	—	*	10	30	11	*9	13	11
151.0	10	*9	*	—	*	*	*6	*	*	*	*
151.8	*8	*	*	—	*	*	*6	*	*	*	*
151.9	14	*8	*6	—	*	*	11	*	*	*5	*
152	*8	*	*	—	*	*	*6	*	*	*	*
153	162	74	88	*	*	42	116	34	41	62	26
153.1	10	*	*6	*	*	*	*6	*	*	*	*
153.2	*7	*	*	—	*	*	*5	*	*	*	*
153.3	31	16	15	—	*	*9	21	*6	*6	11	*8
153.4	27	11	16	—	*	*	21	*5	*8	*8	*5
153.6	23	12	11	—	*	*	18	*	*7	*8	*
153.8	10	*	*6	—	*	*	*7	*	*	*	*
153.9	46	20	26	—	*	13	32	*7	12	23	*
154	68	34	35	*	*8	13	47	22	19	18	10
154.0	24	13	11	*	*	*	18	*6	*	10	*
154.1	39	19	20	—	*5	*7	26	15	12	*6	*6
155	22	10	11	*	*	*7	11	*6	*	*8	*6
155.0	12	*8	*	*	*	*	*5	*	*	*	*
155.2	*7	*	*5	—	*	*	*	*	*	*	*
156	12	*	*9	—	—	*	10	*	*	*	*
156.0	*6	*	*5	—	—	*	*5	*	*	*	*
157	65	35	30	—	*	22	42	14	15	22	14
157.0	18	*9	10	—	*	*6	12	*	*5	*	*6
157.8	9	*	*6	—	*	*	*7	*	*	*	*
157.9	32	20	12	—	*	13	19	*	*8	14	*6
158	11	*	*6	*	*	*	*5	*	*	*	*
158.0	*5	*	*	*	*	*	*	*	*	*	*
158.8	*5	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
161	18	14	*	—	*	*6	10	*	*	*7	*
161.1	*6	*	*	—	*	*	*	*	*	*	*
161.9	*6	*	*	—	*	*	*	*	*	*	*
162	377	210	167	*	17	123	237	93	87	129	68
162.2	28	18	10	—	*	10	17	*	*7	13	*
162.3	83	43	40	—	*	26	55	22	18	27	17
162.4	*8	*	*	*	*	*	*5	*	*	*	*
162.5	46	25	20	*	*	12	31	12	*9	15	9
162.8	37	22	15	—	*	14	21	18	*6	10	*
162.9	176	98	78	*	*9	58	109	36	43	62	34
170	18	13	*6	*9	*6	*	*	10	*	*	*
170.7	*8	*5	*	*	*	*	*	*	*	*	*
171	15	*7	*8	*	*	*	*	*	*	*	*
172	12	*7	*	—	*	*	*6	*	*	*	*
173	26	15	10	*	*	*6	19	*5	*7	10	*
173.3	*9	*5	*	—	*	*	*7	*	*	*	*
174	180	—	180	*	21	72	86	44	40	62	34
174.1	*5	—	*5	—	*	*	*	*	*	*	*
174.2	*7	—	*7	—	*	*	*	*	*	*	*
174.4	26	—	26	—	*	9	13	*6	*9	*8	*
174.8	42	—	42	—	*	17	20	17	*8	9	*7
174.9	86	—	86	*	11	35	40	16	17	37	16
176	15	14	*	—	12	*	*	*	*	*	*6
176.0	*6	*6	*	—	*5	*	—	*	*	*	*
180	27	—	27	*	11	11	*5	*	*5	12	*
180.8	*6	—	*6	—	*	*	*	*	*	*	*
180.9	18	—	18	*	*7	*7	*	*	*	9	*
182	43	—	43	*	*	14	24	10	13	11	*9
182.0	42	—	42	*	*	14	24	10	13	11	*9
183	68	—	68	*	*8	28	32	21	14	22	11
183.0	67	—	67	*	*8	27	32	21	14	22	11
184	*6	—	*6	*	*	*	*	*	*	*	*
185	212	212	—	—	*	40	172	47	53	79	34
186	*6	*6	—	—	*6	*	*	*	*	*	*
186.9	*6	*6	—	—	*6	*	*	*	*	*	*
188	77	57	20	*	*	17	58	24	15	24	15
188.2	*7	*6	*	—	—	*	*6	*	*	*	*
188.4	*7	*6	*	—	*	*	*6	*	*	*	*
188.8	21	16	*	—	*	*5	15	*9	*	*	*
188.9	32	23	9	—	*	*8	24	*9	*7	11	*5
189	50	27	22	*	*	15	30	12	*8	22	*8
189.0	40	22	18	*	*	11	24	10	*6	17	*7
189.1	*6	*	*	—	*	*	*	*	*	*	*
191	48	30	18	*8	13	15	12	12	13	14	*9
191.1	10	*8	*	*	*	*	*	*	*	*	*
191.2	*8	*5	*	*	*	*	*	*	*	*	*
191.3	*5	*	*	*	*	*	*	*	*	*	*
191.8	*7	*	*	*	*	*	*	*	*	*	*
191.9	*8	*	*	*	*	*	*	*	*	*	*
193	21	*	17	*	*7	*7	*7	*7	*	*7	*
194	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
195	14	*6	*8	*	*	*	9	*	*	*	*
196	216	84	131	*	24	76	114	49	53	72	41
196.0	27	15	12	*	*	11	12	*6	*	11	*5
196.1	32	16	16	*	*	12	18	10	*7	10	*
196.2	72	36	36	*	*	19	48	13	18	26	15
196.3	50	*	46	*	10	23	16	11	13	16	10
196.5	9	*5	*	*	*	*	*6	*	*	*	*
196.6	12	*	*8	—	*	*	*7	*	*	*	*
196.9	11	*	*7	*	*	*5	*6	*	*	*	*
197	566	244	322	*9	45	192	320	157	127	173	110
197.0	124	59	65	*6	14	41	63	36	31	36	21
197.1	19	10	9	*	*	*6	9	*	*	*6	*7
197.2	72	29	43	*	*6	25	40	23	15	21	13
197.4	15	*5	9	—	*	*	10	*	*	*	*
197.5	20	*	15	—	*	*8	11	*7	*5	*6	*
197.6	75	22	53	*	*7	25	42	22	15	20	18
197.7	218	104	114	*	12	71	133	57	48	72	40
197.8	21	*9	12	—	*	*9	11	*	*	*7	*5
198	495	219	275	*7	47	173	267	129	119	161	85
198.1	14	9	*	—	*	*5	*9	*	*	*5	*
198.2	16	*5	10	—	*	*	12	*5	*	*	*
198.3	98	45	53	*	*8	44	44	22	25	32	17
198.4	*8	*	*5	*	*	*	*	*	*	*	*
198.5	210	103	107	*	21	61	126	55	56	68	30
198.6	*7	—	*7	—	*	*	*	*	*	*	*
198.7	15	11	*	—	*	*	10	*	*	*6	*
198.8	124	42	82	*	15	45	61	31	25	41	27
198.82	25	*	20	*	*6	*8	11	*8	*	*7	*5
198.89	96	37	59	*	9	36	49	22	20	34	21
199	89	38	51	*	*9	22	57	25	19	30	15
199.0	13	*5	*8	*	*	*	*6	*	*	*	*
199.1	75	32	43	*	*7	17	51	22	17	25	12
200	38	17	21	*	*	*9	22	*9	10	*9	10
200.0	22	10	12	*	*	*6	13	*	*7	*5	*
200.00	9	*	*	*	*	*	*6	*	*	*	*
200.8	*8	*	*6	—	*	*	*6	*	*	*	*
201	21	11	10	*	11	*	*	*	*6	*5	*5
201.9	17	10	*7	*	*9	*	*	*	*5	*	*
201.90	*7	*	*	*	*	*	*	*	*	*	*
202	113	63	49	*	17	36	57	29	29	33	22
202.0	*7	*	*	*	*	*	*	*	*	*	*
202.8	100	56	44	*	15	32	51	24	26	30	20
202.80	56	32	25	*	*8	18	29	14	14	17	12
202.81	10	*6	*	*	*	*	*6	*	*	*	*
202.83	15	9	*5	*	*	*	*8	*	*	*	*
202.88	9	*5	*	—	*	*	*	*	*	*	*
203	58	31	27	—	*	15	39	15	18	13	13
203.0	57	31	26	—	*	14	39	15	18	12	12
203.00	54	29	25	—	*	13	38	14	16	12	12
204	91	47	43	21	9	9	52	30	18	26	17
204.0	33	19	15	21	9	*	*	13	*	10	*6
204.00	25	15	11	15	*8	*	*	*9	*	*9	*
204.01	*8	*	*	*6	*	*	*	*	*	*	*
204.1	56	28	27	*	—	*8	48	17	13	15	10
204.10	53	27	26	—	—	*8	45	16	13	15	10

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
205	42	19	23	*	*7	16	17	15	*7	13	*7
205.0	29	12	17	*	*	10	11	10	*5	10	*
205.00	23	10	13	*	*	*8	10	*8	*	*7	*
205.01	*5	*	*	*	*	*	*	*	*	*	*
205.1	12	*6	*6	*	*	*	*5	*	*	*	*
205.10	12	*6	*6	*	*	*	*5	*	*	*	*
208	12	*	*7	*	*	*	*7	*	*	*7	*
208.0	*6	*	*	*	*	*	*	*	*	*	*
208.00	*6	*	*	*	*	*	*	*	*	*	*
208.9	*5	*	*	*	*	*	*	*	*	*	*
210	*9	*	*	*	*	*	*	*	*	*	*
210.2	*8	*	*	—	*	*	*	*	*	*	*
211	147	63	83	*	*9	33	104	42	33	50	21
211.1	11	*	*8	*	*	*	*7	*	*	*	*
211.3	118	53	64	*	*6	26	86	34	25	41	17
211.4	*9	*	*5	*	*	*	*7	*	*	*	*
212	*6	*	*	*	*	*	*	*	*	*	*
214	22	12	10	*	*	*7	*8	*7	*	*7	*
214.1	*6	*	*	*	*	*	*	*	*	*	*
214.8	*6	*	*	*	*	*	*	*	*	*	*
215	*7	*	*5	*	*	*	*	*	*	*	*
216	22	*6	16	*	12	*	*	*5	*	*8	*
216.5	*7	*	*6	*	*	*	*	*	*	*	*
217	*6	*	*5	—	*	*	*	*	*	*	*
218	435	—	435	*	229	179	27	77	86	170	101
218.0	49	—	49	—	25	22	*	12	10	14	13
218.1	116	—	116	—	62	49	*6	25	24	37	30
218.2	65	—	65	—	35	28	*	12	10	29	15
218.9	205	—	205	*	108	80	17	28	42	90	44
219	*7	—	*7	—	*	*	*	*	*	*	*
220	60	—	60	*	27	24	*8	13	11	21	14
221	*5	—	*5	—	*	*	*	*	*	*	*
225	27	10	16	*	*	*6	14	10	*6	*6	*5
225.2	17	*	12	*	*	*	10	*7	*	*	*
226	16	*	13	*	9	*	*	*	*	*	*
227	28	10	18	*	*7	10	12	*7	*5	*9	*7
227.0	*6	*	*	—	*	*	*	*	*	*	*
227.1	13	*	*9	—	*	*	*5	*	*	*	*
227.3	9	*	*	*	*	*	*	*	*	*	*
228	19	*8	11	*5	*6	*	*	*	*	*	*5
228.0	16	*6	9	*	*5	*	*	*	*	*	*
228.09	*6	*	*	*	*	*	*	*	*	*	*
230	*6	*	*	—	*	*	*	*	*	*	*
233	35	*	32	—	14	11	10	*8	*8	12	*8
233.0	13	—	13	—	*	*6	*5	*	*	*	*
233.1	17	—	17	—	11	*	*	*	*	*7	*
235	23	11	12	*	*	*5	15	*	*	10	*
235.2	11	*5	*6	*	*	*	*	*	*	*5	*
236	*9	*	*7	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
237	18	*8	10	*	*	*	*6	*	*	*7	*
237.5	*6	*	*	*	*	*	*	*	*	*	*
237.7	*9	*	*6	*	*	*	*	*	*	*	*
237.70	*7	*	*	*	*	*	*	*	*	*	*
238	81	50	31	*	*6	11	62	20	23	25	13
238.4	17	11	*6	*	*	*	12	*	*7	*5	*
238.7	56	35	21	*	*	*6	47	16	16	15	10
239	28	14	14	*	*	*	18	*7	*	11	*6
239.5	*6	*	*	—	*	*	*	*	*	*	*
240–279	12,023	4,980	7,043	492	1,705	3,204	6,623	2,522	3,179	4,244	2,078
240	14	*	13	*	*	*	*8	*	*	*6	*
240.9	14	*	13	*	*	*	*8	*	*	*6	*
241	26	*5	21	*	*6	9	10	*7	*5	*8	*5
241.0	*7	*	*6	*	*	*	*	*	*	*	*
241.1	13	*	11	*	*	*	*6	*	*	*	*
241.9	*6	*	*	—	*	*	*	*	*	*	*
242	60	12	48	*	19	20	20	15	12	26	*8
242.0	19	*	15	*	*8	*8	*	*	*	10	*
242.00	18	*	14	*	*8	*8	*	*	*	10	*
242.9	37	*8	29	—	11	10	16	9	*9	13	*5
242.90	35	*7	27	—	10	9	15	*9	*9	12	*
244	935	177	758	*6	94	202	633	213	265	263	194
244.0	38	*	35	—	*	10	26	*5	12	11	9
244.1	10	*	9	—	*	*	*7	*	*	*	*
244.9	880	171	708	*6	89	189	596	205	249	248	177
245	14	*	13	*	*7	*	*	*	*	*	*
245.2	11	*	10	*	*6	*	*	*	*	*	*
246	12	*	10	—	*	*	*	*	*	*	*
246.9	*7	*	*6	—	*	*	*	*	*	*	*
250	4,157	1,875	2,282	32	459	1,276	2,390	905	1,009	1,532	710
250.0	2,693	1,184	1,510	12	244	796	1,641	603	674	1,007	409
250.00	1,622	743	878	*	113	462	1,046	378	409	573	262
250.01	641	264	378	*6	74	201	360	147	159	246	90
250.02	263	110	154	*	25	79	158	48	62	117	36
250.03	167	67	100	*	32	53	78	30	45	72	20
250.1	123	60	63	14	63	32	14	23	28	50	22
250.10	12	*5	*6	*	*5	*	*	*	*	*6	*
250.11	42	20	22	*7	24	*7	*	10	*7	16	*9
250.12	10	*	*7	—	*	*	*	*	*	*	*
250.13	59	31	28	*6	30	17	*5	9	17	25	9
250.2	17	*8	9	—	*	*6	*8	*	*	*7	*
250.22	*6	*	*	—	*	*	*	*	*	*	*
250.3	9	*	*	*	*	*	*	*	*	*	*
250.4	364	166	198	*	41	132	189	70	70	129	94
250.40	140	68	72	*	*	46	90	26	21	53	39
250.41	154	65	89	*	24	62	69	34	34	49	37
250.42	22	*8	13	—	*	*6	14	*	*	*8	*5
250.43	48	24	23	*	12	18	17	*6	11	18	13
250.5	157	67	91	*	25	61	70	34	35	48	39
250.50	38	18	20	—	*	15	20	*8	*7	12	11
250.51	81	33	48	*	14	33	34	19	19	23	21
250.52	14	*	10	—	*	*	*8	*	*	*	*
250.53	25	12	14	—	*8	*9	*9	*6	*6	*9	*5
250.6	305	140	165	*	42	106	156	65	70	106	63

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
250.60	100	49	51	*	*	30	66	19	19	38	24
250.61	133	57	76	*	22	50	61	34	32	44	23
250.62	25	12	13	—	*	*8	16	*5	*6	*8	*5
250.63	47	22	25	—	15	18	14	*7	14	16	10
250.7	196	106	90	*	*9	60	126	40	47	71	39
250.70	76	45	31	*	*	18	57	15	15	28	18
250.71	74	37	36	*	*5	29	39	15	20	23	16
250.72	18	10	*8	—	*	*	13	*6	*	*6	*
250.73	28	14	14	—	*	*8	17	*	*8	14	*
250.8	236	114	122	*	20	63	153	55	62	86	33
250.80	79	40	39	—	*	19	58	19	20	28	12
250.81	76	39	37	*	*9	25	41	21	19	24	12
250.82	45	17	28	—	*	10	33	*8	13	17	*6
250.83	36	18	19	*	*7	*9	20	*6	10	17	*
250.9	57	26	30	*	9	19	27	*8	17	24	*7
250.90	*8	*	*	*	*	*	*	*	*	*	*
250.91	*8	*	*	*	*	*	*	*	*	*	*
250.92	21	9	12	—	*	*8	11	*	*6	*9	*
250.93	19	*8	11	*	*	*7	10	*	*6	*8	*
251	44	17	27	*	9	12	19	12	*6	15	10
251.2	22	*9	13	*	*6	*	9	*6	*	*8	*5
251.8	15	*	11	*	*	*7	*6	*	*	*5	*
252	36	*7	29	*	*	10	21	*8	11	*9	*8
252.0	29	*6	23	—	*	*8	17	*7	10	*6	*7
252.1	*7	*	*6	*	*	*	*	*	*	*	*
253	54	24	30	*5	10	12	27	17	15	15	*8
253.2	*6	*	*	*	*	*	*	*	*	*	*
253.5	*8	*	*	*	*	*	*	*	*	*	*
253.6	32	12	20	*	*	*7	21	11	*8	10	*
255	52	19	34	*	12	16	22	*8	15	19	10
255.0	12	*	10	*	*	*5	*	*	*	*6	*
255.4	25	12	13	*	*	*5	15	*	*9	*7	*6
256	*8	—	*8	*	*6	*	*	*	*	*	*
256.4	*6	—	*6	—	*	—	*	*	*	*	*
259	10	*6	*	*	*	*	*	*	*	*	*
260	16	*7	9	*	*	*	11	*	*5	*	*
261	26	12	14	*	*	*	16	*7	*7	*6	*5
262	13	*9	*	*	*	*	*7	*6	*	*	*
263	344	154	190	*6	31	66	241	73	102	120	49
263.0	20	12	*8	*	*	*	13	*7	*	*7	*
263.1	24	13	11	*	*	*	14	11	*6	*	*
263.8	33	15	18	*	*	*5	23	*	12	13	*
263.9	268	114	153	*	23	52	189	51	80	95	41
266	29	10	19	*	*	*5	22	*8	*8	10	*
266.2	28	10	18	*	*	*5	22	*8	*7	10	*
269	*7	*	*	*	*	*	*5	*	*	*	*
270	*6	*	*	*	*	*	*	*	*	*	*
271	23	9	14	*	*	*6	12	*7	*	*6	*6
271.3	21	*8	13	*	*	*5	12	*6	*	*5	*6

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
272	1,023	526	496	*	114	428	479	241	275	313	194
272.0	572	284	288	*	63	252	257	166	149	165	93
272.1	33	17	16	*	*6	16	11	*	*6	16	*6
272.2	17	*9	*8	*	*	*7	*7	*	*	*6	*5
272.4	379	206	174	*	36	145	198	62	113	120	85
272.9	10	*5	*	*	*	*	*	*	*	*	*
273	55	26	29	*	10	11	30	11	15	21	*7
273.1	*9	*5	*	—	*	*	*8	*	*	*	*
273.8	39	18	21	*	9	*8	18	*5	13	16	*5
274	174	117	57	—	9	40	125	44	43	53	34
274.0	54	35	19	—	*	10	40	12	14	16	12
274.9	117	81	37	—	*	29	84	31	29	36	21
275	204	94	110	*6	47	55	96	43	49	78	34
275.0	*7	*	*	*	*	*	*	*	*	*	*
275.2	69	30	39	*	17	18	33	16	18	27	*8
275.3	37	19	18	*	11	11	13	*5	*6	17	*8
275.4	90	39	51	*	18	24	47	20	21	33	16
276	4,042	1,635	2,407	387	656	747	2,252	746	1,128	1,512	657
276.0	94	37	57	*7	11	13	63	15	27	35	17
276.1	759	322	437	25	94	159	481	123	225	287	124
276.2	155	75	80	22	26	34	73	30	46	49	30
276.3	38	17	21	*	*7	*8	19	*	12	17	*
276.4	19	*8	11	*	*	*5	10	*	*6	*7	*
276.5	1,791	713	1,078	287	311	244	950	352	465	696	278
276.6	61	29	32	*	10	21	28	*8	13	23	18
276.7	242	128	114	10	35	52	145	43	67	83	48
276.8	776	264	513	22	146	192	417	155	225	273	123
276.9	106	42	64	*8	15	17	66	11	42	42	12
277	45	27	18	14	18	*6	*6	15	14	10	*7
277.0	25	15	11	11	14	*	*	*9	*9	*6	*
277.00	25	15	11	11	14	*	*	*9	*9	*6	*
278	567	190	377	*6	162	247	151	108	167	190	101
278.0	556	186	370	*6	159	241	150	105	165	186	99
278.00	429	151	277	*6	111	186	125	84	129	138	78
278.01	127	35	92	*	47	55	25	21	36	48	22
278.8	*5	*	*	*	*	*	*	*	*	*	*
279	19	*8	11	*	*7	*	*	*	*	*	*6
279.0	10	*	*6	*	*	*	*	*	*	*	*
280–289	3,921	1,536	2,385	179	954	815	1,972	839	915	1,417	750
280	583	200	383	19	130	119	314	117	136	228	102
280.0	356	128	228	*	76	79	199	71	78	144	63
280.8	*7	*	*6	*	*	*	*	*	*	*	*
280.9	217	70	147	15	52	40	111	44	56	81	36
281	107	40	67	*	11	18	78	22	25	46	15
281.0	37	12	25	*	*	*	32	10	10	16	*
281.1	10	*	*6	—	*	*	*9	*	*	*	*
281.9	51	20	31	*	*8	10	33	*8	11	22	10
282	130	53	76	29	77	11	13	36	31	46	16
282.4	18	*5	13	*	*6	*	*6	*5	*	*	*
282.5	18	*	14	*	13	*	*	*	*	10	*
282.6	83	39	44	22	55	*	*	24	22	29	*8
282.60	18	*8	10	*7	*8	*	*	*	*	*8	*
282.61	*5	*	*	*	*	*	—	*	*	*	*
282.62	59	27	31	10	46	*	*	18	15	21	*5

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
283	27	9	18	*	*	*6	17	9	*	*7	*8
283.0	16	*	11	*	*	*	13	*6	*	*	*
283.1	*6	*	*	*	*	*	*	*	*	*	*
284	169	83	85	*7	35	43	84	38	41	52	38
284.8	157	78	79	*5	33	41	78	36	37	47	37
284.9	11	*	*6	*	*	*	*6	*	*	*	*
285	2,149	786	1,364	49	523	437	1,141	432	508	782	428
285.1	896	329	567	10	197	194	495	160	231	306	199
285.8	34	15	19	*	*7	*	21	*9	*8	10	*7
285.9	1,214	438	776	37	319	237	620	259	268	465	222
286	114	61	54	*8	28	29	49	34	21	38	21
286.0	11	10	*	*	*	*	*	*	*	*	*
286.4	*7	*	*5	*	*	*	*	*	*	*	*
286.5	*8	*	*	—	*	*	*6	*	*	*	*
286.6	13	*6	*6	*	*	*	*5	*	*	*	*
286.7	23	11	12	*	*	*7	11	*8	*	*7	*
286.9	45	23	21	*	12	12	20	11	*8	16	9
287	286	142	145	20	62	66	138	69	64	99	54
287.3	37	13	23	*7	*7	*	19	13	*9	*9	*7
287.4	54	26	28	*	14	14	24	16	11	16	10
287.5	188	98	90	*9	40	47	93	40	41	72	35
288	259	118	141	29	63	66	100	63	62	88	46
288.0	158	75	84	18	37	45	58	40	42	48	28
288.3	*7	*	*	*	*	*	*	*	*	*	*
288.8	90	37	53	10	25	18	37	21	19	35	16
289	97	44	53	15	23	21	37	19	24	30	23
289.0	*6	*	*	—	*	*	*	*	*	*	*
289.2	19	*8	12	10	*9	*	*	*	*	*8	*
289.3	*6	*	*	*	*	*	*	*	*	*	*
289.4	10	*	*7	*	*	*	*	*	*	*	*
289.5	11	*6	*	*	*	*	*5	*	*	*	*
289.59	10	*6	*	*	*	*	*5	*	*	*	*
289.8	26	14	13	*	*6	*6	14	*6	*7	*5	*8
289.9	18	*7	11	*	*	*	9	*	*	*6	*
290-319	7,351	3,685	3,665	227	3,484	1,627	2,014	1,834	2,004	2,387	1,125
290	453	156	297	*	*	17	435	93	115	174	71
290.0	269	89	180	*	*	*6	262	61	64	97	47
290.1	13	*6	*7	—	*	*	10	*	*	*6	*
290.10	*8	*	*	—	*	*	*6	*	*	*	*
290.2	35	11	25	—	—	*	34	*	16	14	*
290.20	13	*6	*7	—	—	*	13	*	*5	*7	*
290.21	22	*5	17	—	—	*	21	*	11	*8	*
290.3	45	17	27	—	—	*	44	10	11	17	*7
290.4	91	33	58	—	*	*5	85	15	22	40	14
290.40	68	24	44	—	*	*	64	12	15	28	12
290.41	13	*	9	—	*	*	12	*	*6	*	*
290.43	*5	*	*	—	—	*	*	*	*	*	*
291	207	158	48	*	96	68	42	43	71	59	34
291.0	52	45	*7	—	21	22	10	13	13	19	*8
291.2	22	12	10	—	*	*	15	*	*9	*6	*
291.3	*7	*	*	*	*	*	*	*	*	*	*
291.8	120	93	28	*	68	38	15	25	46	31	19

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
292	128	68	60	*	73	21	32	38	37	34	19
292.0	53	28	25	*	41	*9	*	18	18	10	*7
292.1	14	*8	*6	*	*6	*	*6	*	*	*	*
292.11	*7	*	*	—	*	*	*	*	*	*	*
292.12	*8	*	*	*	*	*	*	*	*	*	*
292.8	51	28	23	*	19	10	22	17	12	17	*6
292.81	30	14	16	*	*	*6	20	*8	*7	10	*
292.84	13	*9	*	—	10	*	*	*6	*	*	*
292.89	*7	*	*	—	*6	*	*	*	*	*	*
292.9	9	*	*5	—	*7	*	*	*	*	*	*
293	116	53	63	*	12	15	87	29	29	29	28
293.0	77	35	42	*	*	*8	66	15	20	21	22
293.8	25	11	14	*	*8	*7	10	13	*6	*	*
293.83	19	*9	10	*	*7	*6	*6	12	*	*	*
293.9	13	*6	*7	*	*	*	10	*	*	*	*
294	275	105	169	*	13	12	250	51	80	104	40
294.1	47	19	28	—	*	*	43	*8	14	20	*6
294.8	213	79	134	*	*5	*8	199	40	63	79	30
294.9	14	*7	*7	—	*	*	*6	*	*	*5	*
295	382	201	181	*	216	108	56	120	97	107	58
295.3	104	59	44	*	62	29	13	38	17	29	20
295.30	29	14	15	*	17	*8	*	*6	*6	9	*8
295.32	28	17	11	—	14	*8	*5	12	*5	*6	*5
295.34	44	27	17	—	29	12	*	20	*	13	*7
295.6	40	20	20	—	17	15	*9	14	11	12	*
295.60	*8	*	*	—	*	*	*	*	*	*	*
295.62	17	*7	9	—	*6	*6	*	*	*	*6	*
295.64	15	9	*6	—	*8	*6	*	*6	*	*	*
295.7	143	73	69	*	89	38	15	43	43	36	21
295.70	79	37	42	*	47	23	*8	22	23	23	11
295.72	*9	*	*5	—	*	*	*	*	*	*	*
295.74	53	33	20	—	35	12	*6	19	18	*9	*7
295.9	79	39	40	*	38	23	18	22	23	23	11
295.90	58	28	30	*	26	18	14	11	20	18	9
295.92	*7	*	*	—	*	*	*	*	*	*	*
295.94	13	*7	*6	—	9	*	*	10	*	*	*
296	869	335	534	40	485	184	160	219	241	323	86
296.2	295	124	170	20	161	55	59	69	75	125	26
296.20	152	64	88	*8	80	33	31	41	46	48	16
296.22	26	11	16	*	19	*	*	*	*	19	*
296.23	69	29	40	*6	37	10	16	13	15	37	*
296.24	41	18	23	*	22	*7	9	12	*8	19	*
296.3	291	97	194	10	163	59	58	79	89	107	15
296.30	103	34	69	*	56	22	22	28	38	32	*
296.32	22	*6	16	*	16	*	*	*	9	*9	*
296.33	107	37	70	*6	62	20	19	27	27	47	*7
296.34	52	16	36	*	25	12	14	20	14	16	*
296.4	51	19	31	*	25	16	*8	14	14	12	11
296.40	27	10	17	*	13	*8	*	*	*8	*7	*7
296.43	*5	*	*	—	*	*	*	*	*	*	*
296.44	17	*7	10	*	*8	*6	*	*6	*	*	*
296.5	54	17	37	*	32	13	*7	11	15	20	*8
296.50	32	9	22	*	17	*9	*5	*6	10	11	*
296.53	10	*	*7	*	*6	*	*	*	*	*	*
296.54	9	*	*	—	*8	*	*	*	*	*	*
296.6	52	19	33	*	35	10	*	11	12	23	*6

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
296.60	29	13	17	*	20	*5	*	*5	*9	12	*
296.63	*8	*	*6	*	*6	*	*	*	*	*	*
296.64	12	*	*8	*	*8	*	*	*	*	*	*
296.7	87	38	49	*	47	21	15	22	28	22	16
296.8	26	13	13	*	12	*7	*5	*9	*	*9	*
296.80	15	*7	*8	*	*6	*	*	*	*	*5	*
296.89	*6	*	*	*	*	*	*	*	*	*	*
296.9	11	*6	*5	*	*7	*	*	*	*	*	*
296.90	*8	*	*	*	*6	*	*	*	*	*	*
297	19	*6	13	*	*	*5	10	*5	*6	*	*
297.1	*7	*	*5	—	*	*	*	*	*	*	*
297.9	11	*	*7	*	*	*	*5	*	*	*	*
298	158	78	80	*	60	21	73	43	25	59	30
298.9	146	72	73	*	55	19	68	38	23	55	30
299	*6	*	*	*	*	*	—	*	*	*	*
300	589	205	384	11	222	158	198	140	177	198	74
300.0	298	97	202	*	75	82	139	71	89	104	34
300.00	234	78	156	*	45	63	124	54	74	80	26
300.01	51	14	37	*	25	14	12	15	11	17	*7
300.02	13	*	*8	*	*	*5	*	*	*	*7	*
300.1	29	*7	22	*	20	*6	*	*8	*7	9	*5
300.11	12	*	9	*	*8	*	*	*	*	*	*
300.14	*6	*	*6	*	*	*	—	*	*	*	*
300.15	*6	*	*	*	*	*	—	*	*	*	*
300.2	22	*8	13	*	16	*	*	*6	*7	*7	*
300.21	14	*	9	—	10	*	*	*	*	*	*
300.3	17	*9	*9	*	10	*	*	*5	*	*6	*
300.4	149	51	98	*	57	44	45	32	48	56	14
300.8	*7	*	*6	*	*	*	*	*	*	*	*
300.81	*6	*	*5	*	*	*	*	*	*	*	*
300.9	63	31	32	*	42	14	*	15	20	14	14
301	303	135	168	*5	224	55	19	103	93	71	36
301.0	*5	*	*	—	*	*	*	*	*	*	*
301.2	12	9	*	—	10	*	*	*	*	*	*
301.20	*6	*	*	—	*	*	*	*	*	*	*
301.22	*6	*	*	—	*5	*	*	*	*	*	*
301.5	9	*	*9	*	*	*	*	*	*	*	*
301.50	*9	*	*8	*	*	*	*	*	*	*	*
301.6	39	17	22	*	24	*9	*6	14	13	*8	*
301.7	21	18	*	*	17	*	*	*5	10	*	*
301.8	137	47	89	*	111	22	*	51	45	27	13
301.81	*6	*	*	*	*	*	*	*	*	*	*
301.82	*5	*	*	*	*	*	—	*	*	*	*
301.83	81	19	63	*	69	11	*	22	33	17	*9
301.89	39	20	19	*	30	*7	*	26	*	*	*
301.9	70	34	36	*	50	12	*6	23	17	19	10
302	*5	*	*	*	*	*	*	*	*	*	*
303	772	581	191	*	426	251	93	189	212	244	127
303.0	113	89	24	*	66	38	*8	26	42	26	19
303.00	34	26	*8	—	22	11	*	*	13	11	*6
303.01	77	61	15	*	43	27	*6	20	28	15	13
303.9	659	492	167	*	360	213	85	164	170	217	108
303.90	260	194	66	*	130	87	42	37	70	107	46
303.91	346	263	82	*	208	104	33	112	87	99	47
303.92	11	*8	*	—	*6	*	*	*5	*	*	*
303.93	43	27	16	*	16	17	9	*9	11	10	13

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
304	398	246	152	*	317	65	13	144	90	108	57
304.0	104	64	40	—	76	25	*	50	16	16	22
304.00	29	16	13	—	20	*8	*	*	*6	*8	11
304.01	70	44	26	—	54	14	*	44	*9	*7	10
304.1	21	11	10	—	11	*5	*	*7	*	*6	*
304.10	*8	*	*	—	*	*	*	*	*	*	*
304.11	12	*6	*5	—	*6	*	*	*	*	*	*
304.2	104	68	36	*	91	12	*	40	23	34	*8
304.20	33	19	13	*	27	*	*	*	*8	17	*
304.21	67	46	21	—	59	*7	*	33	14	16	*
304.3	35	24	11	*	30	*	—	*8	20	*6	*
304.30	*8	*5	*	*	*7	*	—	*	*	*	*
304.31	24	17	*7	*	21	*	—	*6	15	*	*
304.4	9	*6	*	—	*9	*	*	*	*	*	*
304.40	*6	*	*	—	*	*	*	*	*	*	*
304.6	11	*5	*6	—	*7	*	*	*	*	*	*
304.61	*7	*	*	—	*	*	*	*	*	*	*
304.7	41	25	16	—	36	*	*	20	*6	*9	*6
304.70	*8	*	*	—	*8	*	—	*	*	*	*
304.71	32	19	13	—	28	*	*	18	*	*6	*
304.8	43	28	15	*	36	*5	*	*7	*6	23	*7
304.80	16	10	*6	*	14	*	*	*	*	9	*
304.81	22	15	*7	*	20	*	*	*5	*	13	*
304.9	28	15	13	*	19	*6	*	*8	*6	10	*
304.90	19	11	*9	*	14	*	*	*6	*6	*7	*
304.91	*7	*	*	—	*	*	*	*	*	*	*
305	1,450	827	623	10	804	443	193	298	360	505	288
305.0	326	229	96	*	191	94	37	89	74	115	48
305.00	204	142	61	*	120	57	24	45	49	77	32
305.01	86	65	21	*	47	28	11	27	20	30	10
305.02	17	11	*6	*	13	*	*	10	*	*	*
305.03	19	11	*8	*	11	*	*	*7	*	*	*
305.1	724	372	352	*	274	296	152	108	175	273	167
305.2	84	51	32	*	77	*	*	22	28	21	12
305.20	55	34	21	*	50	*	*	10	21	16	*7
305.21	18	12	*6	*	17	*	—	*7	*	*	*
305.22	*6	*	*	—	*6	*	—	*	*	*	*
305.4	10	*	*6	*	*6	*	*	*	*	*	*
305.40	*7	*	*	*	*	*	*	*	*	*	*
305.5	35	21	14	—	25	10	*	9	10	9	*7
305.50	23	14	*9	—	16	*6	*	*	*7	*6	*5
305.51	9	*6	*	—	*7	*	*	*	*	*	*
305.6	140	78	62	*	123	17	*	38	38	47	17
305.60	88	51	37	—	76	12	*	17	28	30	12
305.61	38	19	18	*	34	*	—	16	*7	12	*
305.62	*6	*	*	—	*6	*	—	*	*	*	*
305.63	*8	*5	*	—	*7	*	*	*	*	*	*
305.7	13	*	9	—	12	*	*	*	*	*	*7
305.70	10	*	*7	—	10	*	—	*	*	*	*6
305.9	114	64	49	*	92	17	*	27	28	33	25
305.90	78	44	34	*	64	10	*	12	22	28	16
305.91	16	9	*7	*	13	*	*	*7	*	*	*
305.92	*5	*	*	—	*5	*	—	*	*	—	*
305.93	14	*7	*8	*	10	*	*	*	*	*	*8
306	9	*	*5	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
307	59	17	42	*5	35	*6	13	13	17	19	10
307.2	*5	*	*	*	*	*	*	*	*	*	*
307.5	16	*	14	*	14	*	*	*	*	*	*
307.51	9	*	9	*	9	*	—	*	*	*	*
307.8	14	*	12	*	*8	*	*	*	*	*6	*
307.81	11	*	10	*	*6	*	*	*	*	*	*
307.9	15	*8	*7	*	*	*	*9	*	*6	*	*
308	21	*6	15	*	*8	*7	*	*5	*7	*	*
308.3	*8	*	*7	*	*	*	*	*	*	*	*
308.9	*8	*	*6	*	*	*	*	*	*	*	*
309	245	101	144	19	165	36	25	74	77	58	35
309.0	78	34	44	*	55	10	10	23	20	23	13
309.1	*5	*	*	*	*	*	*	*	*	*	*
309.2	38	14	23	*	21	*6	*7	11	12	11	*
309.24	*6	*	*	*	*	*	*	*	*	*	*
309.28	29	11	19	*	18	*6	*	*8	11	*8	*
309.3	*6	*	*	*	*	*	—	*	*	*	*
309.4	29	16	14	*	20	*	*	12	*7	*6	*
309.8	67	21	46	*7	47	12	*	22	25	10	10
309.81	64	19	45	*7	45	12	*	21	24	10	10
309.9	22	*8	14	*	14	*	*	*5	*8	*6	*
310	98	39	59	*	9	*6	83	20	20	43	14
310.1	*9	*	*	—	*	*	*	*	*	*	*
310.2	*6	*	*	*	*	*	*	*	*	*	*
310.9	82	31	51	—	*	*	77	16	16	40	10
311	507	180	327	13	180	115	198	129	147	161	70
312	60	42	18	19	34	*	*	22	19	13	*7
312.3	30	20	10	*6	21	*	*	14	*9	*6	*
312.30	14	*9	*	*	10	*	*	9	*	*	*
312.34	11	*8	*	*	*7	*	*	*	*	*	*
312.8	*8	*5	*	*	*	*	*	*	*	*	*
312.9	17	14	*	*6	9	*	*	*	*8	*	*
313	24	13	10	14	9	—	*	*5	*8	*7	*
313.8	22	13	9	14	*9	—	—	*	*8	*7	*
313.81	22	12	9	13	*8	—	—	*	*8	*7	*
314	53	40	13	34	18	*	*	11	22	12	*7
314.0	50	38	12	34	16	*	*	11	22	12	*5
314.00	14	9	*	*7	*6	*	*	*	*	*5	*
314.01	36	28	*8	26	9	*	—	*8	19	*7	*
315	22	14	*8	15	*5	*	*	*	11	*	*
315.9	*7	*	*	*	*	*	*	*	*	*	*
317	28	13	15	*	14	*7	*	10	*7	*9	*
318	28	12	16	*	16	*	*	*5	10	9	*
318.0	*8	*	*	*	*	*	*	*	*	*	*
318.1	12	*6	*7	*	*5	*	*	*	*6	*	*
318.2	*8	*	*5	*	*6	*	*	*	*	*	*
319	66	40	25	13	23	16	14	15	17	23	11
320–389	3,456	1,567	1,889	416	572	634	1,834	785	842	1,167	662
320	13	*7	*6	*7	*	*	*	*	*	*	*
322	12	*6	*6	*	*	*	*	*	*	*	*
322.9	9	*5	*	*	*	*	*	*	*	*	*
323	10	*	*6	*	*	*	*	*	*	*	*
323.9	*7	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
324	*6	*	*	*	*	*	*	*	*	*	*
331	333	124	209	19	14	19	282	70	83	126	54
331.0	245	83	161	*	*	*5	238	50	58	100	36
331.3	14	*	*9	*	*	*	*8	*	*	*5	*
331.4	54	27	27	17	9	*9	18	12	15	15	12
331.9	16	*7	*9	*	*	*	13	*	*5	*	*
332	212	104	108	*	*	12	199	47	52	80	33
332.0	211	104	107	*	*	12	198	46	52	79	33
333	64	32	32	*	*8	14	41	11	15	24	14
333.0	*6	*	*	*	—	*	*5	*	*	*	*
333.1	21	12	9	—	*	*6	14	*	*6	*7	*
333.8	10	*	*7	—	*	*	*6	*	*	*	*
333.82	9	*	*7	—	*	*	*6	*	*	*	*
333.9	15	*7	*8	*	*	*	*9	*	*	*8	*
333.99	11	*	*6	*	*	*	*7	*	*	*	*
334	*5	*	*	*	*	*	*	*	*	*	*
335	12	*7	*	*	*	*	*7	*	*	*	*
335.2	11	*6	*	*	*	*	*7	*	*	*	*
335.20	*6	*	*	—	*	*	*	*	*	*	*
336	20	12	*8	*	*	*7	*7	*5	*6	*6	*
336.3	*6	*	*	*	*	*	*	*	*	*	*
336.9	*6	*	*	*	*	*	*	*	*	*	*
337	57	22	35	*	19	16	22	14	11	19	13
337.1	39	17	22	*	*9	14	16	*8	*7	14	10
337.2	12	*	10	—	10	*	*	*	*	*	*
340	75	25	50	—	24	32	18	20	21	21	13
341	*8	*6	*	*	*	*	*	*	*	*	*
342	467	215	253	*	24	83	356	91	126	154	97
342.0	16	*7	9	*	*	*	12	*	*5	*	*
342.00	11	*	*6	*	*	*	*7	*	*	*	*
342.1	*6	*	*	*	*	*	*	*	*	*	*
342.8	16	10	*6	*	*	*	10	*	*6	*6	*
342.80	11	*7	*	*	*	*	*6	*	*	*5	*
342.9	429	195	234	*	19	77	330	86	112	142	89
342.90	371	165	206	*	16	66	286	73	97	124	77
342.91	35	18	18	*	*	*7	26	*9	*8	11	*7
342.92	23	12	10	*	*	*	17	*	*7	*7	*5
343	65	35	30	32	22	9	*	18	15	18	13
343.2	11	*6	*	*6	*	*	*	*	*	*	*
343.9	43	24	19	18	16	*6	*	9	11	13	9
344	131	82	49	*	45	34	49	25	34	46	27
344.0	46	33	14	*	18	14	13	*7	11	17	12
344.00	33	24	*9	*	12	11	10	*	*8	12	*8
344.04	*5	*	*	—	*	*	*	*	*	*	*
344.1	44	29	16	*	20	10	14	10	11	15	*9
344.6	12	*7	*6	*	*	*	*	*	*	*	*
344.60	*7	*	*	*	*	*	*	*	*	*	*
344.61	*5	*	*	*	*	*	*	*	*	*	*
344.9	16	*8	*8	*	*	*	11	*	*5	*6	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
345	115	53	62	31	42	18	24	30	28	40	17
345.1	24	10	13	*5	9	*	*5	*6	*7	*9	*
345.10	21	9	12	*	*8	*	*	*	*6	*8	*
345.3	22	11	11	*6	*6	*	*7	*	*5	9	*
345.4	9	*	*6	*	*6	*	*	*	*	*	*
345.40	*6	*	*	*	*	*	*	*	*	*	*
345.5	*7	*	*	*	*	*	*	*	*	*	*
345.50	*6	*	*	*	*	*	*	*	*	*	*
345.9	41	20	21	11	16	*8	*6	10	11	13	*7
345.90	37	18	19	10	13	*7	*6	*9	10	11	*6
346	118	24	94	*	70	36	10	29	26	45	18
346.8	10	*	*7	*	*	*	*	*	*	*	*
346.80	*7	*	*	*	*	*	*	*	*	*	*
346.9	96	19	77	*	58	28	*8	22	22	37	15
346.90	85	18	67	*	53	22	*8	20	20	31	14
346.91	11	*	10	—	*	*6	*	*	*	*6	*
348	161	83	78	15	37	42	68	33	37	58	32
348.0	*7	*	*	*	*	*	*	*	*	*	*
348.1	56	30	26	*	9	20	23	11	13	20	11
348.2	*7	*	*6	*	*	*	*	*	*	*	*
348.3	49	25	24	*	*6	*8	31	10	10	22	*7
348.4	*6	*	*	*	*	*	*	*	*	*	*
348.5	11	*6	*	*	*	*	*	*	*	*	*
348.8	20	11	9	*	*9	*	*	*	*	*7	*5
348.9	*5	*	*	*	*	*	*	*	*	*	*
349	50	14	35	*	29	*9	9	12	10	20	*7
349.0	26	*	22	*	22	*	*	*6	*6	10	*
349.2	*8	*	*	*	*	*	*	*	*	*	*
349.8	13	*6	*6	*	*	*	*5	*	*	*6	*
349.82	*7	*	*	*	*	*	*	*	*	*	*
350	13	*	*8	—	*	*	*8	*	*	*	*
350.1	11	*	*7	—	*	*	*8	*	*	*	*
351	27	12	15	*	*6	*8	13	*	*8	*8	*7
351.0	24	12	12	*	*6	*7	11	*	*8	*6	*6
353	11	*6	*6	*	*5	*	*	*	*	*7	*
353.0	*7	*	*	*	*5	*	*	*	*	*	*
354	29	13	15	*	*7	10	10	10	*5	*9	*
354.0	18	*7	11	*	*	*7	*6	*6	*	*6	*
354.2	*5	*	*	*	*	*	*	*	*	*	*
355	37	15	22	*	*8	*9	20	10	11	9	*7
355.8	16	*6	10	*	*	*	10	*	*6	*	*
355.9	11	*	*7	*	*	*	*	*	*	*	*
356	41	23	18	*	*	*8	27	*8	9	16	*9
356.9	36	20	16	*	*	*7	23	*7	*8	14	*8
357	168	87	81	*	18	60	90	37	41	58	32
357.0	10	*7	*	*	*	*	*	*	*	*	*
357.2	146	74	72	*	13	53	79	34	34	51	28
358	20	11	10	*	*	*	13	*	*7	*7	*
358.0	14	*7	*7	*	*	*	*9	*	*5	*5	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
359	29	15	13	*	12	*8	*7	*	*	10	*9
359.1	10	*8	*	*	*6	*	*	*	*	*6	*
359.4	*6	*	*	*	*	*	*	*	*	*	*
360	*6	*	*	*	*	*	*	*	*	*	*
361	20	10	9	*	*	*5	9	*	*	12	*
361.0	12	*6	*6	*	*	*	*6	*	*	*7	*
361.00	*5	*	*	—	*	*	*	*	*	*	*
362	156	63	93	*7	19	45	85	33	36	54	33
362.0	90	41	50	—	14	36	40	23	18	29	20
362.01	85	38	47	—	13	34	38	21	18	26	20
362.02	*5	*	*	—	*	*	*	*	*	*	*
362.1	*6	*	*	*	*	*	*	*	*	*	*
362.2	10	*	*7	*5	*	*	*	*	*	*6	*
362.21	*5	*	*	*5	—	*	*	*	*	*	*
362.3	14	*	*9	—	*	*	9	*	*	*7	*
362.34	*8	*	*	—	*	*	*	*	*	*	*
362.5	29	*7	22	—	*	*	27	*	12	*6	*6
362.50	25	*6	19	—	*	*	24	*	12	*	*6
363	12	*8	*	*	*8	*	*	*	*	*	*
363.2	9	*6	*	*	*6	*	*	*	*	*	*
363.20	*9	*6	*	*	*6	*	*	*	*	*	*
365	175	62	113	*	*	18	155	52	46	50	26
365.9	158	58	100	*	*	15	141	43	44	47	23
366	49	21	28	*	*	*6	40	14	*6	17	11
366.1	10	*	*6	—	*	*	*9	*	*	*	*
366.8	*5	*	*	—	*	*	*	*	*	*	*
366.9	31	13	18	*	*	*	27	9	*	14	*
368	45	20	26	*	12	10	20	*9	12	15	10
368.1	*6	*	*	*	*	*	*	*	*	*	*
368.2	12	*5	*6	*	*	*	*5	*	*	*	*
368.4	16	*5	11	*	*	*	10	*	*	*6	*
368.46	13	*	*8	*	*	*	*8	*	*	*	*
368.8	*8	*	*	*	*6	*	*	*	*	*	*
369	91	32	59	*	11	13	64	18	15	30	28
369.0	39	13	26	*	*	*6	27	*7	*6	13	12
369.00	38	13	25	*	*	*6	26	*7	*6	13	12
369.4	20	*7	13	*	*	*	14	*6	*	*6	*
369.6	14	*6	*8	*	*	*	11	*	*	*6	*
369.60	13	*6	*7	*	*	*	10	*	*	*5	*
369.9	10	*	*7	*	*	*	*7	*	*	*	*
370	*5	*	*	*	*	*	*	*	*	*	*
371	*5	*	*	*	*	*	*	*	*	*	*
372	46	25	22	13	10	*6	17	*9	12	17	*9
372.0	*9	*6	*	*	*	*	*	*	*	*	*
372.00	*7	*	*	*	*	*	*	*	*	*	*
372.3	25	12	14	9	*	*	9	*5	*7	10	*
372.30	24	11	13	*9	*	*	*9	*	*7	10	*
372.7	*7	*5	*	*	*	*	*	*	*	*	*
372.72	*7	*	*	*	*	*	*	*	*	*	*
373	11	*5	*6	*	*	*	*	*5	*	*	*
373.0	*5	*	*	*	*	*	*	*	*	*	*
373.1	*5	*	*	*	*	*	*	*	*	*	*
374	13	*	9	*	*	*	*6	*	*	*	*
374.3	*8	*	*6	*	*	*	*	*	*	*	*
374.30	*7	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
376	17	*8	9	*7	*	*	*	*	*	*5	*
376.0	11	*6	*	*6	*	*	*	*	*	*	*
376.01	11	*6	*	*6	*	*	*	*	*	*	*
377	12	*	*8	*	*	*	*	*	*	*6	*
377.3	*5	*	*	*	*	*	*	*	*	*	*
378	17	*8	*9	*	*	*	*	*6	*	*	*
378.5	10	*5	*	*	*	*	*	*	*	*	*
379	16	*9	*8	*	*	*	*5	*	*	*	*
379.2	*6	*	*	*	*	*	*	*	*	*	*
380	23	11	12	*5	*6	*	9	*6	*6	*8	*
380.1	15	*7	*8	*	*	*	*5	*	*	*6	*
380.10	13	*6	*7	*	*	*	*	*	*	*	*
380.4	*7	*	*	*	*	*	*	*	*	*	*
381	29	18	11	23	*	*	*	12	*6	*8	*
381.0	*8	*	*	*7	*	*	*	*	*	*	*
381.1	*9	*5	*	*8	*	*	*	*	*	*	*
381.10	*9	*5	*	*8	*	*	*	*	*	*	*
381.4	*7	*	*	*	*	*	*	*	*	*	*
382	201	108	93	177	13	*6	*	43	49	63	46
382.9	192	103	88	171	12	*6	*	41	46	61	43
383	*7	*	*	*	*	*	*	*	*	*	*
384	*5	*	*	*	*	*	*	*	*	*	*
385	*8	*5	*	*	*	*	*	*	*	*	*
386	65	21	43	*	*9	16	39	15	19	22	*9
386.0	17	*6	11	—	*	*	*8	*	*5	*7	*
386.00	12	*	9	—	*	*	*7	*	*5	*	*
386.1	16	*	12	*	*	*6	9	*7	*	*	*
386.11	9	*	*7	*	*	*	*6	*	*	*	*
386.3	28	10	18	*	*	*6	20	*6	*9	10	*
386.30	24	*8	16	*	*	*5	17	*5	*8	*8	*
388	14	9	*	*	*	*	*5	*	*	*	*
389	67	38	29	*6	10	*9	43	15	21	16	14
389.1	9	*6	*	*	*	*	*	*	*	*	*
389.10	*6	*	*	*	*	*	*	*	*	*	*
389.9	49	28	22	*	*6	*5	36	*9	17	13	11
390–459	26,992	13,151	13,841	116	1,561	6,391	18,923	6,148	6,710	9,677	4,458
394	37	*9	28	*	*	11	23	*8	10	13	*7
394.0	17	*	14	*	*	*5	9	*	*	*5	*
394.1	*6	*	*	—	*	*	*	*	*	*	*
394.2	*6	*	*5	—	*	*	*	*	*	*	*
394.9	*9	*	*	—	*	*	*6	*	*	*5	*
396	178	72	106	*	*	18	156	46	40	71	21
396.0	10	*	*7	*	*	*	*8	*	*	*	*
396.1	*6	*	*	*	*	*	*	*	*	*	*
396.2	47	18	29	*	—	*	43	12	12	18	*
396.3	90	39	51	*	*	10	79	22	19	42	*7
396.8	18	*8	11	—	*	*	17	*7	*	*	*
396.9	*7	*	*5	—	—	*	*6	*	*	*	*
397	114	41	74	*	*5	21	88	32	26	46	11
397.0	114	41	73	*	*5	20	87	32	26	45	11

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
398	36	14	22	*	*	9	25	16	*7	*7	*6
398.9	36	14	22	*	*	9	25	16	*7	*7	*6
398.90	15	*7	*8	*	*	*7	*7	*	*	*	*
398.91	21	*7	13	—	*	*	19	12	*	*	*
401	5,152	2,188	2,964	*9	401	1,456	3,286	1,138	1,354	1,816	843
401.0	73	25	47	*	14	23	35	12	14	36	11
401.1	174	74	100	*	10	52	111	107	14	20	33
401.9	4,905	2,089	2,816	*8	377	1,381	3,139	1,019	1,327	1,760	799
402	388	137	251	*	11	68	308	92	82	147	66
402.0	12	*	10	—	*	*	*7	*	*	*9	*
402.01	*7	*	*6	—	*	*	*	*	*	*5	*
402.1	10	*	*7	—	*	*	*8	*5	*	*	*
402.10	*6	*	*	—	*	*	*	*	*	*	*
402.9	366	131	235	*	*8	65	292	85	80	136	64
402.90	188	71	117	*	*	41	143	48	40	69	32
402.91	177	60	117	*	*	24	149	38	41	67	32
403	520	257	262	*	79	159	280	103	99	211	107
403.0	14	*5	*8	—	*	*	*6	*	*	*8	*
403.01	10	*	*6	—	*	*	*	*	*	*7	*
403.1	9	*	*	—	*	*	*7	*	*	*	*
403.11	*7	*	*	—	*	*	*6	*	*	*	*
403.9	497	248	249	*	74	154	266	98	95	200	104
403.90	39	19	20	*	*	10	25	9	*8	17	*
403.91	458	228	229	*	70	144	242	88	87	183	99
404	103	43	60	—	*9	28	66	22	21	38	22
404.0	*7	*	*	—	*	*	*	*	*	*	*
404.9	95	41	54	—	*7	27	61	22	19	34	21
404.91	10	*5	*	—	*	*	*7	*	*	*5	*
404.92	18	*7	11	—	*	*5	9	*	*	*6	*
404.93	64	26	38	—	*	17	43	16	12	21	15
405	15	*	11	*	*	*	*8	*	*	*	*
405.9	13	*	10	*	*	*	*7	*	*	*	*
405.91	9	*	*8	*	*	*	*7	*	*	*	*
410	919	548	371	*	44	302	572	212	227	321	159
410.0	37	22	15	—	*	14	22	9	11	11	*6
410.01	34	21	13	—	*	12	20	*9	10	10	*5
410.1	163	99	64	—	9	63	91	39	36	61	27
410.11	143	88	55	—	*9	55	80	36	34	52	22
410.12	15	*9	*6	—	*	*5	*9	*	*	*5	*
410.2	20	10	10	—	*	*7	12	*5	*	*7	*
410.21	19	10	9	—	*	*6	12	*5	*	*7	*
410.3	21	14	*7	—	*	*8	12	*6	*	*6	*
410.31	20	13	*7	—	*	*7	11	*5	*	*6	*
410.4	194	122	73	*	13	77	103	42	54	68	30
410.41	177	108	69	*	13	71	92	38	49	63	26
410.42	15	12	*	—	*	*6	*9	*	*	*	*
410.5	22	12	10	—	*	*8	14	*7	*	*	*6
410.51	20	10	10	—	*	*8	12	*6	*	*	*6
410.6	*6	*	*	—	*	*	*	*	*	*	*
410.61	*5	*	*	—	*	*	*	*	*	*	*
410.7	307	178	129	*	12	87	207	74	76	101	57
410.70	*7	*	*	—	—	*	*5	*	*	*	*
410.71	279	163	116	*	11	79	188	67	72	88	51
410.72	21	12	*9	—	*	*6	14	*5	*	*8	*
410.8	26	15	10	—	*	*5	20	*8	*	10	*
410.81	24	14	10	—	*	*	18	*7	*	10	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
410.9	123	72	52	*	*	31	88	21	31	50	22
410.90	21	12	9	—	*	*8	13	*	*	*8	*
410.91	94	55	39	*	*	21	70	15	24	39	16
410.92	*8	*	*	—	*	*	*5	*	*	*	*
411	1,072	592	480	*	52	376	642	274	233	381	184
411.0	9	*6	*	—	*	*	*5	*	*	*	*
411.1	987	545	442	*	48	353	586	251	214	352	170
411.8	76	41	35	*	*	20	52	21	18	27	11
411.81	30	22	*8	*	*	12	15	*7	*9	*8	*5
411.89	46	19	27	*	*	*8	36	13	*9	19	*6
412	617	390	228	*	26	189	402	153	164	195	106
413	590	286	304	*	29	162	399	148	159	210	75
413.1	*6	*	*	—	*	*	*	*	*	*	*
413.9	583	282	300	*	27	160	395	145	158	207	74
414	4,144	2,339	1,804	*6	120	1,105	2,913	1,010	1,051	1,437	646
414.0	3,862	2,166	1,696	*	107	1,046	2,705	927	975	1,348	612
414.00	1,198	614	584	*	23	250	923	266	332	468	133
414.01	2,532	1,453	1,080	*	81	752	1,695	637	606	835	454
414.02	130	98	32	—	*	43	85	24	36	45	25
414.1	13	*8	*	*	*	*	*8	*	*	*	*
414.10	*9	*5	*	—	*	*	*6	*	*	*	*
414.8	175	114	61	*	*9	38	128	56	44	48	27
414.9	94	51	43	*	*	18	73	24	29	37	*
415	112	47	65	*	16	27	69	27	31	29	25
415.0	*7	*	*	—	*	*	*	*	*	*	*
415.1	105	44	61	*	15	24	65	25	28	28	24
415.11	16	*8	*8	—	*	*	9	*	*	*	*6
415.19	89	36	52	*	13	20	55	22	25	24	17
416	189	77	112	*	16	49	120	43	37	68	41
416.0	102	38	64	*	10	26	63	20	18	43	21
416.8	31	13	18	*	*	*8	18	10	*6	*6	9
416.9	56	26	30	*	*	15	39	13	13	19	11
420	30	19	12	*	*8	*8	13	*6	*7	11	*6
420.9	26	15	11	*	*7	*8	11	*6	*7	*9	*
420.90	18	11	*8	*	*	*	*9	*	*	*6	*
420.91	*7	*	*	*	*	*	*	*	*	*	*
421	25	13	12	*	*6	*5	13	*8	*	*7	*6
421.0	23	12	11	*	*6	*	12	*7	*	*7	*5
423	40	22	18	*	*9	14	16	*8	*8	15	*9
423.9	32	17	15	*	*7	11	13	*7	*6	12	*7
424	847	324	523	*5	91	161	590	196	198	295	158
424.0	525	176	349	*	81	117	325	116	123	193	93
424.1	269	129	140	*	*7	33	227	69	65	83	52
424.2	27	10	17	*	*	*6	19	*6	*6	*9	*6
424.3	*7	*	*	*	*	*	*	*	*	*	*
424.9	20	*7	13	*	*	*	14	*	*	*7	*6
424.90	19	*6	12	*	*	*	13	*	*	*7	*5
425	427	241	186	*	38	128	258	93	100	156	78
425.1	20	*7	13	*	*	*6	12	*7	*	*	*5
425.4	365	210	155	*	31	106	225	75	87	138	66
425.5	13	10	*	—	*	*7	*	*	*	*	*
425.8	20	11	*9	*	*	*7	12	*5	*	*9	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
426	582	317	264	*5	30	99	447	122	136	219	105
426.0	73	41	32	*	*	9	61	18	15	25	16
426.1	166	94	72	*	*6	23	136	35	44	59	29
426.10	*8	*	*	*	*	*	*6	*	*	*	*
426.11	105	55	50	*	*	15	87	21	27	38	20
426.12	*8	*	*	—	*	*	*6	*	*	*	*
426.13	45	30	16	*	*	*5	37	10	13	15	*7
426.2	44	27	18	—	*	*7	34	*9	9	21	*5
426.3	100	41	59	—	*	18	80	20	23	39	17
426.4	107	61	46	*	10	24	72	22	22	44	19
426.5	47	32	15	*	*	*8	39	*8	13	18	*8
426.52	15	10	*	—	*	*	13	*	*	*6	*
426.53	24	15	*9	*	*	*	20	*	*8	*9	*
426.6	*9	*	*	—	*	*	*	*	*	*	*
426.7	13	*8	*	*	*	*	*	*	*	*	*
426.8	*7	*	*	*	*	*	*	*	*	*	*
426.89	*7	*	*	*	*	*	*	*	*	*	*
426.9	15	*7	*9	*	*	*	14	*	*	*8	*
427	3,383	1,696	1,687	24	157	592	2,609	792	819	1,181	591
427.0	106	41	66	*	9	25	70	22	23	42	19
427.1	224	150	74	*	13	62	148	62	54	71	37
427.2	*6	*	*	*	*	*	*	*	*	*	*
427.3	1,993	975	1,018	*	37	278	1,674	487	499	667	339
427.31	1,814	874	941	*	30	238	1,543	445	453	609	308
427.32	178	102	77	*	*7	39	131	43	47	58	31
427.4	67	43	24	*	*	23	39	13	15	25	14
427.41	65	42	23	*	*	22	38	13	15	24	13
427.5	125	66	59	*	10	28	83	18	25	55	27
427.6	189	101	88	*	13	39	135	43	42	72	33
427.61	51	22	30	*	*	*8	39	13	*9	22	*8
427.69	136	78	58	*	9	31	94	30	32	48	25
427.8	591	275	317	11	62	120	399	133	139	210	109
427.81	132	60	73	*	*	13	117	30	29	54	19
427.89	459	215	244	11	59	107	282	103	110	156	90
427.9	82	44	38	*	*7	16	58	12	21	35	13
428	2,872	1,256	1,616	14	79	449	2,330	603	770	1,085	414
428.0	2,844	1,241	1,603	14	78	444	2,308	597	764	1,073	410
428.1	18	10	*8	*	*	*	16	*	*5	*7	*
428.9	10	*6	*	*	*	*	*6	*	*	*5	*
429	334	155	179	*	19	52	261	73	77	133	52
429.2	178	81	97	—	*	21	153	40	40	76	22
429.3	112	48	64	*	*8	22	81	23	26	41	22
429.4	11	*8	*	*	*	*	*6	*	*	*	*
429.9	18	11	*7	*	*	*	12	*	*6	*6	*
430	27	*7	19	*	*7	12	*8	*	*6	*9	*7
431	80	36	44	*	*7	18	55	14	19	33	14
432	31	19	12	*	*	*6	21	*	*	11	11
432.1	27	17	11	*	*	*	19	*	*	9	10
433	329	181	148	*	*6	66	256	86	83	116	44
433.1	240	131	108	*	*	48	188	64	62	83	30
433.10	214	119	95	*	*	42	168	58	58	72	26
433.11	26	12	14	—	*	*6	19	*6	*	11	*
433.3	77	42	35	—	*	14	61	19	18	28	11
433.30	66	37	30	—	*	12	53	17	14	25	10
433.31	10	*5	*5	—	*	*	*8	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
434	386	178	208	*	13	68	304	89	92	134	71
434.0	20	*7	13	*	*	*	16	*6	*	*7	*
434.01	17	*6	11	*	*	*	14	*6	*	*	*
434.1	55	24	31	—	*	9	45	14	12	14	15
434.10	9	*	*	—	*	*	*8	*	*	*	*
434.11	46	20	26	—	*	*8	37	12	9	11	15
434.9	311	147	165	*	11	56	243	69	75	114	53
434.90	10	*6	*	*	*	*	*8	*	*	*	*
434.91	301	141	160	*	10	54	235	64	73	111	52
435	254	103	151	*	11	48	195	55	68	93	38
435.1	*6	*	*	—	*	*	*5	*	*	*	*
435.3	14	*7	*6	—	*	*	10	*	*	*	*
435.8	10	*	*6	—	*	*	*7	*	*	*	*
435.9	218	88	130	*	10	42	165	48	56	81	34
436	212	88	124	*	*6	31	175	28	51	90	43
437	133	51	82	*	*6	20	106	26	34	53	20
437.0	58	23	35	—	*	*	54	10	14	23	10
437.1	15	*6	10	*	*	*	13	*	*5	*6	*
437.2	10	*	*7	*	*	*	*	*	*	*	*
437.3	13	*6	*7	*	*	*	*5	*	*	*	*
437.9	32	11	21	—	*	*	27	*8	*8	16	*
438	428	197	231	*	14	75	338	82	115	167	63
440	468	244	224	*	*9	98	361	95	115	188	70
440.0	25	11	14	—	*	*6	18	*	*5	*9	*7
440.1	21	10	11	*	*	*6	14	*	*	*9	*
440.2	269	150	119	—	*7	65	198	56	67	103	43
440.20	83	43	40	—	*	18	64	13	22	35	13
440.21	44	30	14	—	*	14	29	9	9	15	10
440.22	19	10	*8	—	*	*	13	*5	*	*6	*
440.23	38	19	19	—	*	10	27	*9	10	13	*7
440.24	78	44	34	—	*	16	59	19	19	32	*9
440.29	*7	*	*	—	*	*	*	*	*	*	*
440.3	10	*5	*	—	*	*	*6	*	*	*	*
440.31	*6	*	*	—	*	*	*	*	*	*	*
440.9	139	66	73	—	*	16	122	28	34	62	15
441	160	105	55	*	*	22	136	38	38	60	25
441.0	15	*8	*6	—	*	*	10	*	*	*	*
441.01	*8	*	*	—	*	*	*5	*	*	*	*
441.2	13	*	*9	*	*	*	10	*	*	*	*
441.3	10	*8	*	—	*	*	*8	*	*	*	*
441.4	116	80	36	*	*	12	103	26	27	45	17
442	42	29	13	*	*	*9	30	*9	10	13	11
442.2	10	*9	*	—	*	*	*8	*	*	*	*
442.3	21	16	*	*	*	*	15	*	*	*6	*6
442.8	*8	*	*	*	*	*	*5	*	*	*	*
442.81	*5	*	*	—	*	*	*	*	*	*	*
443	398	215	182	*	15	90	291	103	85	148	62
443.0	*9	*	*5	—	*	*	*	*	*	*	*
443.8	104	56	48	*	*	30	68	25	20	40	20
443.81	92	51	41	*	*	28	58	22	15	35	19
443.89	12	*	*7	—	*	*	10	*	*	*	*
443.9	283	155	129	*	*6	57	220	74	63	106	41

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
444	118	65	53	*	11	31	75	25	23	46	24
444.0	10	*5	*	*	*	*	*5	*	*	*	*
444.2	82	46	37	*	*8	19	55	18	18	31	15
444.21	14	*5	*9	—	*	*	*8	*	*	*7	*
444.22	69	40	28	*	*	16	48	17	15	25	12
444.8	19	12	*7	*	*	*6	10	*	*	*8	*
444.81	14	*9	*5	*	*	*	*7	*	*	*6	*
446	31	11	19	*	*	*	20	*7	*8	*7	*8
446.4	*5	*	*	—	*	*	*	*	*	*	*
446.5	13	*	10	—	*	*	13	*	*	*	*
447	51	25	26	*	*	19	26	14	9	20	*9
447.1	23	12	12	*	*	*9	14	*7	*	*7	*
447.6	14	*7	*7	*	*	*	*6	*	*	*6	*
447.8	*5	*	*	*	*	*	*	*	*	*	*
448	*6	*	*	*	*	*	*	*	*	*	*
451	99	42	57	*	17	27	54	20	28	36	16
451.0	18	*7	11	*	*	*7	*8	*	*8	*	*
451.1	38	15	23	*	*7	9	22	9	9	13	*6
451.11	13	*	9	*	*	*	*7	*	*	*5	*
451.19	25	11	14	*	*	*7	15	*6	*7	*8	*
451.2	16	*	11	—	*	*	10	*	*	*7	*
451.8	21	12	10	*	*6	*	11	*	*	*8	*5
451.82	*7	*	*	*	*	*	*	*	*	*	*
451.84	*7	*	*	*	*	*	*	*	*	*	*
451.89	*6	*	*	*	*	*	*	*	*	*	*
451.9	*6	*	*	—	*	*	*	*	*	*	*
453	270	122	148	*	46	70	152	57	76	87	50
453.8	249	114	136	*	42	64	142	52	73	80	45
453.9	16	*6	10	*	*	*	*8	*	*	*6	*
454	50	20	30	*	*9	11	29	12	15	13	9
454.0	11	*	*7	*	*	*	*6	*	*	*	*
454.1	*5	*	*	—	*	*	*	*	*	*	*
454.2	14	*6	*8	—	*	*	9	*	*	*	*
454.9	19	*8	11	—	*	*	10	*	*5	*	*
455	142	64	78	*	35	38	69	33	33	58	19
455.0	49	22	28	—	*7	14	29	12	*8	19	10
455.2	23	12	11	—	*6	*5	12	*6	*8	*7	*
455.3	14	*6	*9	—	*	*	*6	*	*	*6	*
455.5	9	*6	*	*	*	*	*	*	*	*	*
455.6	25	11	14	*	*9	*5	11	*5	*5	14	*
455.8	12	*5	*6	—	*	*	*6	*	*	*5	*
456	67	38	29	*	19	29	17	12	11	29	16
456.0	*5	*	*	*	*	*	*	*	*	*	*
456.1	11	*6	*5	*	*	*	*	*	*	*5	*
456.2	41	25	16	*	*8	21	11	*8	*5	19	9
456.20	24	13	10	*	*6	11	*6	*5	*	9	*5
456.21	17	11	*6	*	*	10	*	*	*	9	*
456.8	*6	*	*	*	*	*	*	*	*	*	*
457	24	9	15	*	*8	*7	*9	*7	*6	10	*
457.1	13	*	10	*	*	*	*5	*	*	*5	*
458	332	151	181	*5	34	72	220	64	87	113	68
458.0	97	43	54	*	11	16	69	19	27	30	21
458.2	30	14	16	*	*	*6	21	*	*7	11	*7
458.9	202	92	109	*	19	49	128	39	52	71	40

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
459	108	48	60	*	11	24	70	26	26	39	17
459.0	11	*	*6	*	*	*	*6	*	*	*	*
459.2	12	*6	*6	*	*	*	*	*	*	*	*
459.8	67	30	37	*	*	14	47	18	19	21	*8
459.81	60	27	33	*	*	13	43	16	18	19	*7
459.89	*7	*	*	*	*	*	*	*	*	*	*
459.9	14	*5	9	*	*	*	11	*	*	10	*
460–519	9,368	4,535	4,833	1,098	1,314	1,893	5,063	1,999	2,331	3,475	1,562
461	66	29	37	13	22	15	17	14	18	25	9
461.0	24	12	12	*	*9	*	*5	*	*6	10	*
461.2	*7	*	*	*	*	*	*	*	*	*	*
461.8	*5	*	*	*	*	*	*	*	*	*	*
461.9	28	10	17	*	*8	*8	*7	*7	*8	11	*
462	54	26	28	19	22	*5	*8	9	12	27	*6
463	25	11	14	*9	13	*	*	*6	*	12	*
464	40	24	15	25	*	*6	*5	*8	10	12	10
464.0	*7	*	*	*	*	*	*	*	*	*	*
464.1	*5	*	*	*	*	*	*	*	*	*	*
464.3	*5	*	*	*	*	*	*	*	*	*	*
464.4	21	14	*7	21	*	*	—	*	*6	*7	*6
465	162	69	93	62	41	20	38	34	43	58	27
465.9	160	68	92	61	40	20	38	32	43	58	27
466	384	196	188	184	42	53	104	77	95	143	69
466.0	222	96	126	26	42	51	103	46	54	90	32
466.1	162	100	62	158	*	*	*	31	41	53	37
470	15	9	*6	*	9	*	*	*5	*	*	*
471	*9	*	*	*	*5	*	*	*	*	*	*
472	9	*	*	*	*	*	*	*	*	*	*
472.0	*9	*	*	*	*	*	*	*	*	*	*
473	143	58	86	18	52	34	38	28	33	58	24
473.0	25	10	15	*	10	*	*6	*	*7	9	*
473.2	12	*7	*5	*	*	*	*	*	*	*	*
473.8	11	*6	*5	*	*	*	*	*	*	*	*
473.9	90	33	57	10	33	21	26	18	19	37	16
474	43	22	21	31	11	*	—	23	*	10	*7
474.0	18	*8	11	11	*7	*	—	*8	*	*	*
474.1	24	14	10	20	*	*	—	15	*	*	*
474.10	13	*7	*6	12	*	—	—	9	*	*	*
474.11	*6	*	*	*	*	*	—	*	*	*	*
475	12	*	*8	*	*8	*	*	*	*	*	*
477	24	13	11	*	9	*8	*6	*	*7	12	*
477.9	22	12	10	*	*8	*6	*5	*	*6	11	*
478	87	47	39	20	25	20	22	23	18	31	14
478.0	*7	*	*	—	*6	*	*	*	*	*	*
478.1	18	*8	9	*	*6	*	*	*	*5	*5	*
478.2	18	11	*7	*	*7	*	*	*5	*	*6	*
478.29	14	10	*	*	*6	*	*	*	*	*	*
478.3	14	*6	*8	*	*	*	*7	*	*	*6	*
478.30	*6	*	*	*	*	*	*	*	*	*	*
478.6	*6	*	*	*	*	*	*	*	*	*	*
478.7	17	11	*6	*9	*	*	*	*5	*	*6	*
478.74	*9	*7	*	*6	*	*	*	*	*	*	*
478.75	*5	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
480	51	27	24	39	*	*	*	*6	13	17	15
480.1	26	14	12	26	*	—	*	*	*7	9	*7
480.9	22	11	11	11	*	*	*	*	*6	*7	*7
481	86	50	36	*6	22	22	37	17	21	28	19
482	441	248	193	18	46	69	308	68	115	188	70
482.0	34	18	17	*	*	*	26	*	*7	18	*
482.1	71	43	28	*6	10	13	42	14	16	27	13
482.2	46	23	23	*	*5	11	28	*7	18	11	10
482.3	27	17	11	*	*	*6	17	*	*7	11	*5
482.30	12	*9	*	*	*	*	*7	*	*	*6	*
482.39	10	*6	*	*	*	*	*6	*	*	*	*
482.4	73	40	33	*	*6	12	53	10	15	35	14
482.8	163	94	69	*	11	18	129	17	47	79	19
482.82	13	*8	*	*	*	*	10	*	*	*7	*
482.83	100	59	41	*	*7	9	80	*8	38	44	*9
482.89	49	25	24	*	*	*7	37	*6	*7	27	*9
482.9	27	14	13	*	*7	*	13	11	*	*7	*
483	12	*6	*6	*	*	*	*	*	*	*	*
483.0	9	*	*	*	*	*	*	*	*	*	*
484	*8	*6	*	*	*	*	*	*	*	*	*
485	56	22	35	22	*6	*	24	*7	22	22	*6
486	1,259	599	660	171	167	199	721	279	310	455	215
487	38	14	24	*7	12	*5	14	*5	10	16	*7
487.0	*7	*	*	*	*	*	*	*	*	*	*
487.1	26	11	15	*	9	*	*9	*	*8	13	*
490	82	32	50	14	22	17	28	12	16	40	14
491	824	375	449	*	45	212	565	163	211	339	112
491.0	*7	*	*	—	*	*	*	*	*	*	*
491.2	773	353	420	*	34	194	543	152	195	320	106
491.20	124	60	64	*	10	36	79	25	27	50	21
491.21	648	293	356	*	25	158	464	126	168	270	84
491.8	*5	*	*	*	*	*	*	*	*	*	*
491.9	37	15	22	*	*5	15	16	*8	13	12	*
492	241	136	105	*	*9	61	170	55	66	82	39
492.0	20	15	*6	—	*	*9	*8	*	*	*7	*
492.8	220	121	99	*	*6	53	162	50	62	75	34
493	1,183	432	751	298	359	246	281	317	307	354	206
493.0	60	31	28	46	*9	*	*	17	9	*5	28
493.00	32	15	17	22	*5	*	*	*7	*7	*	14
493.01	28	16	12	24	*	*	—	10	*	*	14
493.2	236	75	160	*	23	70	142	56	66	71	43
493.20	199	65	134	*	16	59	123	44	60	60	36
493.21	36	10	26	*	*6	11	19	12	*6	11	*7
493.9	885	325	560	251	326	171	136	243	231	276	134
493.90	673	223	450	133	268	142	130	187	167	213	106
493.91	212	102	110	119	58	29	*6	56	64	63	29
494	33	11	22	*	*	*9	20	*9	*7	11	*6
496	1,683	906	777	*	41	381	1,259	349	429	633	272
500	11	11	*	—	*	*	*9	*	*	*	*
501	13	12	*	—	*	*	10	*6	*	*	*
506	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
507	271	155	116	11	22	31	207	53	59	110	50
507.0	267	153	114	*9	22	31	205	52	57	109	50
508	*7	*	*	*	*	*	*	*	*	*	*
510	24	15	10	*	*7	*6	10	*6	*	*8	*6
510.9	20	13	*7	*	*6	*5	*8	*6	*	*6	*5
511	410	193	218	10	49	88	263	92	95	147	76
511.0	27	12	15	*	*5	11	11	*	*6	12	*5
511.8	35	17	18	*	*	*6	24	*	10	10	11
511.9	345	161	184	*8	40	71	226	82	79	125	59
512	112	62	50	*	26	28	55	22	23	47	20
512.1	53	28	25	*	*5	13	33	10	11	21	11
512.8	54	31	23	*	19	13	21	11	11	25	*8
513	13	10	*	*	*	*	*6	*	*	*5	*
513.0	11	9	*	*	*	*	*	*	*	*	*
514	28	9	19	*	*7	*7	12	*	*	16	*6
515	190	85	105	*	16	33	141	30	70	70	21
516	29	13	16	*	*	*6	19	*	*9	12	*
516.3	18	*9	9	*	*	*	12	*	*	*8	*
516.8	10	*	*6	*	*	*	*6	*	*	*	*
517	15	*	11	*	*5	*	*	*	*	*	*
517.8	14	*	10	*	*	*	*	*	*	*	*
518	1,079	533	546	66	147	260	606	224	240	424	192
518.0	367	177	190	29	58	82	198	68	86	155	59
518.1	10	*7	*	*	*	*	*	*	*	*	*
518.3	19	*8	11	*	*	*6	11	*	*	*8	*
518.4	26	11	16	*	*	*7	14	*5	*6	*8	*7
518.5	64	34	30	*	14	13	32	13	13	25	12
518.8	592	296	296	32	63	150	347	133	130	222	107
518.81	442	217	225	13	44	113	271	98	95	168	81
518.82	91	47	44	14	11	21	45	20	21	35	15
518.89	58	32	26	*	*8	15	31	14	14	19	11
519	76	35	41	27	11	14	23	16	24	24	12
519.0	*9	*5	*	*	*	*	*	*	*	*	*
519.1	36	15	20	12	*7	*6	11	*7	13	12	*
519.8	16	*7	*8	*	*	*	*6	*	*	*6	*
519.9	9	*	*	*7	*	*	*	*	*	*	*
520–579	8,156	3,538	4,618	435	1,733	2,096	3,892	1,758	1,950	2,968	1,480
521	16	11	*5	*	*9	*	*	*5	*	*	*
521.0	16	11	*5	*	*9	*	*	*5	*	*	*
522	18	9	*8	*	*7	*	*	*	*	*5	*
522.5	14	*7	*6	*	*6	*	*	*	*	*	*
523	12	*8	*	*	*	*	*	*	*	*	*
524	33	13	20	*	26	*	*	9	9	11	*
524.0	10	*	*6	*	*8	*	*	*	*	*	*
524.1	10	*	*5	*	*8	*	*	*	*	*	*
524.10	*8	*	*5	*	*7	*	*	*	*	*	*
524.6	*7	*	*6	*	*5	*	*	*	*	*	*
525	*8	*	*	*	*	*	*	*	*	*	*
526	*7	*5	*	*	*	*	*	*	*	*	*
527	25	*9	16	*	*	*7	13	*6	*	10	*7
527.2	15	*6	9	*	*	*	*8	*	*	*7	*

* Figure does not meet standard of reliability or precision.
– Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
528	50	23	28	*8	17	12	13	12	14	16	*8
528.0	24	*9	15	*	*7	*7	*7	*7	*7	*8	*
528.3	*8	*	*	*	*	*	*	*	*	*	*
528.9	11	*5	*5	*	*	*	*	*	*	*	*
529	*5	*	*	*	*	*	*	*	*	*	*
530	929	403	526	80	138	242	469	205	216	355	153
530.1	257	113	144	*8	44	70	134	51	58	98	50
530.10	102	47	55	*	21	32	45	24	19	41	18
530.11	115	50	65	*	16	27	69	21	30	43	21
530.19	40	16	24	*	*8	11	20	*6	*9	14	11
530.2	53	33	20	*	11	14	28	14	13	14	13
530.3	52	23	29	*	*	*8	42	11	11	20	11
530.5	26	11	15	*	*	11	13	*5	*	12	*5
530.6	10	*5	*5	—	*	*	*9	*	*	*	*
530.7	25	16	9	*	*5	*9	11	*5	*5	10	*
530.8	497	198	300	69	72	129	227	116	119	196	66
530.81	475	187	288	69	68	124	215	110	113	189	63
530.82	13	*7	*6	*	*	*	*8	*	*	*	*
530.89	*8	*	*6	*	*	*	*	*	*	*	*
531	170	80	90	*	19	47	102	31	49	54	35
531.0	35	14	21	*	*	11	20	*	10	11	9
531.00	33	13	19	*	*	11	18	*	*8	11	9
531.3	*7	*	*	*	*	*	*	*	*	*	*
531.30	*6	*	*	*	*	*	*	*	*	*	*
531.4	63	35	28	*	*	16	41	11	17	21	14
531.40	62	35	27	*	*	16	40	10	17	21	14
531.5	*8	*	*	—	*	*	*5	*	*	*	*
531.50	*8	*	*	—	*	*	*	*	*	*	*
531.7	*7	*	*	*	*	*	*	*	*	*	*
531.70	*6	*	*	*	*	*	*	*	*	*	*
531.9	47	20	28	*	*8	12	27	11	14	15	*7
531.90	46	20	27	*	*8	12	26	11	14	15	*7
532	131	76	55	*	19	37	75	27	31	50	23
532.0	27	16	10	*	*	*8	16	*	*7	10	*6
532.00	26	16	10	*	*	*8	15	*	*6	10	*6
532.1	*6	*	*	—	*	*	*	*	*	*	*
532.10	*5	*	*	—	*	*	*	*	*	*	*
532.3	*6	*	*	—	*	*	*	*	*	*	*
532.30	*6	*	*	—	*	*	*	*	*	*	*
532.4	46	29	17	*	*7	9	30	10	10	18	*9
532.40	45	28	17	*	*7	*9	29	9	10	18	*9
532.9	34	17	16	*	*	12	17	*8	*8	12	*6
532.90	32	17	15	*	*	12	16	*8	*8	11	*6
533	201	91	110	*	28	58	114	46	54	69	32
533.4	17	*6	11	*	*	*	11	*	*5	*8	*
533.40	17	*6	11	*	*	*	10	*	*5	*8	*
533.7	26	13	13	—	*	*7	15	*9	*6	*7	*
533.70	26	13	13	—	*	*7	15	*9	*6	*7	*
533.9	148	67	81	*	21	45	82	34	40	49	25
533.90	147	66	81	*	20	45	82	34	40	49	25

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
535	620	280	340	10	122	162	326	129	142	250	100
535.0	93	39	54	*	20	22	49	19	23	38	13
535.00	69	27	42	*	17	18	33	15	17	29	*8
535.01	24	12	12	*	*	*	16	*	*5	9	*6
535.1	69	28	41	*	9	15	45	16	16	24	13
535.10	59	23	35	*	*9	13	37	13	15	20	11
535.11	10	*	*6	*	*	*	*8	*	*	*	*
535.3	26	20	*6	*	11	11	*	*6	*6	9	*5
535.30	17	13	*	—	*8	*6	*	*5	*	*	*
535.31	10	*7	*	*	*	*5	*	*	*	*	*
535.4	143	62	80	*	27	30	84	30	36	50	27
535.40	101	40	61	*	21	24	54	20	28	34	18
535.41	42	22	19	*	*5	*6	31	9	*8	16	*8
535.5	220	94	127	*	43	62	111	43	44	101	32
535.50	187	77	110	*	38	54	91	36	38	88	25
535.51	34	17	17	*	*	*8	20	*7	*6	14	*7
535.6	67	36	31	*	13	22	32	16	16	26	10
535.60	53	26	27	*	11	18	23	10	14	21	*8
535.61	14	10	*	*	*	*	*9	*6	*	*5	*
536	180	70	109	10	40	52	78	32	44	70	33
536.2	24	11	13	*5	*6	*	9	*7	*	11	*
536.3	62	23	39	*	16	22	23	12	10	27	13
536.8	78	30	48	*	15	21	38	11	27	24	15
536.9	13	*5	*8	*	*	*	*6	*	*	*7	*
537	50	22	28	*	*5	14	30	10	10	19	11
537.0	22	11	12	*	*	*7	13	*	*	*7	*6
537.8	20	*8	12	*	*	*	13	*	*	*7	*
537.83	*6	*	*	—	*	*	*	*	*	*	*
537.89	*9	*	*	*	*	*	*5	*	*	*	*
540	231	141	91	59	122	33	18	42	51	71	68
540.0	48	30	18	14	18	9	*6	9	10	12	15
540.1	31	20	11	*5	12	*8	*6	*	*6	11	*9
540.9	153	91	62	39	93	16	*	29	34	47	43
541	13	*	*8	*	10	*	*	*	*	*	*
542	*8	*	*5	*	*	*	*	*	*	*	*
543	15	*	12	*	*8	*	*	*	*	*5	*
543.9	13	*	10	*	*6	*	*	*	*	*	*
550	106	89	18	16	15	18	57	34	22	34	16
550.1	29	25	*	*	*	*	19	*8	*8	*9	*
550.10	20	17	*	*	*	*	12	*7	*	*6	*
550.11	*6	*6	*	*	*	*	*6	*	*	*	*
550.9	76	63	13	11	13	16	37	26	14	25	12
550.90	53	41	11	*6	10	12	25	19	*8	17	*9
550.91	*9	*8	*	*	*	*	*5	*	*	*	*
550.92	14	13	*	*	*	*	*6	*5	*	*	*
552	59	18	40	*	9	17	32	15	16	18	9
552.1	18	*6	12	*	*	*8	*6	*5	*5	*	*
552.2	25	*8	18	*	*	*7	13	*6	*7	*9	*
552.20	*8	*	*5	—	*	*	*	*	*	*	*
552.21	14	*	11	*	*	*	*8	*	*	*5	*
552.3	*5	*	*	—	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
553	543	195	348	*8	59	149	327	111	143	213	76
553.1	48	21	27	*6	*8	16	18	11	10	19	*7
553.2	92	35	57	*	13	28	52	19	26	36	11
553.20	31	10	21	*	*	*9	19	*	9	13	*
553.21	53	23	31	*	*8	18	27	12	14	21	*6
553.29	*8	*	*5	*	*	*	*	*	*	*	*
553.3	395	137	258	*	37	102	253	78	106	155	55
555	80	33	47	*	32	23	22	22	21	24	13
555.0	16	*6	10	*	*8	*5	*	*	*	*	*
555.1	16	*8	*9	*	*5	*	*7	*	*6	*5	*
555.2	*9	*	*6	*	*	*	*	*	*	*	*
555.9	38	16	22	*	16	12	10	12	10	10	*6
556	53	26	27	*	27	12	13	14	14	15	10
556.8	*6	*	*	*	*	*	*	*	*	*	*
556.9	36	17	19	*	18	*7	9	*8	12	11	*5
557	65	24	40	*	*	15	45	15	18	22	10
557.0	37	14	23	*	*	*9	24	*9	9	13	*6
557.1	12	*6	*6	*	*	*	9	*	*	*	*
557.9	15	*	11	*	*	*	11	*	*	*6	*
558	400	154	247	124	95	59	122	81	97	164	58
558.1	10	*	*8	—	*	*	*7	*	*	*	*
558.9	386	149	237	124	94	54	114	77	93	161	53
560	630	274	356	18	108	147	357	129	145	231	125
560.1	273	127	146	*8	60	64	142	54	58	110	52
560.2	21	11	10	*	*	*	14	*	*	9	*
560.3	71	26	45	*	*8	*8	53	16	14	30	10
560.39	67	24	43	*	*7	*8	50	16	13	27	10
560.8	131	53	78	*	18	33	78	27	37	36	30
560.81	86	34	52	*	14	23	48	17	24	24	21
560.89	45	19	25	*	*	10	30	10	13	12	9
560.9	128	53	76	*	20	38	69	26	30	44	28
562	488	184	304	*	31	98	358	117	116	179	75
562.0	10	*	*8	—	*	*	*8	*	*	*	*
562.00	9	*	*7	—	*	*	*6	*	*	*	*
562.1	477	181	296	*	30	96	351	113	113	176	75
562.10	213	74	139	*	*6	38	168	55	51	75	31
562.11	174	72	102	*	22	49	102	42	41	63	28
562.12	68	28	40	—	*	*7	60	12	15	29	12
562.13	23	*7	16	—	*	*	20	*	*6	*9	*
564	293	102	191	14	65	63	151	59	75	99	60
564.0	179	74	105	10	33	36	99	35	46	64	34
564.1	80	12	68	*	25	19	36	17	22	24	18
564.7	*6	*	*	—	*	*	*6	*	*	*	*
564.8	12	*	*8	*	*	*	*	*	*	*	*
565	18	11	*7	*	*7	*	*	*	*	*7	*
565.0	*8	*	*	*	*	*	*	*	*	*	*
565.1	9	*7	*	*	*	*	*	*	*	*	*
566	27	19	*7	*	12	*7	*7	*	*6	10	*6
567	91	47	44	*	24	30	33	20	25	24	21
567.2	58	31	27	*	17	22	17	13	13	16	15
567.8	10	*	*5	*	*	*	*	*	*	*	*
567.9	22	12	11	*	*	*6	11	*5	*8	*7	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
568	164	38	126	*	58	46	58	37	41	60	26
568.0	143	34	109	*	48	42	52	32	37	52	22
568.8	20	*	16	*	10	*	*6	*	*	*8	*
568.81	11	*	10	*	*7	*	*	*	*	*	*
568.89	*7	*	*	*	*	*	*	*	*	*	*
569	215	93	122	*7	38	47	122	51	57	73	34
569.0	12	*5	*7	*	*	*	*7	*	*	*5	*
569.1	*9	*	*8	*	*	*	*6	*	*	*	*
569.3	39	16	23	*	*7	*8	22	9	11	14	*
569.4	23	11	12	*	*	*5	12	*7	*6	*7	*
569.41	*7	*	*	*	*	*	*	*	*	*	*
569.49	14	*7	*7	*	*	*	*7	*	*	*5	*
569.5	15	*8	*7	*	*	*	*7	*	*5	*	*
569.6	23	11	12	*	*5	*6	11	*5	*6	*7	*
569.61	*7	*	*	*	*	*	*	*	*	*	*
569.69	15	*7	*8	*	*	*	*9	*	*	*5	*
569.8	84	36	49	*	12	18	51	19	21	28	16
569.81	14	*	10	*	*	*	*	*	*	*6	*
569.82	*6	*	*	*	*	*	*	*	*	*	*
569.83	21	10	11	*	*	*5	12	*5	*	*7	*
569.84	*9	*	*	—	—	*	*8	*	*	*	*
569.85	15	*7	*8	—	*	*	12	*	*	*7	*
569.89	20	9	11	*	*	*	11	*6	*7	*	*
569.9	*6	*	*	*	*	*	*	*	*	*	*
570	20	10	9	*	*7	*6	*	*6	*	*7	*
571	356	217	140	*	91	152	110	71	69	137	79
571.1	42	33	9	*	19	19	*	10	9	14	9
571.2	124	88	36	—	27	67	30	26	24	42	33
571.3	38	27	11	—	17	15	*6	*7	10	11	11
571.4	15	*7	*7	*	*	*5	*5	*	*	*7	*
571.40	*7	*	*	*	*	*	*	*	*	*	*
571.49	*7	*	*	*	*	*	*	*	*	*	*
571.5	101	45	56	*	14	34	51	15	18	51	17
571.6	*8	*	*7	*	*	*	*6	*	*	*	*
571.8	20	10	10	*	*7	*7	*5	*6	*	*7	*
572	125	70	55	*	26	57	39	26	18	51	30
572.0	*7	*	*	*	*	*	*	*	*	*	*
572.2	50	29	21	*	11	20	18	11	*7	20	12
572.3	42	24	18	*	*8	22	11	*7	*7	20	*7
572.4	*7	*	*	—	*	*	*	*	*	*	*
572.8	18	*9	9	*	*	*9	*	*	*	*8	*5
573	64	29	35	*	14	19	30	12	18	24	11
573.0	*8	*	*6	—	*	*	*6	*	*	*	*
573.3	21	10	10	*	*7	*7	*6	*6	*	*8	*
573.8	22	*8	14	*	*	*7	12	*	*5	*8	*
573.9	12	*8	*	*	*	*	*6	*	*	*5	*
574	695	221	474	*	182	205	304	161	156	245	134
574.0	151	54	97	*	39	48	62	36	29	51	34
574.00	133	46	87	*	36	42	54	32	26	45	30
574.01	17	*8	9	—	*	*6	*8	*	*	*7	*
574.1	292	85	207	*	84	100	107	76	69	94	54
574.10	278	79	199	*	80	96	101	72	64	90	52
574.11	14	*6	*8	*	*	*	*6	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
574.2	154	49	105	*	29	35	88	30	37	59	27
574.20	150	46	103	*	28	35	85	29	36	59	26
574.3	23	*7	16	*	*7	*	11	*	*	11	*
574.30	13	*	*9	—	*5	*	*5	*	*	*6	*
574.31	10	*	*7	*	*	*	*5	*	*	*	*
574.4	35	12	23	*	14	*8	12	9	9	12	*
574.40	19	*	15	*	10	*	*6	*6	*	*7	*
574.41	16	*8	*8	*	*	*5	*7	*	*6	*	*
574.5	41	14	27	*	9	*8	24	*6	*9	17	*9
574.50	20	*7	13	—	*	*5	10	*	*	*8	*
574.51	21	*7	14	*	*	*	14	*	*	9	*
575	126	48	79	*	43	31	51	22	35	47	22
575.0	40	18	21	—	11	11	17	*5	12	13	10
575.1	49	18	32	*	18	12	19	*8	12	22	*8
575.3	*6	*	*	*	*	*	*	*	*	*	*
575.6	*7	*	*	*	*	*	*	*	*	*	*
575.8	18	*	14	*	*9	*	*5	*	*	*7	*
576	99	45	54	*	19	24	52	19	21	35	24
576.1	40	19	21	*	*7	*9	22	*7	9	12	11
576.2	32	14	18	*	*	9	17	*	*7	12	*8
576.8	23	11	12	*	*5	*	12	*7	*	*8	*
577	338	159	179	*5	134	103	95	64	95	121	58
577.0	229	103	126	*	87	70	69	42	62	86	40
577.1	72	37	35	*	32	24	15	13	24	25	11
577.2	16	*8	*7	*	*7	*	*	*	*	*5	*
577.8	18	10	*8	*	*7	*	*	*	*	*	*
578	340	154	185	*8	36	67	229	80	81	109	70
578.0	42	20	22	*	*8	*8	24	13	11	*7	11
578.1	87	41	46	*5	10	15	56	22	20	26	18
578.9	211	94	117	*	18	44	149	44	49	76	42
579	31	13	18	*8	*	*7	12	*7	10	9	*
579.3	12	*7	*5	*	*	*	*	*	*5	*	*
579.8	10	*	*6	*	*	*	*	*	*	*	*
580-629	6,812	2,285	4,527	155	1,997	1,589	3,071	1,383	1,602	2,632	1,194
580	12	*	*7	*	*	*	*	*	*	*5	*
580.9	*7	*	*	*	*	*	*	*	*	*	*
581	42	16	26	*	12	12	14	*9	*8	17	*8
581.8	18	*	13	*	*	*8	*6	*	*	*8	*
581.81	17	*	13	*	*	*8	*5	*	*	*7	*
581.9	20	*9	12	*	*6	*	*7	*	*	*9	*
582	49	23	26	*	*9	12	28	15	*8	15	12
582.8	13	*	*9	—	*	*	*	*	*	*	*7
582.81	9	*	*8	—	*	*	*	*	*	*	*6
582.9	31	17	15	*	*	*6	22	11	*6	11	*
583	182	84	97	*	30	54	95	46	38	61	37
583.8	165	75	90	*	23	51	91	41	34	58	33
583.81	157	69	88	*	19	49	89	38	33	56	30
583.89	*8	*6	*	*	*	*	*	*	*	*	*
583.9	14	*8	*6	*	*6	*	*	*	*	*	*
584	298	156	141	*	37	60	196	76	67	106	49
584.5	30	19	11	*	*	*6	19	10	*6	11	*
584.8	*7	*	*	*	*	*	*	*	*	*	*
584.9	261	134	126	*	31	52	174	64	60	93	43

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
585	305	169	136	*	42	79	180	77	66	98	64
586	70	34	35	*	*5	10	52	12	14	31	13
588	19	*8	11	*	*	*6	*6	*	*	9	*
588.8	15	*6	*8	*	*	*	*6	*	*	*7	*
590	204	43	161	20	97	38	49	35	46	79	44
590.0	10	*	*7	*	*	*	*	*	*	*	*
590.00	10	*	*7	*	*	*	*	*	*	*	*
590.1	116	25	91	12	57	17	30	21	29	40	26
590.10	116	25	91	12	57	17	30	21	29	40	26
590.8	72	13	59	*7	35	17	14	12	14	35	12
590.80	72	12	59	*7	35	17	13	12	14	35	11
591	126	65	61	*5	42	32	47	32	27	46	21
592	251	144	106	*	108	85	55	54	60	98	39
592.0	97	48	49	*	37	32	27	24	22	36	15
592.1	149	94	55	*	69	52	27	30	37	59	23
593	387	200	187	18	42	69	257	76	110	126	74
593.2	21	11	9	*	*	*	12	*	*5	*7	*
593.3	10	*6	*	*	*	*	*	*	*	*	*
593.4	32	14	18	*	*8	*8	14	*8	*8	12	*
593.7	14	*5	*9	11	*	*	*	*	*	*	*
593.70	11	*	*7	9	*	*	*	*	*	*	*
593.8	35	17	18	*	*6	10	17	*8	12	*7	9
593.89	30	16	14	*	*6	*8	15	*6	10	*7	*8
593.9	270	144	126	*	17	40	210	47	78	94	51
594	16	14	*	*	*	*	10	*	*	*	*
594.1	13	11	*	—	*	*	*9	*	*	*	*
595	70	25	44	*	10	13	45	15	20	25	10
595.0	21	*7	15	*	*	*	14	*	*8	*7	*
595.1	*7	*	*6	*	*	*	*	*	*	*	*
595.2	*8	*	*	*	*	*	*7	*	*	*	*
595.8	13	*8	*	*	*	*	9	*	*	*	*
595.82	*7	*	*	*	*	*	*6	*	*	*	*
595.9	19	*5	14	*	*	*	13	*	*	*8	*
596	179	128	51	*	22	33	120	42	47	54	36
596.0	59	58	*	*	*	10	48	14	16	18	11
596.1	*8	*	*	*	*	*	*5	*	*	*	*
596.3	*6	*	*	*	*	*	*	*	*	*	*
596.4	*8	*	*	*	*	*	*	*	*	*	*
596.5	73	40	33	*	16	13	42	17	21	23	12
596.54	66	37	29	*	16	12	36	15	19	22	10
596.8	20	14	*6	*	*	*	12	*	*	*6	*5
598	42	34	*8	*	*6	11	24	11	9	16	*7
598.9	34	28	*6	*	*	*9	20	*9	*7	13	*
599	1,851	619	1,232	54	251	288	1,257	382	462	746	261
599.0	1,551	425	1,125	46	196	226	1,083	320	391	626	214
599.6	27	23	*	*	*	*	22	*5	*6	9	*6
599.7	260	168	92	*7	53	56	145	55	61	105	39
599.8	*6	*	*	*	*	*	*	*	*	*	*
600	351	351	—	—	*	63	285	83	100	110	58
601	55	55	—	—	*	14	37	10	17	19	10
601.0	19	19	—	—	*	*	13	*	*5	*8	*
601.1	23	23	—	—	*	*	18	*	*9	*	*5
601.9	11	11	—	—	*	*	*5	*	*	*5	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
602	*8	*8	—	—	*	*	*6	*	*	*	*
603	10	10	—	*	*	*	*	*	*	*	*
603.9	*8	*8	—	*	*	*	*	*	*	*	*
604	21	21	—	*	*6	*6	*8	*5	*	10	*
604.9	20	20	—	*	*6	*6	*8	*	*	9	*
604.90	12	12	—	*	*	*	*	*	*	*5	*
604.99	*8	*8	—	*	*	*	*	*	*	*	*
605	12	12	—	*5	*	*	*	*	*	*	*
607	29	29	—	*	*6	13	*8	*8	*	10	*6
607.8	18	18	—	*	*	10	*5	*	*	*6	*
607.84	12	12	—	—	*	*8	*	*	*	*	*
608	22	22	—	*	*6	*7	*7	*6	*	*7	*5
608.4	*6	*6	—	*	*	*	*	*	*	*	*
608.8	10	10	—	*	*	*	*	*	*	*	*
610	37	*	36	—	11	14	11	13	*	13	*6
610.1	29	*	29	—	*8	11	10	9	*	11	*5
611	53	*	49	*	26	15	11	15	11	16	11
611.0	*7	*	*7	*	*	*	*	*	*	*	*
611.1	21	*	18	*	15	*5	*	*8	*	*5	*
611.7	12	*	11	—	*	*	*5	*	*	*	*
611.72	10	*	9	—	*	*	*	*	*	*	*
611.8	10	—	10	—	*	*	*	*	*	*	*
614	299	—	299	*	219	61	17	45	56	142	55
614.0	12	—	12	*	10	*	*	*	*	*7	*
614.1	32	—	32	*	23	*8	*	*	*5	15	*7
614.2	13	—	13	—	10	*	*	*	*	*6	*
614.3	11	—	11	*	10	*	*	*	*	*	*
614.4	*8	—	*8	*	*6	*	*	*	*	*	*
614.6	192	—	192	*	134	46	12	27	36	93	37
614.9	27	—	27	*	24	*	*	*5	*6	12	*
615	24	—	24	*	21	*	*	*	*6	9	*6
615.0	*7	—	*7	—	*6	*	—	*	*	*	*
615.9	12	—	12	*	12	*	*	*	*	*5	*
616	160	—	160	*	93	52	14	32	35	66	27
616.0	124	—	124	—	68	47	*8	26	28	49	22
616.1	27	—	27	*	20	*	*	*5	*5	13	*
616.10	27	—	27	*	20	*	*	*5	*5	13	*
617	292	—	292	*	189	92	11	50	64	120	58
617.0	163	—	163	—	89	66	*9	25	36	68	34
617.1	41	—	41	*	32	*8	*	*8	10	13	11
617.2	10	—	10	—	*8	*	*	*	*	*	*
617.3	48	—	48	*	37	10	*	11	12	18	*7
617.9	21	—	21	—	17	*	*	*	*	13	*
618	267	—	267	*	70	99	97	37	59	115	56
618.0	83	—	83	*	16	29	38	11	18	34	20
618.1	28	—	28	—	14	*7	*7	*5	*7	12	*
618.2	28	—	28	—	11	13	*	*	*7	11	*7
618.3	24	—	24	—	*	*7	13	*6	*6	*8	*
618.4	22	—	22	—	10	*8	*	*	*5	11	*
618.5	30	—	30	—	*	14	14	*	*	14	*8
618.6	24	—	24	—	*	10	11	*	*6	*8	*6
618.8	25	—	25	*	*9	11	*	*	*	15	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
619	9	—	9	*	*	*	*	*	*	*	*
619.1	*7	—	*7	*	*	*	*	*	*	*	*
620	293	—	293	*6	191	81	16	56	62	118	57
620.0	47	—	47	*	31	15	*	*8	13	19	*7
620.1	75	—	75	*	54	18	*	12	15	34	15
620.2	116	—	116	*	78	26	*9	25	21	47	23
620.5	*7	—	*7	*	*	*	*	*	*	*	*
620.8	43	—	43	*	22	17	*	10	10	14	9
621	89	—	89	*	34	35	20	21	21	31	15
621.0	35	—	35	—	11	15	*9	11	*6	12	*6
621.2	*8	—	*8	—	*	*	*	*	*	*	*
621.3	21	—	21	—	*	11	*6	*5	*5	*6	*
621.5	9	—	9	—	*7	*	*	*	*	*	*
621.8	10	—	10	*	*	*	*	*	*	*	*
622	39	—	39	—	23	12	*	*7	10	16	*5
622.1	26	—	26	—	18	*7	*	*	*7	12	*
622.7	*8	—	*8	—	*	*	*	*	*	*	*
623	19	—	19	*	10	*	*5	*	*	*7	*
623.8	13	—	13	*	*7	*	*	*	*	*	*
625	318	—	318	*	186	96	35	30	69	147	72
625.0	19	—	19	—	16	*	—	*	*	10	*
625.3	65	—	65	*	52	13	—	*6	13	33	14
625.5	*5	—	*5	—	*	*	—	*	*	*	*
625.6	143	—	143	*	44	66	33	13	32	64	35
625.8	*8	—	*8	*	*6	*	*	*	*	*	*
625.9	74	—	74	*	60	12	*	*8	14	37	15
626	237	—	237	*	153	79	*	38	50	105	44
626.2	162	—	162	*	107	55	*	26	33	76	28
626.6	*5	—	*5	*	*	*	*	*	*	*	*
626.8	54	—	54	*	33	19	*	*9	12	20	13
626.9	*7	—	*7	*	*	*	*	*	*	*	*
627	44	—	44	—	*5	22	17	*7	14	16	*6
627.0	*6	—	*6	—	*	*	—	*	*	*	*
627.1	25	—	25	—	*	12	13	*	*8	*8	*
627.2	*6	—	*6	—	*	*	*	*	*	*	*
628	*7	—	*7	*	*6	*	—	*	*	*	*
630–677	8,265	—	8,265	24	8,234	*7	—	1,457	1,722	2,877	2,209
632	17	—	17	*	17	—	—	*	*	*7	*
633	39	—	39	—	39	—	—	*7	*8	17	*6
633.1	36	—	36	—	36	—	—	*7	*7	16	*6
634	31	—	31	*	31	—	—	12	*	11	*
634.9	24	—	24	*	24	—	—	11	*	*7	*
634.91	19	—	19	*	19	—	—	*9	*	*5	*
635	11	—	11	*	11	*	—	*	*	*	*
635.9	*8	—	*8	—	*8	*	—	*	*	*	*
637	*8	—	*8	—	*8	*	—	*	*	*	*
639	16	—	16	—	16	—	—	*	*	*7	*
639.1	*7	—	*7	—	*7	—	—	*	*	*	*
640	9	—	9	—	9	—	—	*	*	*	*
640.0	*8	—	*8	—	*8	—	—	*	*	*	*
640.03	*8	—	*8	—	*8	—	—	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
641	94	—	94	*	94	*	—	18	23	25	29
641.0	*7	—	*7	—	*7	—	—	*	*	*	*
641.01	*6	—	*6	—	*6	—	—	*	*	*	*
641.1	22	—	22	—	22	—	—	*6	*	*	*8
641.11	15	—	15	—	15	—	—	*	*	*	*6
641.13	*7	—	*7	—	*7	—	—	*	*	*	*
641.2	43	—	43	*	43	*	—	*6	14	12	12
641.21	35	—	35	*	35	*	—	*5	9	10	11
641.23	*8	—	*8	—	*8	—	—	*	*	*	*
641.9	13	—	13	—	13	—	—	*	*	*	*
641.91	*6	—	*6	—	*6	—	—	*	*	*	*
641.93	*7	—	*7	—	*7	—	—	*	*	*	*
642	300	—	300	*	298	*	—	48	65	129	58
642.0	22	—	22	*	22	*	—	*	*7	*6	*5
642.01	18	—	18	*	18	*	—	*	*5	*	*
642.3	97	—	97	*	97	*	—	19	18	44	17
642.31	85	—	85	*	85	*	—	16	15	39	15
642.33	10	—	10	—	10	—	—	*	*	*	*
642.4	94	—	94	*	93	*	—	14	24	38	17
642.41	82	—	82	*	81	—	—	11	22	33	16
642.43	*9	—	*9	—	*9	*	—	*	*	*	*
642.5	30	—	30	*	30	—	—	*	*7	13	*6
642.51	26	—	26	*	25	—	—	*	*6	10	*5
642.7	9	—	9	*	9	—	—	*	*	*5	*
642.71	*7	—	*7	*	*7	—	—	*	*	*	*
642.9	40	—	40	*	40	*	—	*	*7	19	10
642.91	36	—	36	*	36	*	—	*	*6	17	*9
643	44	—	44	—	44	—	—	*9	9	22	*
643.0	13	—	13	—	13	—	—	*	*	*7	*
643.03	12	—	12	—	12	—	—	*	*	*6	*
643.1	22	—	22	—	22	—	—	*	*	10	*
643.13	21	—	21	—	21	—	—	*	*	10	*
644	394	—	394	*	392	*	—	68	83	153	90
644.0	122	—	122	*	122	*	—	22	27	45	28
644.03	122	—	122	*	121	*	—	22	27	45	28
644.1	28	—	28	—	28	—	—	*6	*6	11	*
644.13	28	—	28	—	28	—	—	*6	*6	11	*
644.2	244	—	244	*	242	*	—	40	50	97	56
644.21	244	—	244	*	242	*	—	40	50	97	56
645	145	—	145	*	144	—	—	19	29	51	45
645.0	145	—	145	*	144	—	—	19	29	51	45
645.01	142	—	142	*	142	—	—	19	27	51	45
646	250	—	250	*	249	*	—	45	48	108	48
646.1	19	—	19	*	19	—	—	*	*	*9	*6
646.11	17	—	17	*	17	—	—	*	*	*8	*6
646.2	*5	—	*5	—	*5	—	—	*	*	*	*
646.6	180	—	180	*	179	*	—	32	33	81	34
646.61	104	—	104	*	104	—	—	18	18	48	19
646.62	11	—	11	*	10	—	—	*	*	*	*
646.63	60	—	60	*	59	*	—	11	11	25	12
646.8	35	—	35	*	35	—	—	*8	*8	13	*6
646.81	16	—	16	—	16	—	—	*	*	*6	*
646.83	16	—	16	*	16	—	—	*	*	*7	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
647	94	—	94	*	93	*	—	13	17	40	23
647.6	40	—	40	*	39	—	—	*7	*7	16	11
647.61	34	—	34	*	33	—	—	*5	*	14	10
647.63	*5	—	*5	—	*5	—	—	*	*	*	*
647.8	44	—	44	*	44	*	—	*	*8	19	11
647.81	31	—	31	*	31	—	—	*	*5	14	*9
647.83	10	—	10	—	10	*	—	*	*	*	*
648	908	—	908	*	903	*	—	176	184	312	236
648.0	35	—	35	—	35	*	—	*7	*	16	*8
648.01	21	—	21	—	21	*	—	*	*	11	*6
648.03	13	—	13	—	13	—	—	*	*	*	*
648.1	34	—	34	—	34	*	—	*8	*5	11	10
648.11	28	—	28	—	28	*	—	*6	*	10	*8
648.13	*6	—	*6	—	*6	—	—	*	*	*	*
648.2	373	—	373	*	370	*	—	59	75	134	105
648.21	210	—	210	*	208	*	—	27	41	84	58
648.22	126	—	126	*	125	*	—	27	25	33	40
648.23	26	—	26	—	26	*	—	*	*7	11	*
648.24	*8	—	*8	—	*8	—	—	*	*	*	*
648.3	18	—	18	—	18	—	—	*5	*	*	*
648.31	13	—	13	—	13	—	—	*	*	*	*
648.33	*6	—	*6	—	*6	—	—	*	*	*	*
648.4	87	—	87	*	86	*	—	17	18	28	23
648.41	60	—	60	*	60	*	—	9	13	20	19
648.43	20	—	20	*	20	*	—	*7	*	*6	*
648.6	38	—	38	*	37	*	—	*8	*7	13	10
648.61	32	—	32	—	32	*	—	*8	*6	10	*9
648.8	137	—	137	—	137	*	—	28	26	49	34
648.81	118	—	118	—	117	*	—	24	22	41	30
648.83	18	—	18	—	18	—	—	*	*	*7	*
648.9	180	—	180	*	178	*	—	41	43	55	40
648.91	106	—	106	*	106	—	—	23	26	29	27
648.92	*8	—	*8	—	*8	—	—	*	*	*	*
648.93	62	—	62	*	62	*	—	16	14	22	10
650	631	—	631	*	630	*	—	93	124	250	164
651	64	—	64	*	64	*	—	15	13	22	15
651.0	59	—	59	*	59	*	—	14	12	20	13
651.01	44	—	44	*	44	*	—	10	*8	15	11
651.03	15	—	15	—	15	*	—	*	*	*5	*
652	265	—	265	*	264	*	—	52	51	91	71
652.1	*7	—	*7	—	*7	—	—	*	*	*	*
652.11	*6	—	*6	—	*6	—	—	*	*	*	*
652.2	149	—	149	*	148	*	—	23	28	57	41
652.21	142	—	142	*	142	*	—	23	28	52	39
652.23	*5	—	*5	—	*5	—	—	*	*	*	*
652.3	28	—	28	*	28	*	—	*7	*	*9	*8
652.31	27	—	27	*	27	*	—	*6	*	*9	*8
652.4	*6	—	*6	—	*6	—	—	*	*	*	*
652.41	*6	—	*6	—	*6	—	—	*	*	*	*
652.5	25	—	25	*	24	—	—	*7	*5	*9	*
652.51	25	—	25	*	24	—	—	*7	*5	*9	*
652.6	*9	—	*9	—	*9	—	—	*	*	*	*
652.61	*8	—	*8	—	*8	—	—	*	*	*	*
652.8	37	—	37	*	37	*	—	9	*7	9	11
652.81	37	—	37	*	37	*	—	9	*7	9	11

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
653	154	—	154	*	154	*	—	26	25	64	38
653.4	123	—	123	*	123	*	—	21	21	53	29
653.41	123	—	123	*	123	*	—	21	21	53	29
653.5	22	—	22	—	22	*	—	*	*	*8	*7
653.51	22	—	22	—	22	*	—	*	*	*8	*7
654	558	—	558	*	557	*	—	89	114	216	139
654.0	*8	—	*8	—	*8	—	—	*	*	*	*
654.01	*7	—	*7	—	*7	—	—	*	*	*	*
654.1	24	—	24	—	24	*	—	*6	*	*8	*7
654.11	21	—	21	—	21	*	—	*5	*	*6	*6
654.2	468	—	468	—	467	*	—	68	99	183	118
654.21	458	—	458	—	457	*	—	66	97	180	115
654.23	9	—	9	—	9	—	—	*	*	*	*
654.4	17	—	17	—	17	—	—	*6	*	*6	*
654.41	13	—	13	—	13	—	—	*	*	*	*
654.5	23	—	23	*	23	*	—	*5	*	*9	*
654.51	15	—	15	*	15	—	—	*	*	*6	*
654.53	*7	—	*7	—	*7	*	—	*	*	*	*
654.6	*9	—	*9	—	*9	—	—	*	*	*	*
654.61	*8	—	*8	—	*8	—	—	*	*	*	*
655	21	—	21	*	21	—	—	*	*	*6	*6
655.8	10	—	10	*	10	—	—	*	*	*	*
655.81	*8	—	*8	*	*8	—	—	*	*	*	*
656	602	—	602	*	599	*	—	120	129	190	163
656.1	65	—	65	—	65	—	—	*8	20	23	15
656.11	63	—	63	—	63	—	—	*8	19	23	14
656.3	342	—	342	*	340	*	—	72	64	112	94
656.31	339	—	339	*	338	*	—	71	63	111	94
656.4	18	—	18	—	18	—	—	*	*	*8	*
656.41	17	—	17	—	17	—	—	*	*	*8	*
656.5	56	—	56	*	56	—	—	10	10	19	17
656.51	54	—	54	*	53	—	—	9	10	19	16
656.6	83	—	83	—	83	—	—	16	20	21	25
656.61	82	—	82	—	82	—	—	16	20	21	25
656.7	22	—	22	*	22	—	—	9	*	*	*
656.71	22	—	22	*	21	—	—	9	*	*	*
656.8	13	—	13	*	13	—	—	*	*7	*	*
656.81	12	—	12	*	12	—	—	*	*7	*	*
657	17	—	17	—	17	—	—	*	*	*6	*
657.0	17	—	17	—	17	—	—	*	*	*6	*
657.01	14	—	14	—	14	—	—	*	*	*	*
658	429	—	429	*	426	*	—	82	81	143	122
658.0	101	—	101	*	100	—	—	18	19	38	26
658.01	97	—	97	*	96	—	—	17	18	36	25
658.1	172	—	172	*	171	*	—	35	33	54	50
658.11	164	—	164	*	163	*	—	33	31	52	48
658.13	*7	—	*7	—	*7	—	—	*	*	*	*
658.2	70	—	70	*	69	—	—	13	15	21	21
658.21	67	—	67	*	66	—	—	12	15	19	21
658.4	78	—	78	*	77	*	—	16	11	29	22
658.41	76	—	76	*	75	*	—	16	11	28	21
658.8	*6	—	*6	—	*6	—	—	—	*	*	*
658.81	*6	—	*6	—	*6	—	—	—	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
659	194	—	194	*	193	*	—	32	43	54	65
659.1	41	—	41	—	40	*	—	*6	10	12	13
659.11	35	—	35	—	35	*	—	*5	*9	10	11
659.13	*6	—	*6	—	*6	—	—	*	*	*	*
659.2	45	—	45	*	44	*	—	*8	10	16	10
659.21	44	—	44	*	43	*	—	*8	10	16	10
659.4	14	—	14	—	14	*	—	*	*	*	*
659.41	14	—	14	—	14	*	—	*	*	*	*
659.5	*7	—	*7	—	*6	*	—	*	*	*	*
659.51	*6	—	*6	—	*6	*	—	*	*	*	*
659.6	43	—	43	—	43	*	—	*	*7	12	20
659.61	41	—	41	—	41	*	—	*	*6	11	20
659.8	37	—	37	—	37	—	—	*8	9	*8	12
659.81	32	—	32	—	32	—	—	*7	9	*	11
659.83	*6	—	*6	—	*6	—	—	*	*	*	*
659.9	*5	—	*5	—	*5	—	—	*	*	*	*
660	244	—	244	*	243	*	—	30	56	72	86
660.0	47	—	47	*	47	*	—	*8	*8	12	19
660.01	47	—	47	*	47	*	—	*8	*8	12	19
660.1	50	—	50	*	50	*	—	*6	*9	14	21
660.11	50	—	50	*	50	*	—	*6	*9	14	21
660.3	48	—	48	*	48	—	—	*5	14	13	16
660.31	48	—	48	*	48	—	—	*5	14	13	16
660.4	59	—	59	—	59	—	—	*5	15	19	20
660.41	59	—	59	—	59	—	—	*5	15	19	20
660.6	21	—	21	*	20	—	—	*	*7	*7	*
660.61	20	—	20	*	20	—	—	*	*7	*7	*
660.7	13	—	13	*	13	—	—	*	*	*5	*
660.71	13	—	13	*	13	—	—	*	*	*5	*
661	383	—	383	*	381	*	—	73	79	128	103
661.0	78	—	78	*	78	—	—	13	16	24	26
661.01	77	—	77	*	77	—	—	12	16	24	25
661.1	68	—	68	*	68	—	—	16	11	26	15
661.11	68	—	68	*	68	—	—	16	11	26	15
661.2	124	—	124	*	123	*	—	26	28	45	25
661.21	124	—	124	*	123	*	—	26	28	45	25
661.3	93	—	93	*	92	—	—	15	20	27	30
661.31	92	—	92	*	92	—	—	15	20	27	30
661.9	16	—	16	—	16	—	—	*	*	*6	*6
661.91	15	—	15	—	15	—	—	*	*	*6	*6
662	57	—	57	—	57	—	—	14	12	14	17
662.0	14	—	14	—	14	—	—	*	*	*	*
662.01	13	—	13	—	13	—	—	*	*	*	*
662.1	*7	—	*7	—	*7	—	—	*	*	*	*
662.11	*7	—	*7	—	*7	—	—	*	*	*	*
662.2	36	—	36	—	36	—	—	10	*9	*8	9
662.21	36	—	36	—	36	—	—	10	*9	*8	9
663	664	—	664	*	662	*	—	128	132	203	201
663.0	13	—	13	—	13	—	—	*	*	*	*
663.01	13	—	13	—	13	—	—	*	*	*	*
663.1	95	—	95	*	95	*	—	18	20	24	33
663.11	95	—	95	*	95	*	—	18	20	24	33
663.2	30	—	30	—	30	*	—	*6	*6	9	9

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
663.21	30	—	30	—	30	*	—	*6	*6	9	9
663.3	490	—	490	*	489	*	—	94	98	153	144
663.31	490	—	490	*	489	*	—	94	98	153	144
663.4	13	—	13	—	13	—	—	*	*	*	*
663.41	13	—	13	—	13	—	—	*	*	*	*
663.8	15	—	15	*	15	—	—	*	*	*7	*
663.81	15	—	15	*	15	—	—	*	*	*7	*
664	1,043	—	1,043	*	1,041	*	—	180	225	325	314
664.0	434	—	434	*	433	*	—	80	96	135	123
664.01	433	—	433	*	432	*	—	80	96	134	122
664.1	401	—	401	*	401	—	—	67	85	114	135
664.11	400	—	400	*	400	—	—	67	85	113	134
664.2	123	—	123	*	123	—	—	21	28	42	33
664.21	123	—	123	*	123	—	—	21	28	42	33
664.3	59	—	59	*	59	—	—	10	13	24	12
664.31	59	—	59	*	59	—	—	10	13	24	12
664.4	11	—	11	—	11	—	—	*	*	*	*
664.41	11	—	11	—	11	—	—	*	*	*	*
664.9	9	—	9	—	9	—	—	*	*	*	*
664.91	9	—	9	—	9	—	—	*	*	*	*
665	160	—	160	*	160	—	—	25	49	51	36
665.3	11	—	11	—	11	—	—	*	*	*	*
665.31	11	—	11	—	11	—	—	*	*	*	*
665.4	46	—	46	*	45	—	—	*8	10	16	11
665.41	46	—	46	*	45	—	—	*8	10	16	11
665.5	91	—	91	*	91	—	—	13	31	27	20
665.51	91	—	91	*	91	—	—	13	31	27	20
666	93	—	93	*	93	—	—	14	15	31	33
666.0	19	—	19	—	19	—	—	*	*	*7	*6
666.02	18	—	18	—	18	—	—	*	*	*6	*6
666.1	57	—	57	*	56	—	—	*8	10	16	22
666.12	56	—	56	*	56	—	—	*8	10	16	21
666.2	11	—	11	*	11	—	—	*	*	*	*
666.22	*7	—	*7	*	*7	—	—	*	*	*	*
666.3	*6	—	*6	—	*6	—	—	*	*	*	*
666.32	*6	—	*6	—	*6	—	—	*	*	*	*
667	20	—	20	—	20	—	—	*	*	*5	*5
667.0	12	—	12	—	12	—	—	*	*	*	*
667.02	11	—	11	—	11	—	—	*	*	*	*
667.1	*8	—	*8	—	*8	—	—	*	*	*	*
667.12	*7	—	*7	—	*7	—	—	*	*	*	*
668	*6	—	*6	—	*6	—	—	*	*	*	*
669	183	—	183	*	182	—	—	18	52	66	47
669.4	*9	—	*9	—	*9	—	—	*	*	*	*
669.42	*6	—	*6	—	*6	—	—	*	*	*	*
669.5	110	—	110	*	109	—	—	*9	40	41	21
669.51	110	—	110	*	109	—	—	*9	40	41	21
669.7	10	—	10	—	10	—	—	*	*	*6	*
669.71	10	—	10	—	10	—	—	*	*	*6	*
669.8	45	—	45	—	45	—	—	*6	*8	12	19
669.81	40	—	40	—	40	—	—	*6	*7	10	18

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
670	33	—	33	*	32	—	—	*	*8	13	*8
670.0	33	—	33	*	32	—	—	*	*8	13	*8
670.02	24	—	24	*	24	—	—	*	*6	10	*6
670.04	*8	—	*8	—	*8	—	—	*	*	*	*
671	25	—	25	*	25	*	—	*7	*6	*8	*
671.8	13	—	13	—	13	*	—	*	*	*	*
671.81	*6	—	*6	—	*6	*	—	*	*	*	*
671.82	*7	—	*7	—	*7	*	—	*	*	*	*
672	12	—	12	—	12	—	—	*	*	*	*
672.0	12	—	12	—	12	—	—	*	*	*	*
672.02	11	—	11	—	11	—	—	*	*	*	*
674	36	—	36	—	35	*	—	*6	10	12	*7
674.3	11	—	11	—	11	—	—	*	*	*	*
674.32	*8	—	*8	—	*8	—	—	*	*	*	*
674.8	18	—	18	—	18	*	—	*	*7	*6	*
674.82	15	—	15	—	15	—	—	*	*5	*6	*
676	*6	—	*6	*	*5	—	—	*	*	*	*
680–709	1,531	747	784	125	330	360	716	363	391	503	274
680	*6	*	*	*	*	*	*	*	*	*	*
681	50	28	21	*	12	13	20	11	12	17	10
681.0	20	13	*7	*	*9	*5	*	*6	*	*5	*
681.00	17	11	*6	*	*8	*	*	*	*	*5	*
681.1	29	15	14	*	*	*7	16	*	*8	11	*6
681.10	25	13	11	*	*	*7	13	*	*7	*8	*5
682	620	321	299	38	161	167	255	143	156	201	121
682.0	33	19	14	*9	10	*8	*6	*8	*7	*8	10
682.1	10	*	*5	*	*5	*	*	*	*	*	*
682.2	63	28	35	*	19	20	20	11	16	20	16
682.3	77	41	36	*	35	17	21	18	19	23	17
682.4	30	19	12	*	11	*6	10	10	*6	10	*
682.5	13	*6	*7	*	*6	*	*	*	*	*	*
682.6	271	129	142	*9	52	76	134	63	71	89	47
682.7	104	64	39	*	16	30	54	24	29	35	16
682.8	*7	*	*	*	*	*	*	*	*	*	*
682.9	11	*6	*6	*	*	*	*	*	*	*6	*
683	*8	*	*	*6	*	*	*	*	*	*	*
685	*6	*	*	*	*	*	*	*	*	*	*
686	16	*8	*9	*	*6	*	*	*	*	*6	*
686.9	11	*5	*6	*	*	*	*	*	*	*	*
690	*8	*6	*	*	*	*	*	*	*	*	*
690.1	*6	*	*	*	*	*	*	*	*	*	*
691	36	18	18	33	*	*	*	*7	10	*9	10
691.0	31	15	16	30	*	—	*	*6	*9	*7	9
691.8	*5	*	*	*	*	*	*	*	*	*	*
692	43	21	22	10	10	10	13	13	12	13	*5
692.9	35	18	18	9	*8	*7	11	10	10	11	*
693	46	22	24	*7	11	9	19	13	11	13	9
693.0	44	20	24	*6	11	*9	19	13	11	12	9
694	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
695	28	12	16	*	*6	10	*8	*7	*7	*8	*5
695.1	*7	*	*	*	*	*	*	*	*	*	*
695.4	*5	*	*	—	*	*	*	*	*	*	*
695.8	*6	*	*	*	*	*	*	*	*	*	*
695.89	*6	*	*	*	*	*	*	*	*	*	*
695.9	*5	*	*	*	*	*	*	*	*	*	*
696	32	20	12	*	*7	10	14	10	*8	*8	*6
696.0	*6	*	*	—	*	*	*	*	*	*	*
696.1	25	16	9	*	*6	*8	11	*7	*6	*7	*
698	21	*8	12	*	*8	*	*7	*	*6	*8	*
698.9	15	*7	*8	*	*6	*	*	*	*	*7	*
701	19	*	14	*	*8	*	*6	*	*	*8	*
701.8	*6	*	*5	*	*	*	*	*	*	*	*
702	10	*	*5	*	*	*	*8	*	*	*6	*
702.1	*7	*	*	—	*	*	*	*	*	*	*
702.19	*5	*	*	—	*	*	*	*	*	*	*
703	13	*7	*6	*	*	*	*8	*	*	*	*
703.0	9	*6	*	*	*	*	*	*	*	*	*
704	*8	*	*	*	*	*	*	*	*	*	*
705	*6	*	*	*	*	*	*	*	*	*	*
706	23	11	12	*	*9	*6	*6	*	*7	*5	*5
706.1	*8	*5	*	*	*5	*	*	*	*	*	*
706.2	12	*	*7	*	*	*	*	*	*	*	*
707	462	208	254	*	46	92	321	110	120	164	67
707.0	256	108	149	*	23	34	198	62	60	101	34
707.1	189	92	97	*	21	54	113	45	56	57	31
707.8	11	*5	*6	*	*	*	*6	*	*	*	*
707.9	*5	*	*	—	*	*	*	*	*	*	*
708	19	*8	11	*	*7	*	*	*	*7	*6	*
708.0	11	*5	*6	*	*	*	*	*	*	*	*
708.9	*6	*	*	*	*	*	*	*	*	*	*
709	37	15	22	*	15	9	*9	10	10	11	*7
709.0	*5	*	*	*	*	*	*	*	*	*	*
709.2	15	*5	10	*	*8	*	*	*	*	*	*
709.8	*7	*	*	*	*	*	*	*	*	*	*
709.9	*7	*	*	*	*	*	*	*	*	*	*
710–739	4,541	1,617	2,924	64	774	1,071	2,633	924	1,186	1,590	841
710	145	22	123	*	60	40	42	25	21	68	32
710.0	104	12	92	*	49	27	26	18	13	50	24
710.1	16	*	12	—	*	*6	*5	*	*	*6	*
710.2	*7	*	*5	—	*	*	*	*	*	*	*
710.4	*6	*	*	—	*	*	*	*	*	*	*
710.8	*5	*	*	—	*	*	*	—	*	*	*
711	30	17	13	*	10	*6	10	*6	*6	12	*6
711.0	27	16	11	*	*8	*6	10	*5	*6	11	*
711.06	9	*6	*	*	*	*	*	*	*	*	*
712	11	*6	*6	—	*	*	*9	*5	*	*	*
712.3	*8	*	*	—	*	*	*6	*	*	*	*
713	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
714	211	60	151	*	16	49	144	44	60	68	39
714.0	196	55	141	*	12	48	136	39	54	66	37
714.3	*7	*	*6	*	*	*	*	*	*	*	*
714.30	*6	*	*5	*	*	*	*	*	*	*	*
715	1,153	367	786	*	36	205	910	220	350	383	199
715.0	50	14	36	—	*	*	46	11	13	17	10
715.00	*8	*	*6	—	*	*	*7	*	*	*5	*
715.09	42	12	29	—	*	*	39	10	11	12	*9
715.1	44	16	29	—	*	9	31	*8	12	18	*7
715.15	13	*	*8	—	*	*	9	*	*	*	*
715.16	28	10	18	—	*	*5	20	*	*8	12	*
715.2	*6	*	*	—	*	*	*	*	*	*	*
715.3	223	85	138	*	*8	53	162	46	71	65	41
715.31	14	*6	*8	—	*	*	10	*	*	*	*
715.35	62	26	36	*	*	13	46	13	21	14	14
715.36	138	50	88	—	*	33	101	27	43	44	23
715.8	53	17	35	—	*	*7	45	9	18	15	10
715.89	52	17	35	—	*	*7	45	9	18	15	10
715.9	777	232	545	*	22	131	624	145	236	267	130
715.90	453	123	330	*	*7	63	382	85	138	167	63
715.91	18	*7	11	—	*	*	11	*	*7	*	*
715.95	90	34	56	—	*5	19	66	20	29	24	16
715.96	188	60	128	—	*7	38	143	30	51	63	44
715.98	17	*5	12	—	*	*	14	*	*5	*6	*
716	199	70	129	*	22	50	127	40	55	76	28
716.1	9	*6	*	—	*	*	*	*	*	*	*
716.5	10	*	*8	*	*	*	*	*	*	*	*
716.59	9	*	*8	*	*	*	*	*	*	*	*
716.6	*7	*	*5	*	*	*	*	*	*	*	*
716.9	171	60	111	*	12	43	116	34	48	65	24
716.90	119	41	78	—	*7	27	86	22	34	47	16
716.91	*7	*	*	—	*	*	*	*	*	*	*
716.96	17	*6	11	*	*	*7	*9	*	*	*6	*
716.99	13	*	9	*	*	*	*9	*	*	*6	*
717	45	27	18	*	27	11	*6	18	*8	11	*7
717.4	*6	*	*	*	*	*	*	*	*	*	*
717.7	*9	*	*	*	*	*	*	*	*	*	*
717.8	18	13	*5	*	16	*	*	10	*	*	*
717.83	12	*8	*	*	11	*	*	*6	*	*	*
718	58	27	32	*6	25	11	17	18	13	17	10
718.3	14	*7	*7	*	*9	*	*	*	*	*	*
718.31	*8	*	*	*	*5	*	*	*	*	*	*
718.4	22	9	13	*	*6	*	10	*6	*5	*8	*
718.46	*5	*	*	*	*	*	*	*	*	*	*
718.49	*6	*	*	*	*	*	*	*	*	*	*
718.8	*8	*	*	*	*5	*	*	*	*	*	*
719	143	56	87	*	28	24	87	29	43	46	25
719.0	13	*8	*5	*	*	*	*8	*	*	*	*
719.06	11	*6	*	*	*	*	*6	*	*	*	*
719.4	77	32	45	*	19	15	41	11	25	22	18
719.40	*5	*	*	*	*	*	*	*	*	*	*
719.41	21	*7	14	*	*	*5	14	*	*8	*5	*6
719.45	32	15	17	*	*6	*	20	*	12	10	*7

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
719.46	10	*	*5	*	*6	*	*	*	*	*	*
719.7	37	10	27	—	*	*	32	11	11	12	*
719.70	*8	*	*6	—	*	*	*8	*	*	*	*
719.75	14	*	11	—	*	*	12	*	*	*6	*
719.76	*8	*	*	—	*	*	*5	*	*	*	*
720	18	12	*6	*	*	*8	*7	*6	*	*8	*
720.0	12	10	*	—	*	*7	*	*5	*	*	*
721	218	97	121	*	24	62	130	36	62	86	34
721.0	53	28	25	*	*6	21	26	*7	14	24	*9
721.1	19	11	*8	—	*	*8	*8	*	*6	*7	*
721.2	*6	*	*	—	*	*	*	*	*	*	*
721.3	82	33	49	*	*9	17	56	15	27	28	12
721.9	50	19	31	*	*	11	35	*9	11	21	*8
721.90	50	19	31	*	*	11	35	*9	11	21	*8
722	473	244	229	*	189	174	110	89	108	194	82
722.0	63	35	29	*	30	29	*	9	13	34	*7
722.1	243	134	109	*	115	88	40	51	54	97	42
722.10	239	132	108	*	114	87	39	50	52	95	41
722.2	*7	*	*	—	*	*	*	*	*	*	*
722.4	19	*8	11	—	*	*6	*9	*	*	10	*
722.5	71	30	41	—	12	24	35	9	22	24	17
722.52	69	30	39	—	12	23	33	*9	20	23	17
722.6	*9	*	*6	—	*	*	*	*	*	*	*
722.7	20	12	*8	—	*5	*8	*6	*5	*	*7	*
722.71	11	*7	*	—	*	*6	*	*	*	*	*
722.73	*8	*	*	—	*	*	*	*	*	*	*
722.8	22	10	12	—	10	*7	*	*	*	10	*
722.83	19	9	10	—	10	*6	*	*	*	*9	*
722.9	19	10	*9	—	*6	*7	*6	*	*	*8	*
722.93	14	*7	*7	—	*	*5	*	*	*	*6	*
723	56	26	30	*	16	24	16	11	14	19	12
723.0	16	9	*7	—	*	*8	*5	*	*5	*	*
723.1	18	*5	13	*	*6	*6	*5	*	*	*6	*6
723.4	17	10	*8	*	*	9	*	*6	*	*6	*
724	338	136	202	*	75	99	163	60	97	108	73
724.0	102	42	60	*	*6	25	71	21	27	33	21
724.00	15	*	12	—	—	*	14	*	*6	*	*
724.02	84	39	45	*	*6	24	54	19	21	27	17
724.2	92	36	56	*	28	33	31	13	27	30	23
724.3	16	*6	10	—	*	*	*8	*	*	*	*
724.4	36	14	22	—	10	13	13	*8	10	11	*7
724.5	70	26	44	*	17	18	34	10	20	24	16
724.6	*6	*	*	—	*	*	*	*	*	*	*
724.8	9	*5	*	*	*	*	*	*	*	*	*
725	41	*6	35	—	—	*	39	10	12	*9	10
726	107	54	53	*	25	40	40	31	26	26	24
726.0	*6	*	*	—	*	*	*	*	*	*	*
726.1	27	13	14	—	*	10	13	*7	*7	*6	*7
726.10	23	11	12	—	*	*7	12	*5	*7	*	*7
726.2	26	12	14	—	*7	12	*7	9	*6	*	*6
726.3	14	10	*	*	*6	*5	*	*	*	*	*
726.33	12	*9	*	*	*5	*	*	*	*	*	*
726.5	12	*	11	—	*	*	*8	*	*	*6	*
726.6	*8	*7	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
726.65	*6	*5	*	*	*	*	*	*	*	*	*
726.7	*5	*	*	*	*	*	*	*	*	*	*
726.9	*8	*6	*	*	*	*	*	*	*	*	*
726.91	*6	*	*	—	*	*	*	*	*	*	*
727	60	25	35	*	22	18	17	16	12	17	15
727.0	32	13	18	*	13	10	*7	*9	*8	10	*5
727.00	13	*6	*8	*	*	*	*	*	*	*	*
727.05	10	*	*5	*	*6	*	*	*	*	*	*
727.4	*7	*	*	—	*	*	*	*	*	*	*
727.8	*8	*	*	*	*	*	*	*	*	*	*
728	81	40	40	*	23	23	32	23	14	27	16
728.8	52	26	25	*	17	17	16	15	*8	15	14
728.85	14	*6	*8	*	*6	*	*	*	*	*	*6
728.86	*6	*	*	—	*	*	*	*	*	*	*
728.89	28	16	12	*	*9	*8	11	9	*5	*7	*6
728.9	11	*7	*	*	*	*	*6	*	*	*	*
729	166	54	113	*	47	53	65	27	49	54	35
729.1	54	10	44	*	21	20	13	10	17	18	9
729.2	10	*	*7	—	*	*	*	*	*	*	*
729.5	46	18	29	*	13	15	18	*9	17	11	10
729.8	48	20	28	*	*8	12	28	*7	13	19	10
729.81	*8	*	*5	*	*	*	*	*	*	*	*
729.82	*7	*	*	—	*	*	*	*	*	*	*
729.89	34	16	18	*	*	*7	22	*5	*8	13	*8
730	111	59	52	*	22	33	52	28	25	38	20
730.0	33	20	14	*	*6	10	15	10	*7	10	*6
730.07	17	11	*6	*	*	*7	*8	*6	*	*5	*
730.08	*5	*	*	*	*	*	*	*	*	*	*
730.1	26	13	14	*	*6	*6	13	*7	*7	*8	*5
730.15	*6	*	*	*	*	*	*	*	*	*	*
730.17	*9	*	*	*	*	*	*	*	*	*	*
730.18	*6	*	*	*	*	*	*	*	*	*	*
730.2	47	22	25	*	*8	15	22	11	10	18	*9
730.26	*6	*	*	*	*	*	*	*	*	*	*
730.27	23	10	13	*	*	*7	13	*	*6	*9	*
730.28	*8	*	*	*	*	*	*	*	*	*	*
731	36	20	16	—	*	*7	27	9	12	11	*
731.0	19	9	10	—	—	*	18	*6	*7	*	*
731.8	17	11	*6	—	*	*6	9	*	*	*7	*
732	11	*6	*	*7	*	*	*	*	*	*	*
733	641	127	514	*	57	93	488	122	149	245	125
733.0	297	33	265	*	*	22	272	54	71	112	60
733.00	264	30	234	*	*	18	244	48	65	103	48
733.01	20	*	18	—	—	*	19	*	*	*	*9
733.09	12	*	11	*	*	*	*8	*	*	*	*
733.1	182	36	145	*	*7	21	153	34	40	72	35
733.10	*5	*	*	—	*	*	*	*	*	*	*
733.11	*5	*	*	—	*	*	*	*	*	*	*
733.13	116	25	92	*	*	11	103	23	28	47	18
733.14	18	*	14	*	*	*	13	*	*	*6	*
733.15	11	*	*9	*	*	*	*8	*	*	*	*
733.19	22	*	18	*	*	*	19	*	*	9	*6
733.4	46	20	26	*	14	16	16	*9	10	20	*7
733.42	35	15	19	*	10	13	12	*6	*6	18	*
733.49	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
— Quantity zero.NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
733.6	29	*8	20	*	*7	11	10	*7	*9	10	*
733.8	47	17	29	*	17	14	16	9	*7	21	9
733.81	*8	*	*	*	*	*	*	*	*	*	*
733.82	39	14	25	*	13	12	14	*7	*6	17	*8
733.9	38	12	26	*	*8	*9	20	*8	12	10	*9
733.90	23	*6	17	*	*	*	16	*	*7	*6	*5
733.99	12	*	*8	*	*	*	*	*	*	*	*
735	30	*6	24	*	*5	*7	17	*7	*5	11	*6
735.0	12	*	11	—	*	*	*8	*	*	*7	*
735.4	10	*	*8	*	*	*	*6	*	*	*	*
736	41	17	24	*5	10	*8	18	12	*9	13	*6
736.4	*8	*	*	*	*	*	*	*	*	*	*
736.7	20	*7	12	*	*	*	*8	*8	*	*	*
736.79	16	*6	10	*	*	*	*7	*7	*	*	*
736.8	*7	*	*	*	*	*	*	*	*	*	*
737	65	16	49	*6	13	*6	38	14	15	24	12
737.1	12	*	*9	*	*	*	10	*	*	*5	*
737.10	10	*	*7	*	*	*	*9	*	*	*	*
737.3	45	11	34	*6	11	*	23	11	10	15	*8
737.30	39	*9	30	*	11	*	20	10	9	14	*6
737.4	*6	*	*5	*	*	*	*5	*	*	*	*
738	36	14	21	*	*8	*9	17	*8	*9	10	*9
738.4	28	11	17	*	*	*7	15	*5	*8	*7	*7
739	9	*	*6	*	*	*	*	*	*	*	*
740–759	577	294	283	333	111	71	62	195	126	141	114
741	21	12	9	9	10	*	*	*7	*6	*	*
741.0	9	*5	*	*5	*	—	—	*	*	*	*
741.9	12	*6	*5	*	*6	*	*	*	*	*	*
741.90	*8	*	*	*	*	*	*	*	*	*	*
742	25	12	13	18	*	*	*	11	*7	*6	*
742.3	9	*5	*	*6	*	*	*	*	*	*	*
742.5	*5	*	*	*	*	*	*	*	*	*	*
743	*6	*	*	*	*	*	*	*	*	*	*
744	*8	*	*	*6	*	*	*	*	*	*	*
745	82	42	40	63	11	*	*	33	15	22	12
745.1	*8	*6	*	*6	*	—	—	*	*	*	—
745.2	10	*5	*5	9	*	*	—	*7	*	*	*
745.4	29	14	15	23	*	*	*	10	*5	*9	*5
745.5	26	12	14	16	*	*	*	*8	*6	*8	*
745.6	*5	*	*	*	*	—	—	*	*	*	*
746	58	30	28	34	10	*8	*5	19	11	15	12
746.0	*6	*	*	*5	*	*	*	*	*	*	*
746.4	10	*8	*	*	*	*	*	*	*	*	*
746.8	23	11	12	16	*	*	*	*8	*	*7	*
746.89	12	*5	*6	10	*	*	—	*	*	*	*
746.9	*7	*	*	*	*	*	*	*	*	*	*
747	64	32	32	41	*8	*6	*8	23	11	17	13
747.0	16	9	*6	15	*	—	—	*	*	*	*
747.1	*6	*	*	*	*	*	*	*	*	*	*
747.10	*5	*	*	*	*	*	*	*	*	*	*
747.2	*7	*	*	*	*	*	*	*	*	*	*
747.22	*5	*	*	*	*	*	*	*	*	*	*
747.3	15	*6	9	11	*	*	*	10	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
747.6	*6	*	*	*	*	*	*	*	*	*	*
747.8	9	*	*6	*	*	*	*	*	*	*	*
747.81	*6	*	*	*	*	*	*	*	*	*	*
748	13	*6	*7	11	*	—	—	*5	*	*	*
748.3	*7	*	*	*7	—	—	—	*	*	*	*
749	17	11	*6	16	*	—	*	*5	*	*	*6
749.0	*7	*	*	*6	*	—	*	*	*	*	*
749.2	*8	*5	*	*7	*	—	—	*	*	*	*
750	18	14	*	15	*	*	*	*6	*6	*	*
750.5	11	*9	*	11	—	—	—	*	*	*	*
751	31	17	14	18	*	*	*	11	*7	*6	*8
751.0	*7	*	*	*	*	*	*	*	*	*	*
751.2	*5	*	*	*5	—	—	—	*	*	*	*
751.6	*8	*	*5	*	*	*	*	*	*	*	*
752	31	*8	23	*8	14	*8	*	*8	*9	10	*
752.1	16	—	16	*	*9	*6	*	*	*6	*6	*
752.11	15	—	15	*	*8	*6	*	*	*6	*6	*
753	51	25	26	17	*8	10	16	13	13	13	12
753.0	*6	*	*	*	*	*	*	*	*	*	*
753.1	27	12	14	*	*	*8	12	*	*8	*8	*6
753.10	*8	*	*	*	*	*	*6	*	*	*	*
753.12	10	*	*6	*	*	*	*	*	*	*	*
753.2	*8	*5	*	*6	*	*	*	*	*	*	*
754	17	*8	*9	12	*	*	*	*6	*	*	*
755	16	*9	*8	*9	*	*	*	*8	*	*	*
755.6	9	*	*	*	*	*	*	*	*	*	*
755.63	*5	*	*	*	*	*	*	*	*	*	*
756	48	26	21	21	*8	11	*8	13	10	14	10
756.0	10	*7	*	9	*	—	—	*	*	*	*
756.1	23	12	11	*	*	9	*7	*5	*	*9	*
756.12	12	*	*7	*	*	*5	*	*	*	*	*
756.19	*5	*	*	*	*	*	*	*	*	*	*
756.5	*5	*	*	*	*	*	*	*	*	*	*
757	*7	*	*5	*	*	*	*	*	*	*	*
758	31	19	13	17	*9	*5	*	9	*8	*8	*6
758.0	24	15	9	12	*8	*	*	*7	*6	*6	*5
759	31	14	17	13	9	*	*	14	*6	*7	*5
759.8	17	*8	*9	*8	*6	*	*	10	*	*	*
759.89	13	*6	*7	*6	*5	*	*	*8	*	*	*
760–779	416	247	169	412	*	*	*	75	109	109	123
760	*5	*	*	*	*	*	*	*	*	*	*
765	43	26	17	43	—	—	—	*6	11	12	14
765.0	*8	*	*	*8	—	—	—	*	*	*	*
765.1	35	21	14	35	—	—	—	*5	10	9	11
765.18	11	*6	*	11	—	—	—	*	*	*	*
765.19	*8	*6	*	*8	—	—	—	*	*	*	*
767	9	*5	*	9	—	—	—	*	*	*	*
769	27	16	11	27	—	—	—	*	*5	*8	11

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
770	85	51	34	85	—	—	—	15	27	23	20
770.1	*7	*	*	*7	—	—	—	*	*	*	*
770.2	*7	*5	*	*7	—	—	—	*	*	*	*
770.6	11	*7	*	11	—	—	—	*	*	*	*
770.7	19	12	*7	19	—	—	—	*	*8	*	*
770.8	34	19	15	34	—	—	—	*7	9	10	*8
771	37	23	14	37	*	—	*	*8	9	*8	12
771.8	29	18	11	28	—	—	*	*6	*7	*7	*9
772	*8	*	*	*8	—	—	—	*	*	*	*
773	*8	*	*	*8	—	—	—	*	*	*	*
773.1	*6	*	*	*6	—	—	—	*	*	*	*
774	75	43	32	75	—	—	—	12	19	22	22
774.2	24	15	10	24	—	—	—	*	*6	*6	*8
774.6	47	27	20	47	—	—	—	*8	12	15	13
775	31	21	10	31	—	—	—	*6	*7	*9	*8
775.5	13	*8	*	13	—	—	—	*	*	*	*
775.6	*6	*	*	*6	—	—	—	*	*	*	*
775.7	*5	*	*	*5	—	—	—	*	*	*	*
776	17	*9	*8	16	—	—	*	*	*	*6	*
776.6	*7	*	*	*7	—	—	—	*	*	*	*
777	*9	*5	*	*9	*	—	—	*	*	*	*
778	17	11	*6	16	*	*	—	*	*6	*	*
778.4	*6	*	*	*6	—	—	—	*	*	*	*
779	30	17	14	30	*	—	*	*5	*8	*6	12
779.0	*7	*	*	*7	—	—	—	*	*	*	*
779.3	17	10	*7	17	—	—	—	*	*5	*	*7
780–799	5,998	2,694	3,303	512	1,373	1,487	2,626	1,288	1,552	2,077	1,080
780	1,750	824	926	187	377	399	787	417	440	585	307
780.0	76	30	46	*	20	13	40	14	16	28	18
780.01	20	*8	12	*	*	*	11	*	*	*7	*6
780.09	49	19	30	*	14	*7	25	9	11	18	10
780.1	13	*7	*6	*	*	*	*8	*	*	*5	*
780.2	319	144	175	*	41	60	214	87	73	106	51
780.3	711	356	355	89	189	181	252	174	177	244	116
780.4	99	33	66	*	16	20	62	20	31	31	18
780.5	102	63	39	10	26	37	30	28	23	29	22
780.52	12	*5	*7	*	*	*	*	*	*	*	*
780.57	81	54	28	*8	21	31	22	25	18	23	16
780.6	268	127	141	74	61	55	78	56	63	96	52
780.7	106	41	66	*	13	19	71	22	39	26	19
780.9	54	21	33	*	*9	11	32	13	13	20	*8
781	137	59	78	*7	14	25	91	25	48	47	17
781.0	15	*5	10	*	*	*	*6	*	*5	*	*
781.2	72	30	42	*	*	11	57	13	29	24	*6
781.3	41	20	20	*	*	*8	26	*7	11	16	*7
781.9	*5	*	*	*	*	*	*	*	*	*	*
782	157	66	91	17	36	40	64	31	50	44	32
782.0	26	12	15	*	9	*9	*8	*	*8	*9	*6
782.1	35	13	22	*8	10	*8	*9	*8	10	10	*6
782.3	60	21	39	*	*7	15	37	13	21	16	10
782.4	21	14	*8	*	*8	*7	*5	*	*7	*6	*
782.7	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
783	141	65	76	63	13	16	48	35	35	39	32
783.0	28	*8	20	*	*	*	19	*5	*7	13	*
783.2	33	13	20	*	*6	9	14	*	15	10	*
783.3	12	*7	*5	*9	*	*	*	*	*	*	*
783.4	63	35	28	48	*	*	11	19	11	11	22
784	329	132	197	10	70	84	166	63	91	102	73
784.0	115	36	78	*	51	37	24	22	25	43	24
784.2	*8	*	*	*	*	*	*	*	*	*	*
784.3	92	39	53	*	*	20	67	17	27	27	21
784.5	58	29	29	*	*	11	43	*7	20	14	16
784.7	42	21	20	*	*7	10	22	12	12	10	*8
785	272	138	135	17	44	71	140	67	67	93	46
785.0	44	18	26	*	13	10	17	*7	13	16	*8
785.1	23	*7	16	*	*	*7	12	*6	*6	*8	*
785.2	18	*7	12	*	*5	*	*5	*	*5	*8	*
785.4	71	44	27	*	*6	23	42	16	15	27	13
785.5	74	44	30	*	*5	17	48	24	16	22	11
785.50	*8	*6	*	*	*	*	*	*	*	*	*
785.51	32	18	14	*	*	*7	23	11	*7	10	*
785.59	34	20	14	*	*	*7	21	12	*8	10	*
785.6	33	15	18	*5	*9	*8	10	9	*8	9	*7
785.9	*9	*	*6	*	—	*	*6	*	*	*	*
786	1,019	487	531	61	204	380	374	197	249	402	171
786.0	142	65	78	46	20	25	51	21	35	54	33
786.01	*7	*	*	—	*	*	*	*	*	*	*
786.09	133	60	73	45	17	23	48	20	33	48	32
786.2	15	*6	9	*5	*	*	*	*	*	*	*
786.3	51	32	18	*	12	12	27	12	14	19	*5
786.5	774	364	411	*	166	332	273	153	186	314	121
786.50	610	286	324	*	124	271	213	111	148	258	93
786.51	22	13	*9	—	*7	*8	*7	*6	*5	*7	*
786.52	83	36	47	*	23	29	31	17	23	32	12
786.59	59	29	31	*	13	24	22	19	10	18	13
786.6	21	11	10	*	*	*5	14	*5	*6	*7	*
787	532	184	348	66	125	102	239	100	136	187	109
787.0	267	75	192	40	78	58	91	43	67	102	55
787.01	154	40	114	11	47	38	58	22	42	62	29
787.02	59	13	46	*	20	14	23	10	15	21	14
787.03	53	21	32	27	10	*5	11	12	10	19	13
787.2	114	52	62	*	*9	14	87	22	33	37	22
787.3	9	*	*5	*	*	*	*	*	*	*	*
787.6	9	*	*6	*	*	*	*	*	*	*	*
787.9	130	49	81	20	32	27	50	29	30	42	29
787.91	129	49	80	20	32	27	50	29	29	42	29
788	402	241	161	*8	67	75	252	98	114	122	68
788.0	25	14	11	*	13	*8	*	*9	*	9	*
788.1	*7	*	*	*	*	*	*	*	*	*	*
788.2	252	175	77	*	41	43	165	63	67	80	42
788.20	206	143	63	*	34	35	135	53	51	68	34
788.21	*6	*	*	*	*	*	*	*	*	*	*
788.29	40	27	12	*	*7	*6	26	9	14	11	*6
788.3	75	26	49	*	*8	13	50	14	28	19	14

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
788.30	57	20	37	*	*7	10	38	11	21	14	10
788.33	*6	—	*6	—	*	*	*	*	*	*	*
788.39	*6	*	*	*	*	*	*	*	*	*	*
788.4	11	*6	*	*	*	*	*6	*	*	*	*
788.41	*8	*	*	*	*	*	*	*	*	*	*
788.9	23	11	12	*	*	*	20	*7	*7	*5	*
789	442	168	275	33	157	125	127	94	120	150	78
789.0	314	100	214	30	128	73	83	61	93	103	57
789.00	118	35	82	13	42	26	35	17	30	50	21
789.01	31	10	22	*	13	*7	11	*8	10	*9	*
789.02	*7	*	*	*	*	*	*	*	*	*	*
789.03	53	19	35	*8	33	*8	*	15	14	12	12
789.04	16	*	12	*	*8	*	*	*	*6	*	*
789.05	*6	*	*	*	*	*	*	*	*	*	*
789.06	32	10	22	*	9	9	12	*6	11	10	*
789.07	14	*6	*8	*	*	*	*	*	*	*	*
789.09	36	10	26	*	13	10	11	*7	13	10	*6
789.1	*8	*	*	*	*	*	*	*	*	*	*
789.2	10	*6	*	*	*	*5	*	*	*	*	*
789.3	15	*	12	*	*7	*	*5	*	*	*6	*
789.30	*8	*	*7	—	*	*	*	*	*	*	*
789.5	93	54	39	*	19	39	33	24	18	35	16
790	308	119	189	12	105	66	125	62	81	102	64
790.2	90	11	79	*	69	*7	14	12	18	36	25
790.4	*8	*	*	*	*	*	*	*	*	*	*
790.5	26	13	13	*	*6	10	*9	*5	*7	*8	*6
790.6	108	49	59	*	13	31	62	25	30	34	19
790.7	33	16	17	*5	*8	*6	14	10	12	*8	*
790.9	34	21	13	*	*	*7	22	*7	*8	13	*6
790.92	20	10	10	*	*	*	12	*5	*5	*6	*
790.93	11	11	—	—	—	*	*9	*	*	*	*
791	20	*7	13	*5	*7	*	*	*	*6	*8	*
791.0	11	*	*7	*	*	*	*	*	*	*	*
792	55	*9	46	*	37	*	13	13	16	15	11
792.1	19	*8	11	*	*	*	13	*	*7	*5	*
792.3	35	—	35	*	35	—	—	*8	9	10	*8
793	13	*5	*8	*	*	*	*7	*	*	*	*
793.1	*8	*	*6	*	*	*	*	*	*	*	*
794	73	37	36	*	24	22	25	16	16	29	12
794.3	31	16	16	*	*9	*9	14	*8	*8	12	*
794.31	22	*9	13	*	*6	*5	10	*6	*	*8	*
794.39	*8	*5	*	—	*	*	*	*	*	*	*
794.8	34	19	15	*	13	12	*8	*7	*5	14	*8
795	31	14	18	*	21	*5	*	9	*6	*9	*8
795.5	22	11	11	*	13	*	*	*8	*	*	*6
796	39	17	22	*	14	*9	16	*6	14	13	*6
796.2	20	*6	14	*	*8	*6	*6	*	*8	*	*
796.3	*8	*6	*	*	*	*	*	*	*	*	*
796.4	*9	*	*5	*	*	*	*5	*	*	*	*
797	9	*	*6	—	—	—	9	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
799	265	119	146	18	56	56	135	46	59	123	37
799.0	99	41	58	9	16	23	50	12	33	40	14
799.1	16	*7	10	*	*	*	11	*	*	*7	*
799.3	23	9	14	—	*	*	19	*	9	*8	*
799.4	39	22	16	*	*8	9	21	11	*6	14	*6
799.8	*7	*	*	*	*	*	*	*	*	*	*
799.9	79	35	44	*5	26	16	31	16	*	49	9
800–999	5,919	3,095	2,824	436	2,027	1,189	2,267	1,280	1,402	1,994	1,243
800	15	12	*	*6	*7	*	*	*	*	*	*
800.0	*5	*	*	*	*	*	*	*	*	*	*
801	37	25	12	*7	17	*6	*6	*9	*7	12	*9
801.0	14	*8	*6	*	*6	*	*	*	*	*	*
801.01	*7	*	*	*	*	*	*	*	*	*	*
801.1	*5	*	*	*	*	*	*	*	*	*	*
801.2	11	*7	*	*	*6	*	*	*	*	*	*
802	114	84	30	*9	74	15	17	32	22	31	29
802.0	21	12	9	*	10	*	*7	*9	*	*	*
802.2	31	26	*	*	24	*	*	*6	*7	*9	9
802.22	*5	*	*	*	*	*	—	*	*	*	*
802.3	*5	*	*	*	*	*	*	*	*	*	*
802.4	20	15	*	*	14	*	*	*6	*	*7	*
802.6	15	10	*5	*	10	*	*	*5	*	*	*
802.8	14	10	*	*	*7	*	*	*5	*	*	*
803	10	*8	*	*	*6	*	*	*	*	*	*
805	130	66	65	*	43	22	64	19	35	43	33
805.0	28	19	9	*	13	*7	*7	*	*7	*7	11
805.02	9	*6	*	*	*	*	*5	*	*	*	*
805.2	35	16	18	*	*9	*	22	*6	*9	13	*8
805.4	53	25	28	*	16	*8	29	*7	16	17	13
805.6	11	*	*6	*	*	*	*	*	*	*5	*
806	10	*7	*	*	*7	*	*	*	*	*	*
807	119	70	50	*	41	27	48	20	30	42	28
807.0	104	60	44	*	36	23	42	17	27	35	25
807.00	*6	*	*	*	*	*	*	*	*	*	*
807.01	25	14	11	*	12	*	*8	*	*7	*8	*6
807.02	17	*9	*8	*	*5	*	*9	*	*	*6	*
807.03	18	11	*7	*	*7	*	*7	*	*	*6	*6
807.04	*7	*	*	*	*	*	*	*	*	*	*
807.05	*8	*6	*	—	*	*	*	*	*	*	*
807.09	18	11	*7	*	*5	*5	*7	*	*	9	*
807.2	*7	*	*	*	*	*	*	*	*	*	*
807.4	*6	*	*	—	*	*	*	*	*	*	*
808	87	34	53	*	25	16	44	19	23	30	16
808.0	20	11	9	*	*7	*5	*8	*6	*	*7	*
808.2	40	10	30	*	*9	*5	25	*8	11	13	*8
808.4	16	11	*	*	*6	*5	*	*	*	*5	*
808.8	10	*	*8	*	*	*	*7	*	*	*	*
810	24	13	12	*	11	*	*7	*	*5	9	*7
810.0	24	12	12	*	11	*	*7	*	*5	*9	*7
810.00	11	*	*7	*	*	*	*	*	*	*	*
810.02	*7	*5	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
811	12	*9	*	*	*7	*	*	*	*	*	*
811.0	12	*8	*	*	*7	*	*	*	*	*	*
811.00	*5	*	*	*	*	*	*	*	*	*	*
812	98	30	68	15	15	19	49	20	24	31	23
812.0	52	9	43	*	*	12	34	11	13	19	10
812.00	17	*	14	*	*	*5	*9	*	*	*9	*
812.01	20	*	17	*	*	*	15	*	*6	*6	*
812.09	9	*	*8	*	*	*	*6	*	*	*	*
812.2	14	*	10	*	*	*	*8	*	*	*7	*
812.20	*5	*	*	*	*	*	*	*	*	*	*
812.21	9	*	*6	*	*	*	*	*	*	*	*
812.4	25	11	14	13	*	*	*5	*7	*6	*	*8
812.41	14	*6	*8	9	*	*	*	*	*	*	*5
813	96	47	49	15	34	17	30	23	23	30	20
813.0	17	9	*7	*	*8	*	*5	*5	*	*	*
813.2	9	*6	*	*	*	*	*	*	*	*	*
813.23	*7	*	*	*	*	*	*	*	*	*	*
813.4	46	17	29	*5	13	*8	20	11	12	15	*8
813.41	11	*	*9	*	*	*	*6	*	*	*	*
813.42	22	*8	14	*	*8	*	9	*5	*6	*7	*
813.44	11	*6	*6	*	*	*	*	*	*	*	*
813.5	*7	*5	*	*	*	*	*	*	*	*	*
813.8	*8	*	*5	*	*	*	*	*	*	*	*
814	13	*8	*6	*	*	*	*5	*	*	*	*
814.0	13	*7	*6	*	*	*	*5	*	*	*	*
814.00	*5	*	*	*	*	*	*	*	*	*	*
814.01	*5	*	*	—	*	*	*	*	*	*	*
815	16	11	*5	*	10	*	*	*	*	*6	*
815.0	11	*8	*	*	*7	*	*	*	*	*	*
815.00	*6	*	*	*	*	*	*	*	*	*	*
815.1	*5	*	*	*	*	*	*	*	*	*	*
816	28	22	*6	*	15	*6	*	*9	*6	*9	*
816.0	14	10	*	*	*9	*	*	*	*	*	*
816.01	10	*7	*	*	*6	*	*	*	*	*	*
816.1	14	12	*	*	*6	*	*	*	*	*	*
816.11	*9	*7	*	*	*	*	*	*	*	*	*
820	390	92	298	*	12	26	348	74	102	140	74
820.0	124	29	95	*	*	*9	111	25	38	37	25
820.00	*5	*	*	—	—	*	*	*	*	*	*
820.02	*7	*	*	—	—	*	*6	*	*	*	*
820.03	10	*	*8	*	*	*	9	*	*	*	*
820.09	100	21	79	—	*	*6	91	22	30	31	17
820.2	180	44	136	*	*6	12	160	37	47	65	31
820.20	12	*	*9	*	*	*	11	*	*	*	*
820.21	148	33	115	*	*	9	134	28	39	55	26
820.22	19	*7	12	*	*	*	15	*5	*	*7	*
820.8	81	17	64	*	*	*6	74	12	17	36	18
821	66	34	32	12	20	*8	26	14	17	27	*9
821.0	39	23	17	10	12	*	13	*8	10	16	*6
821.00	10	*	*6	*	*	*	*6	*	*	*6	*
821.01	29	18	11	*8	11	*	*7	*6	*8	11	*
821.2	22	*7	15	*	*	*	12	*	*6	*9	*
821.21	*5	*	*	*	*	*	*	*	*	*	*
821.23	10	*	*8	*	*	*	*6	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
822	25	10	15	*	*9	*	11	*7	*	10	*
822.0	22	*8	13	*	*6	*	11	*7	*	*8	*
823	94	57	38	10	41	22	21	17	23	32	22
823.0	32	16	16	*	13	*8	9	*7	9	*8	*8
823.00	25	12	13	*	10	*7	*7	*6	*7	*7	*6
823.2	20	14	*6	*	11	*	*	*	*6	*8	*
823.20	*7	*	*	*	*	*	*	*	*	*	*
823.22	12	9	*	*	*6	*	*	*	*	*	*
823.3	*6	*6	*	*	*	*	*	*	*	*	*
823.8	27	14	13	*	9	*7	*8	*	*5	12	*6
823.80	*9	*	*6	*	*	*	*	*	*	*6	*
823.82	16	*9	*7	*	*6	*	*	*	*	*	*
823.9	*6	*	*	*	*	*	*	*	*	*	*
824	127	57	71	*	55	38	30	27	28	49	23
824.0	10	*7	*	*	*6	*	*	*	*	*	*
824.2	10	*5	*5	*	*	*	*	*	*	*	*
824.4	33	10	23	*	13	10	10	*8	*6	13	*6
824.6	31	10	21	*	12	11	*8	*7	*7	12	*5
824.8	32	18	14	*	15	*8	*7	*6	*6	13	*6
825	42	27	15	*	26	9	*6	12	*9	13	*8
825.0	14	*8	*6	*	*7	*	*	*	*	*	*
825.2	24	16	*8	*	17	*	*	*7	*	*7	*5
825.21	*6	*	*	*	*5	*	*	*	*	*	*
825.25	11	*8	*	*	*7	*	*	*	*	*	*
831	*7	*	*	*	*	*	*	*	*	*	*
831.0	*7	*	*	*	*	*	*	*	*	*	*
835	*6	*	*	*	*	*	*	*	*	*	*
835.0	*6	*	*	*	*	*	*	*	*	*	*
836	29	16	13	*	20	*6	*	*6	*8	11	*
836.0	12	*6	*6	*	*8	*	*	*	*	*6	*
836.1	*9	*	*	—	*6	*	*	*	*	*	*
839	*8	*6	*	*	*	*	*	*	*	*	*
840	53	29	24	—	11	22	20	10	17	16	10
840.4	44	24	20	—	*6	21	18	*8	13	15	*8
843	*6	*	*	*	*	*	*	*	*	*	*
844	41	22	20	*	26	*8	*6	12	9	12	*8
844.2	21	11	10	*	17	*	*	*	*	*8	*
844.8	*8	*	*	*	*	*	*	*	*	*	*
844.9	*8	*	*	*	*	*	*	*	*	*	*
845	20	11	*9	*	11	*	*	*5	*	*6	*
845.0	19	11	*8	*	11	*	*	*5	*	*5	*
845.00	10	*	*6	*	*	*	*	*	*	*	*
846	12	*6	*7	*	*5	*	*	*	*	*	*
846.0	10	*5	*5	—	*	*	*	*	*	*	*
847	39	21	18	*	17	11	11	*6	13	15	*
847.0	18	9	9	*	11	*	*	*	*6	*5	*
847.2	14	*8	*6	—	*	*	*6	*	*	*7	*
848	*8	*	*	—	*	*	*	*	*	*	*
850	59	35	24	*8	29	*7	14	24	9	15	11
850.0	20	12	*8	*	*8	*	*6	*6	*	*	*
850.1	20	10	10	*	11	*	*	10	*	*5	*
850.5	11	*8	*	*	*7	*	*	*6	*	*	*
850.9	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
851	19	14	*	*	9	*	*	*	*	*	*7
851.8	11	*8	*	*	*5	*	*	*	*	*	*
852	46	27	18	*	9	*8	26	*7	10	20	*9
852.0	*9	*5	*	*	*	*	*	*	*	*	*
852.2	34	21	14	*	*6	*6	21	*	*7	16	*7
852.20	16	10	*6	*	*	*	12	*	*	10	*
852.21	10	*	*6	*	*	*	*7	*	*	*	*
853	*9	*5	*	*	*	*	*	*	*	*	*
853.0	*9	*	*	*	*	*	*	*	*	*	*
854	72	45	27	11	33	10	18	10	20	20	22
854.0	71	45	27	11	32	10	18	10	20	19	22
854.00	23	14	9	*	*8	*	*8	*	*7	*6	*7
854.01	15	10	*5	*	*	*	*6	*	*5	*	*
854.02	18	12	*5	*	11	*	*	*	*	*	*8
854.06	11	*5	*5	*	*6	*	*	*	*	*	*
860	60	44	16	*	33	15	11	11	13	23	14
860.0	31	21	10	*	15	9	*6	*	*8	13	*6
860.2	*7	*	*	*	*	*	*	*	*	*	*
860.4	11	*7	*	*	*6	*	*	*	*	*	*
861	39	23	17	*5	21	*6	*7	*7	*7	15	10
861.0	*6	*	*	*	*	*	*	*	*	*	*
861.01	*6	*	*	*	*	*	*	*	*	*	*
861.2	31	17	14	*	16	*	*5	*	*	12	9
861.21	30	17	14	*	16	*	*5	*	*	12	9
862	*7	*6	*	*	*	*	*	*	*	*	*
863	23	19	*	*	16	*	*	*5	*	10	*
864	22	14	*8	*	13	*	*	*	*6	*9	*
864.0	17	9	*7	*	*8	*	*	*	*5	*6	*
864.01	*6	*	*	*	*	*	*	*	*	*	*
864.1	*5	*	*	*	*	*	—	*	*	*	*
865	20	11	9	*	12	*	*	*	*5	*8	*
865.0	18	9	*8	*	10	*	*	*	*	*7	*
866	14	10	*	*	*9	*	*	*	*	*6	*
866.0	12	*8	*	*	*7	*	*	*	*	*5	*
866.01	*7	*	*	*	*	*	*	*	*	*	*
867	13	*9	*	*	*	*	*7	*	*	*	*
867.0	11	*8	*	*	*	*	*6	*	*	*	*
868	12	*9	*	*	*6	*	*	*	*	*	*
868.0	9	*7	*	*	*	*	*	*	*	*	*
870	19	14	*5	*	*8	*6	*	*	*6	*5	*
870.8	11	*8	*	*	*	*	*	*	*	*	*
871	11	9	*	*	*	*	*	*	*	*	*
872	16	11	*6	*	*8	*	*	*	*	*	*6
872.0	11	*7	*	*	*	*	*	*	*	*	*
872.00	*5	*	*	*	*	*	*	*	*	*	*
873	202	129	73	21	99	29	53	40	49	66	47
873.0	52	33	19	*	24	*8	17	12	14	13	13
873.2	10	*7	*	*	*5	*	*	*	*	*	*
873.20	*8	*6	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
873.4	99	63	36	10	47	14	29	19	22	36	22
873.40	10	*6	*	*	*	*	*	*	*	*	*
873.42	50	30	19	*	21	*7	20	11	10	16	13
873.43	17	10	*7	*	*8	*	*	*	*	*7	*
873.44	10	*8	*	*	*6	*	*	*	*	*	*
873.49	9	*7	*	*	*5	*	*	*	*	*	*
873.5	*6	*	*	*	*	*	*	*	*	*	*
873.6	22	14	*7	*	14	*	*	*	*	*8	*6
873.63	12	*8	*	*	10	*	*	*	*	*	*
873.8	*5	*	*	*	*	*	*	*	*	*	*
874	*8	*7	*	*	*	*	*	*	*	*	*
875	*8	*6	*	*	*	*	*	*	*	*	*
875.0	*6	*5	*	*	*	*	*	*	*	*	*
879	17	13	*	*	13	*	*	*	*	*8	*
879.2	*6	*5	*	—	*6	*	*	*	*	*	*
880	11	10	*	*	*8	*	*	*	*	*	*
880.0	*9	*7	*	*	*6	*	*	*	*	*	*
881	41	24	17	*	22	*7	9	*7	15	11	*8
881.0	27	15	12	*	13	*5	*7	*	12	*7	*
881.00	12	*8	*	*	*7	*	*	*	*6	*	*
881.02	10	*	*6	*	*	*	*	*	*	*	*
881.1	*7	*	*	*	*	*	*	*	*	*	*
881.2	*6	*5	*	*	*	*	*	*	*	*	*
882	21	16	*5	*	13	*	*	*	*	*6	*6
882.0	11	9	*	*	*7	*	*	*	*	*	*
882.1	*7	*	*	*	*	*	*	*	*	*	*
883	36	25	10	*	18	*	11	*7	10	12	*7
883.0	15	10	*	*	*	*	*7	*	*	*	*
883.1	9	*7	*	*	*7	*	*	*	*	*	*
883.2	12	*8	*	*	*7	*	*	*	*	*5	*
884	10	*6	*	*	*5	*	*	*	*	*	*
884.0	*8	*6	*	*	*	*	*	*	*	*	*
886	12	11	*	*	*	*5	*	*	*	*	*
886.0	11	11	*	*	*	*	*	*	*	*	*
890	11	*9	*	*	*9	*	*	*	*	*6	*
890.0	*6	*5	*	*	*	*	*	*	*	*	*
891	38	25	12	*	21	*7	*7	*8	*7	13	9
891.0	20	12	*8	*	12	*	*	*	*	*6	*6
891.1	15	11	*	*	*8	*	*	*	*	*7	*
892	10	*8	*	*	*	*	*	*	*	*	*
892.1	*6	*	*	*	*	*	*	*	*	*	*
903	*7	*6	*	*	*5	*	—	*	*	*	*
904	*5	*	*	*	*	*	*	*	*	*	*
905	58	26	32	*	24	11	21	13	21	10	14
905.2	14	*6	*8	*	*6	*	*6	*	*6	*	*
905.3	15	*6	9	—	*	*	10	*	*6	*	*
905.4	14	*6	*8	*	*6	*	*	*	*	*5	*
906	16	*8	*8	*	10	*	*	*	*6	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
907	37	30	*7	*	20	9	*7	*6	10	13	*9
907.0	16	13	*	*	*7	*	*	*	*	*6	*
907.2	20	17	*	—	12	*	*	*	*6	*6	*
908	10	*7	*	*	*	*	*	*	*	*	*
908.9	*6	*	*	—	*	*	*	*	*	*	*
909	*6	*	*	*	*	*	*	*	*	*	*
910	24	16	*9	*	10	*	*6	*	*8	*6	*6
910.0	22	15	*7	*	10	*	*5	*	*7	*	*6
911	*9	*6	*	*	*5	*	*	*	*	*	*
911.0	*7	*5	*	*	*	*	*	*	*	*	*
912	*5	*	*	*	*	*	*	*	*	*	*
913	15	9	*5	*	*6	*	*5	*	*	*	*5
913.0	13	*8	*	*	*5	*	*5	*	*	*	*
914	*5	*	*	*	*	*	*	*	*	*	*
916	25	17	*8	*	9	*	*8	*	*6	*8	*7
916.0	19	13	*6	*	*8	*	*6	*	*	*6	*6
918	*6	*	*	*	*	*	*	*	*	*	*
919	19	11	*8	*	10	*	*	*	*6	*6	*
919.0	17	10	*7	*	9	*	*	*	*	*5	*
920	46	20	26	*9	13	*	22	*9	13	14	10
921	16	*9	*7	*	*7	*	*	*	*	*6	*
921.1	*6	*	*	*	*	*	*	*	*	*	*
922	54	28	27	*	23	10	19	15	16	15	*8
922.1	27	14	13	*	12	*6	*9	*9	*6	*8	*
922.2	14	*7	*7	*	*7	*	*	*	*5	*	*
922.3	*8	*	*	*	*	*	*	*	*	*	*
923	28	14	14	*	*8	*	15	*6	11	*7	*5
923.0	13	*6	*7	*	*5	*	*6	*	*	*	*
923.00	11	*6	*6	*	*	*	*	*	*	*	*
923.1	*6	*	*	*	*	*	*	*	*	*	*
924	85	34	52	*	27	13	41	18	23	30	14
924.0	22	9	13	*	*	*	16	*6	*7	*6	*
924.01	17	*8	10	*	*	*	14	*	*6	*	*
924.1	18	*7	11	*	*5	*	10	*	*6	*	*
924.10	*5	*	*	*	*	*	*	*	*	*	*
924.11	13	*	*9	*	*	*	*8	*	*	*	*
924.5	*6	*	*	*	*	*	*	*	*	*	*
924.8	32	14	18	*	15	*	11	*5	*6	15	*7
933	*7	*	*	*	*	*	*	*	*	*	*
933.1	*6	*	*	*	*	*	*	*	*	*	*
934	14	*7	*7	*	*	*	*7	*	*	*	*
934.8	*7	*	*	*	*	*	*	*	*	*	*
935	16	*8	*9	*6	*	*	*	*6	*	*6	*
935.1	12	*6	*6	*6	*	*	*	*	*	*	*
941	11	10	*	*	*	*	*	*	*	*	*
941.2	*8	*7	*	*	*	*	*	*	*	*	*
942	13	*8	*5	*	*	*	*	*	*	*	*
942.2	*7	*	*	*	*	*	*	*	*	*	*
943	12	*6	*5	*	*6	*	*	*	*	*	*
943.2	*7	*	*	*	*	*	*	*	*	*	*
944	12	*9	*	*	*7	*	*	*	*	*	*
944.2	*5	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
945	16	9	*7	*5	*	*	*	*	*5	*	*
945.2	10	*5	*	*	*	*	*	*	*	*	*
946	13	10	*	*	*5	*	*	*	*	*8	*
946.2	*9	*7	*	*	*	*	*	*	*	*6	*
948	27	20	*7	*6	11	*7	*	*8	*6	*9	*
948.0	17	12	*	*5	*6	*	*	*6	*	*	*
948.00	17	12	*	*5	*6	*	*	*6	*	*	*
948.1	*5	*	*	*	*	*	*	*	*	*	*
952	*9	*7	*	*	*	*	*	*	*	*	*
955	15	13	*	*	12	*	*	*	*	*	*
955.2	*5	*	*	*	*	*	*	*	*	*	*
958	39	29	10	*	18	*8	11	*7	*9	12	11
958.3	10	*6	*	*	*	*	*	*	*	*	*
958.4	*6	*	*	*	*	*	*	*	*	*	*
958.7	*8	*6	*	*	*	*	*	*	*	*	*
958.8	13	11	*	*	*9	*	*	*	*	*	*
959	39	18	21	*	19	*	12	*8	9	13	*8
959.1	17	*8	*8	*	10	*	*	*	*	*7	*
959.8	*6	*	*	*	*	*	*	*	*	*	*
962	*6	*	*	*	*	*	*	*	*	*	*
963	11	*	*8	*	*8	*	*	*	*	*	*
963.0	9	*	*6	*	*7	*	*	*	*	*	*
965	76	24	52	*7	56	*8	*	16	17	26	17
965.0	21	10	11	*	15	*	*	*	*	*7	*5
965.00	*5	*	*	*	*	*	*	*	*	*	*
965.09	10	*	*7	*	*7	*	*	*	*	*	*
965.1	13	*	9	*	*8	*	*	*	*	*5	*
965.4	29	*7	22	*	23	*	*	*7	*6	10	*6
965.6	*7	*	*6	*	*6	*	*	*	*	*	*
966	16	*9	*7	*	*8	*	*	*	*	*6	*
966.1	*8	*	*	*	*	*	*	*	*	*	*
966.3	*7	*	*	*	*	*	—	*	*	*	*
967	15	*6	9	*	*8	*	*	*	*	*	*5
967.0	*8	*	*	*	*5	*	*	*	*	*	*
968	13	*8	*6	*	12	*	*	*	*	*	*
968.5	*8	*6	*	*	*7	*	*	*	*	*	*
969	96	37	59	*5	68	15	*8	19	28	31	17
969.0	31	11	20	*	21	*5	*	*7	*9	10	*
969.1	*6	*	*	*	*	*	*	*	*	*	*
969.4	41	18	23	*	31	*5	*	*7	14	12	*7
969.7	*7	*	*	*	*	*	—	*	*	*	*
971	*5	*	*	*	*	*	*	*	*	*	*
972	17	*7	11	*	*	*	10	*	*7	*	*
972.1	10	*	*7	*	*	*	9	*	*	*	*
977	18	9	9	*	14	*	*	*	*5	10	*
977.8	*8	*	*	*	*6	*	*	*	*	*	*
977.9	9	*	*5	*	*7	*	*	*	*	*5	*
980	19	10	*9	*	15	*	*	*	*5	*6	*
980.0	16	*9	*7	*	13	*	*	*	*	*5	*
986	*5	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
987	9	*7	*	*	*	*	*	*	*	*	*
987.9	*5	*	*	*	*	*	*	*	*	*	*
989	11	*8	*	*	*	*	*	*	*	*6	*
989.5	*9	*7	*	*	*	*	*	*	*	*5	*
990	*5	*	*	—	*	*	*	*	*	*	*
991	13	*8	*6	*	*5	*	*6	*	*	*5	*
991.6	*7	*	*	*	*	*	*	*	*	*	*
994	10	*7	*	*	*	*	*	*	*	*	*
994.1	*5	*	*	*	*	*	*	*	*	*	*
995	72	30	42	*7	13	16	35	19	21	24	*8
995.1	*8	*	*	*	*	*	*	*	*	*	*
995.2	43	16	27	*	*7	*6	29	13	13	14	*
996	613	315	297	44	126	162	280	144	112	228	128
996.0	42	25	18	*	*	*8	31	11	9	13	10
996.01	29	15	14	*	*	*	23	*8	*5	9	*6
996.02	*5	*	*	*	*	*	*	*	*	*	*
996.04	*5	*	*	—	*	*	*	*	*	*	*
996.1	26	11	14	*	*	*8	10	*6	*5	12	*
996.2	23	13	10	12	*	*	*	*6	*6	*7	*
996.3	9	*7	*	*	*	*	*	*	*	*	*
996.39	*7	*5	*	*	*	*	*	*	*	*	*
996.4	81	37	44	*	14	21	45	18	18	30	15
996.5	32	12	20	*	*6	*8	14	*7	*6	11	*7
996.59	25	11	15	*	*	*6	13	*6	*	*8	*6
996.6	175	92	83	14	39	42	80	40	35	64	37
996.61	*6	*	*	*	*	*	*	*	*	*	*
996.62	85	45	40	*9	23	22	31	24	15	28	18
996.64	18	*8	10	*	*	*	14	*	*	*6	*
996.66	15	*8	*8	*	*	*	9	*	*	*6	*
996.67	13	*6	*7	*	*	*	*5	*	*	*7	*
996.69	28	16	12	*	*5	*7	12	*	*	11	*9
996.7	187	97	90	*5	38	56	88	42	29	71	46
996.72	*7	*	*	—	*	*	*5	*	*	*	*
996.73	71	36	35	*	19	23	29	14	10	27	20
996.74	55	28	28	*	*7	15	30	13	*8	22	12
996.76	14	10	*	*	*	*	*8	*	*	*5	*
996.77	11	*5	*5	—	*	*	*6	*	*	*	*
996.78	12	*6	*6	*	*5	*	*	*	*	*	*
996.79	11	*	*6	*	*	*	*	*	*	*	*
996.8	37	21	15	*	17	13	*	12	*	17	*5
996.81	23	14	9	*	11	9	*	*6	*	11	*
997	686	353	333	26	114	174	372	158	154	217	158
997.0	31	13	18	*	*	*7	18	*8	*	*8	9
997.02	18	*8	9	*	*	*	13	*6	*	*	*6
997.09	9	*	*6	*	*	*	*	*	*	*	*
997.1	147	85	62	*	*7	38	98	47	30	41	29
997.2	31	17	14	*	*	9	18	*6	10	10	*6
997.3	154	77	77	*8	23	39	83	31	36	46	41
997.4	203	93	110	10	55	49	89	40	45	71	48
997.5	85	46	39	*	15	20	49	18	20	26	20
997.6	23	15	*8	*	*	*8	13	*5	*	10	*
997.62	16	10	*6	*	*	*	9	*	*	*6	*
997.69	*5	*	*	—	*	*	*	*	*	*	*
997.9	12	*6	*5	*	*	*	*	*	*	*	*
997.91	*5	*	*	*	*	*	*	*	*	*	*
997.99	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
998	631	290	341	24	154	176	276	146	138	225	122
998.1	195	92	102	*5	37	51	101	50	46	63	36
998.2	77	31	46	*	26	23	27	16	16	30	16
998.3	45	25	19	*	9	11	21	*8	12	16	*8
998.5	170	79	92	*6	46	46	72	41	39	59	31
998.6	13	*5	*8	*	*	*	*5	*	*	*	*
998.8	122	52	69	*7	31	37	46	24	21	48	28
998.81	*6	*	*	*	*	*	*	*	*	*	*
998.89	115	49	66	*7	30	36	42	22	20	46	26
999	54	25	28	*	12	11	27	13	13	16	12
999.2	24	11	12	*	*	*	17	*6	*	*9	*
999.3	9	*	*	*	*	*	*	*	*	*	*
999.8	10	*	*6	*	*	*	*5	*	*	*	*
999.9	*6	*	*	*	*	*	*	*	*	*	*
V01–V82	8,372	1,910	6,462	232	4,949	990	2,202	1,619	2,143	2,686	1,924
V02	32	*	30	*	27	*	*	*7	*9	*5	11
V02.5	25	*	24	—	22	*	*	*5	*7	*	9
V02.6	*7	*	*6	*	*	*	*	*	*	*	*
V04	9	—	9	—	9	—	—	*	*	*	*
V04.3	9	—	9	—	9	—	—	*	*	*	*
V05	*5	*	*	*	*	—	—	*	*	*	*
V07	24	*	22	*	18	*	*	10	*	*	*7
V07.2	17	*	16	*	16	—	—	*8	*	*	*
V08	37	27	10	*	27	*8	*	20	*	11	*
V09	22	10	13	*	*6	*	13	*5	*8	*7	*
V09.0	18	*8	10	*	*	*	11	*	*6	*6	*
V10	873	349	523	*	53	217	599	209	254	239	172
V10.0	193	95	98	—	*6	42	145	48	59	52	34
V10.02	11	*8	*	—	*	*5	*6	*	*	*	*
V10.04	*8	*	*	—	*	*	*6	*	*	*	*
V10.05	141	65	76	—	*	27	110	33	48	36	24
V10.06	17	10	*7	—	*	*	13	*6	*	*5	*
V10.1	53	28	25	*	*	15	37	14	12	20	*8
V10.11	52	27	25	*	*	15	36	13	12	20	*8
V10.2	19	14	*5	—	*	*	14	*	*5	*7	*
V10.21	17	13	*	—	*	*	12	*	*	*7	*
V10.3	250	*	246	—	15	74	162	59	59	75	57
V10.4	194	111	83	—	13	37	143	45	67	41	40
V10.41	21	—	21	—	*6	*6	*9	*	*8	*	*
V10.42	31	—	31	—	*	*7	23	*7	11	*6	*7
V10.43	22	—	22	—	*	10	9	*6	*8	*5	*
V10.44	*9	—	*9	—	*	*	*5	*	*	*	*
V10.46	106	106	—	—	*	9	97	23	38	24	21
V10.5	63	44	19	*	*	13	48	17	19	17	10
V10.51	36	27	*9	—	*	*5	31	11	12	9	*
V10.52	24	15	9	*	*	*7	16	*6	*7	*7	*
V10.7	22	*8	15	*	*	*7	10	*5	*8	*5	*
V10.72	*6	*	*	—	*	*	*	*	*	*	*
V10.79	16	*6	10	*	*	*6	*8	*	*7	*	*
V10.8	72	42	30	*	10	23	36	16	22	19	14
V10.82	19	13	*6	—	*	*8	9	*	*7	*5	*
V10.83	24	16	*8	—	*	*6	17	*	10	*6	*
V10.85	*8	*	*	*	*	*	*	*	*	*	*
V10.87	9	*	*7	*	*	*	*	*	*	*	*
V10.89	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
V11	39	18	21	*	13	10	16	*	15	10	10
V11.3	20	11	*8	—	*6	*6	*8	*	*8	*5	*
V11.8	12	*	*9	*	*	*	*6	*	*	*	*
V12	503	228	275	*9	59	135	300	90	174	147	92
V12.0	36	16	19	*	10	9	16	*6	9	12	*9
V12.01	13	*6	*7	*	*	*	*9	*	*	*	*
V12.02	*8	*	*	*	*	*	*	*	*	*	*
V12.09	13	*5	*8	*	*6	*	*	*	*	*5	*
V12.2	17	*6	11	*	*	*7	*7	*	*5	*5	*
V12.4	14	*6	*8	*	*7	*	*5	*	*6	*	*
V12.5	280	123	157	*	19	68	192	51	107	74	47
V12.50	27	11	16	*	*	*6	19	*5	11	10	*
V12.51	46	19	26	—	*7	11	28	10	19	*9	*8
V12.52	*7	*	*	—	*	*	*	*	*	*	*
V12.59	200	90	111	*	*8	49	141	35	74	55	37
V12.6	16	*8	*8	*	*	*6	*6	*	*	*6	*
V12.7	137	67	70	*	19	43	72	22	42	44	28
V12.71	74	38	36	*	*9	24	41	13	18	27	15
V12.72	12	*8	*	—	*	*	10	*	*	*	*
V12.79	47	19	28	*	9	16	19	*6	19	13	10
V13	65	34	31	13	13	10	28	11	28	16	9
V13.0	28	17	11	*	*5	*5	17	*5	12	*8	*
V13.01	15	11	*	—	*	*	*7	*	*6	*	*
V13.09	12	*6	*6	*	*	*	9	*	*6	*	*
V13.5	10	*	*7	*	*	*	*7	*	*	*	*
V13.7	10	*7	*	10	*	—	—	—	*8	*	*
V14	20	*7	13	*	*	*	11	*	*8	*	*
V14.0	11	*	*7	*	*	*	*7	*	*	*	*
V15	352	196	155	20	87	112	133	56	97	94	104
V15.1	18	11	*8	13	*	*	*	10	*	*	*
V15.3	18	10	*8	*	*	*6	*8	*	*	*	*7
V15.5	17	*8	9	*	*7	*	*6	*	*7	*5	*
V15.8	292	165	127	*5	71	101	114	38	83	80	91
V15.81	104	55	48	*	48	31	22	17	29	38	20
V15.82	166	101	65	*	15	66	85	19	48	36	63
V15.89	20	*7	13	*	*8	*	*	*	*5	*5	*7
V16	12	*	10	—	*	*	*	*	*	*	*
V17	39	23	16	*	12	18	*9	*6	16	*6	11
V17.3	28	15	12	—	*7	14	*6	*	11	*	10
V17.4	*9	*7	*	—	*	*	*	*	*	*	*
V18	*7	*	*	*	*	*	*	*	*	*	*
V22	37	—	37	*	37	—	—	11	10	*7	9
V22.1	*8	—	*8	—	*8	—	—	*	*	*	*
V22.2	29	—	29	*	28	—	—	*6	9	*7	*6
V23	86	—	86	*	86	*	—	*8	17	43	18
V23.4	*7	—	*7	—	*7	—	—	*	*	*	*
V23.7	64	—	64	*	64	—	—	*	11	35	14
V23.8	*8	—	*8	*	*8	*	—	*	*	*	*
V24	*7	—	*7	—	*7	—	—	*	*	*	*
V24.0	*6	—	*6	—	*6	—	—	*	*	*	*
V25	323	*	322	—	321	*	—	40	47	161	75
V25.2	317	*	316	—	315	*	—	40	47	155	75

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
V27	3,831	—	3,831	10	3,818	*	—	624	785	1,385	1,036
V27.0	3,184	—	3,184	*8	3,173	*	—	539	617	1,145	882
V27.1	18	—	18	*	18	—	—	*	*	*8	*
V27.2	41	—	41	*	40	*	—	9	*6	14	11
V27.9	585	—	585	*	584	*	—	72	158	216	138
V29	22	13	9	22	—	—	—	*	9	*	*6
V29.0	21	12	*9	21	—	—	—	*	*8	*	*6
V30	*8	*	*	*8	—	—	—	*	*	*	*
V30.1	*8	*	*	*8	—	—	—	*	*	*	*
V42	49	26	23	*9	15	17	*8	21	12	*8	*7
V42.0	23	12	11	*	*8	11	*	*6	*8	*	*
V42.1	*6	*	*	*	*	*	*	*	*	*	*
V42.7	*7	*	*	*	*	*	*	*	*	*	*
V42.8	*6	*	*	*	*	*	—	*	*	*	*
V43	132	49	84	*	*5	22	104	34	58	24	16
V43.3	43	21	21	*	*	9	31	13	15	9	*6
V43.6	85	25	60	*	*	13	70	20	41	14	10
V43.64	41	11	30	*	*	*6	33	11	19	*6	*5
V43.65	42	14	29	*	*	*6	36	*8	22	*8	*
V44	75	39	35	18	11	15	32	24	31	11	*9
V44.0	16	10	*6	*6	*	*	*5	*	*7	*	*
V44.1	22	11	11	*8	*	*	*9	*8	*8	*	*
V44.2	10	*	*7	*	*	*	*	*	*5	*	*
V44.3	19	10	*9	*	*	*	12	*6	*8	*	*
V45	600	363	237	20	42	148	389	151	257	111	82
V45.0	100	51	49	*	*	11	85	31	36	21	12
V45.01	89	42	46	*	*	*8	78	28	31	19	11
V45.02	9	*8	*	—	*	*	*6	*	*	*	*
V45.1	32	18	14	*	*7	13	12	*6	*	10	11
V45.2	14	*7	*6	9	*	*	*	*5	*	*	*
V45.4	*9	*	*	*	*	*	*	*	*	*	*
V45.8	441	282	159	10	27	118	285	107	205	76	53
V45.81	260	178	82	*	*	65	190	66	119	46	29
V45.82	107	71	36	*	*7	38	62	29	45	16	16
V45.89	74	32	41	9	16	15	33	12	40	13	*8
V46	12	*6	*5	*	*	*	*	*	*	*	*
V46.1	11	*6	*5	*	*	*	*	*	*	*	*
V49	56	40	16	*	*9	19	27	11	18	16	11
V49.6	*6	*	*	*	*	*	*	*	*	*	*
V49.7	49	33	15	*	*7	17	25	*9	16	15	*9
V49.72	*6	*	*	—	*	*	*	*	*	*	*
V49.75	21	15	*5	*	*	*7	10	*	*7	*6	*
V49.76	13	*7	*6	*	*	*	*8	*	*	*5	*
V51	*7	*	*	*	*	*	*	*	*	*	*
V53	*7	*	*	*	*	*	*	*	*	*	*
V53.3	*6	*	*	*	*	*	*	*	*	*	*
V54	28	13	15	*	*7	*7	13	*	*7	10	*7
V54.0	*8	*5	*	*	*	*	*	*	*	*	*
V54.8	11	*	*7	*	*	*	*7	*	*	*	*
V54.9	*9	*	*5	*	*	*	*	*	*	*6	*
V55	45	22	23	*8	9	*8	19	16	9	14	*5
V55.1	*6	*	*	*	*	*	*	*	*	*	*
V55.3	30	13	17	*	*8	*6	15	*7	*7	12	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
V57	374	147	227	*	37	57	278	45	122	138	69
V57.1	123	44	78	*	*9	18	95	16	28	60	18
V57.2	31	11	21	*	*	*6	24	*	*9	16	*
V57.21	31	11	21	*	*	*6	24	*	*9	16	*
V57.8	207	86	120	*	25	31	149	26	83	53	44
V57.89	206	86	120	*	25	31	149	26	83	53	44
V57.9	9	*	*7	*	*	*	*7	*	*	*7	*
V58	349	154	195	32	62	110	145	114	66	102	67
V58.0	45	19	27	*	*7	11	26	12	*8	17	*9
V58.1	232	107	125	29	43	83	77	87	42	62	41
V58.4	17	*7	10	*	*	*	*8	*5	*	*	*
V58.49	16	*6	10	*	*	*	*8	*	*	*	*
V58.6	34	16	18	*	*6	*9	20	*8	*8	10	*9
V58.61	31	14	16	*	*	*8	18	*8	*7	*9	*8
V58.8	18	*6	13	*	*	*	13	*	*	9	*
V58.89	17	*	12	*	*	*	13	*	*	*8	*
V59	*8	*	*	*	*	*	*	*	*	*	*
V60	21	10	11	*	13	*	*	*7	*6	*	*
V60.0	9	*5	*	—	*7	*	*	*	*	*	*
V60.2	*6	*	*	*	*5	*	*	*	*	*	*
V61	57	18	39	10	37	*7	*	11	11	25	*9
V61.0	*6	*	*5	*	*	*	*	*	*	*	*
V61.1	17	*7	9	—	12	*	*	*	*	9	*
V61.2	12	*6	*6	*6	*6	*	—	*	*	*	*
V61.20	*7	*	*	*	*	*	—	*	*	*	*
V61.5	*9	—	*9	—	*8	—	*	*	*	*6	*
V61.8	*7	*	*	*	*	*	—	*	*	*	*
V62	35	19	15	*	26	*	*	10	*9	10	*6
V62.5	*8	*6	*	*	*7	*	*	*	*	*6	*
V62.8	18	*9	10	*	13	*	*	*6	*7	*	*
V62.81	*6	*	*	*	*5	*	*	*	*	*	*
V62.89	11	*6	*6	*	*8	*	*	*	*	*	*
V63	*5	*	*	*	*	*	*	*	*	*	*
V64	42	16	26	*	13	9	18	9	10	16	*6
V64.1	19	*8	11	*	*5	*	*8	*	*	*7	*
V64.2	14	*6	*8	*	*5	*	*6	*	*	*5	*
V64.3	*9	*	*6	*	*	*	*	*	*	*	*
V65	26	12	14	*	15	*	*	*9	*6	*5	*6
V65.3	12	*7	*5	*	*	*	*	*	*	*	*5
V65.4	10	*	*7	—	*8	*	*	*7	*	*	*
V65.44	*7	*	*6	—	*5	*	*	*7	—	—	—
V66	16	*6	10	*	*	*	12	*	*	*	10
V66.5	*6	*	*	—	—	*	*6	*	*	*	*5
V67	10	*	*6	*	*	*	*	*	*	*	*
V67.0	*7	*	*5	*	*	*	*	*	*	*	*
V70	12	*7	*	*	*8	*	*	*	*	*	*
V70.7	11	*7	*	*	*7	*	*	*	*	*	*
V71	26	14	11	9	*8	*	*	*7	*9	*	*7
V71.4	*8	*5	*	*	*	*	*	*	*	*	*
V71.8	11	*5	*6	*8	*	*	*	*	*7	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
All codes	40,397	15,997	24,400	1,950	14,603	9,302	14,542	9,870	8,771	12,973	8,783
01–05	937	431	506	199	317	196	225	222	217	290	208
01	120	67	53	16	32	35	37	27	25	42	27
01.0	*9	*6	*	*	*	*	*	*	*	*	*
01.02	*6	*	*	*	*	*	*	*	*	*	*
01.1	19	13	*6	*	*8	*6	*	*	*	*6	*
01.18	10	*7	*	*	*	*	*	*	*	*	*
01.2	22	12	10	*	*6	*5	*8	*5	*	*6	*6
01.24	15	*9	*6	*	*	*	*5	*	*	*	*
01.3	32	17	15	*	*7	*8	14	*	*6	15	*6
01.31	19	10	9	*	*	*	11	*	*	*8	*
01.39	12	*6	*6	*	*	*5	*	*	*	*6	*
01.5	36	18	18	*	10	13	*9	9	*9	10	*8
01.51	*5	*	*	*	*	*	*	*	*	*	*
01.59	28	15	13	*	*8	11	*6	*7	*8	*8	*6
02	101	53	48	29	35	18	20	27	26	28	21
02.0	13	*9	*	*	*6	*	*	*	*	*	*
02.06	*5	*	*	*	*	*	*	*	*	*	*
02.1	*7	*	*	*	*	*	*	*	*	*	*
02.12	*6	*	*	*	*	*	*	*	*	*	*
02.2	20	9	11	*	*7	*5	*6	*6	*	*7	*
02.3	21	11	10	*6	*6	*	*6	*	*	*5	*7
02.34	17	*9	*8	*	*	*	*	*	*	*	*6
02.4	26	15	11	15	*	*	*	*7	10	*6	*
02.42	22	13	*9	13	*	*	*	*6	*7	*	*
02.9	14	*7	*7	*	*9	*	*	*	*	*	*
02.94	9	*	*	*	*6	*	*	*	*	*	*
03	653	281	373	150	224	126	152	153	153	199	148
03.0	111	59	52	*	19	42	49	26	27	37	22
03.09	108	57	51	*	17	42	48	25	26	35	21
03.3	313	158	155	137	82	40	54	66	74	93	79
03.31	310	157	153	137	80	39	54	65	74	92	79
03.4	10	*	*	*	*	*	*	*	*	*	*
03.5	18	10	*8	*	*6	*6	*	*	*	*7	*
03.59	12	*6	*6	*	*	*	*	*	*	*5	*
03.6	*8	*	*	*	*	*	*	*	*	*5	*
03.8	*5	*	*	*	*	*	*	*	*	*	*
03.9	185	42	143	*	110	31	40	49	43	52	41
03.90	85	21	64	*	50	16	19	26	24	26	*9
03.91	51	*5	46	*	39	*7	*5	12	*9	9	22
03.92	29	11	18	*	*8	*6	13	*7	*6	*8	*8
03.95	11	*	10	—	10	*	*	*	*	*6	*
04	51	26	25	*	20	16	12	14	12	16	9
04.0	9	*	*	*	*	*	*	*	*	*	*
04.07	*5	*	*	*	*	*	*	*	*	*	*
04.3	*5	*	*	*	*	*	*	*	*	*	*
04.4	18	*6	11	*	*7	*7	*	*6	*	*6	*
04.43	*8	*	*	*	*	*	*	*	*	*	*
04.49	*7	*	*	*	*	*	*	*	*	*	*
04.8	11	*5	*6	—	*	*	*	*	*	*	*
04.81	11	*5	*5	—	*	*	*	*	*	*	*
05	11	*	*7	—	*6	*	*	*	*	*	*
05.3	*7	*	*5	—	*	*	*	*	*	*	*
05.31	*6	*	*	—	*	*	*	*	*	*	*
06–07	101	29	71	*	33	37	28	31	17	31	21

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
06	86	22	64	*	29	32	23	27	14	26	18
06.1	*7	*	*	*	*	*	*	*	*	*	*
06.2	21	*	18	*	10	*8	*	*6	*	*5	*
06.3	15	*	11	*	*7	*	*	*	*	*6	*
06.39	12	*	9	*	*6	*	*	*	*	*5	*
06.4	10	*	*9	*	*	*	*	*	*	*	*
06.8	20	*5	14	*	*	*8	*8	*6	*	*	*6
06.89	17	*	13	—	*	*7	*7	*	*	*	*
07	15	*8	*7	*	*	*	*	*	*	*5	*
07.6	*6	*	*	*	*	*	*	*	*	*	*
08–16	188	94	95	10	35	44	99	52	19	84	33
08	39	23	16	*	14	11	12	12	*6	13	*8
08.8	25	16	*9	*	10	*6	*7	*7	*	*8	*5
08.81	22	15	*6	*	10	*	*6	*7	*	*6	*
11	*5	*	*	*	*	*	*	*	*	*	*
12	15	*6	10	*	*	*6	*6	*5	*	*9	*
12.9	*7	*	*5	*	*	*	*	*	*	*	—
13	51	21	29	*	*	*8	40	15	*	17	16
13.4	18	*7	11	*	*	*	15	*	*	*7	*6
13.41	17	*6	11	*	*	*	14	*	*	*7	*6
13.5	*6	*	*	—	*	*	*	*	*	*	*
13.59	*6	*	*	—	*	*	*	*	*	*	*
13.7	23	9	14	*	*	*	19	*7	*	*8	*8
13.71	23	9	14	*	*	*	19	*7	*	*8	*8
14	56	27	30	*	9	13	33	13	*6	34	*
14.2	*7	*	*5	*	*	*	*5	*	*	*6	*
14.24	*6	*	*	*	*	*	*	*	*	*	*
14.4	*8	*5	*	*	*	*	*	*	*	*	*
14.49	*6	*	*	*	*	*	*	*	*	*	*
14.5	9	*	*6	*	*	*	*6	*	*	*7	*
14.7	22	11	11	*	*	*	13	*7	*	12	*
14.74	14	*6	*8	*	*	*	*8	*	*	*8	*
14.9	*5	*	*	*	*	*	*	*	*	*	*
16	13	*8	*5	*	*	*	*	*	*	*7	*
18–20	58	35	23	35	13	*	*6	21	*8	12	17
18	18	12	*6	*5	*7	*	*	*	*	*	*9
18.4	*7	*	*	*	*	*	*	*	*	*	*
19	*7	*	*	*	*	*	*	*	*	*	*
20	34	20	14	27	*	*	*	14	*5	*8	*7
20.0	25	15	10	23	*	*	*	11	*	*6	*
20.01	23	14	9	22	*	*	*	10	*	*5	*
21–29	323	189	135	69	121	72	61	106	52	100	65
21	87	56	31	*5	37	21	24	27	17	27	16
21.0	28	16	13	*	*5	*7	15	10	*7	*6	*
21.01	10	*6	*	*	*	*	*6	*	*	*	*
21.02	*8	*	*	*	*	*	*	*	*	*	*
21.03	*8	*	*	*	*	*	*	*	*	*	*
21.3	*6	*	*	*	*	*	*	*	*	*	*
21.5	*7	*	*	*	*	*	*	*	*	*	*
21.6	*9	*6	*	*	*6	*	*	*	*	*	*
21.69	*6	*	*	*	*	*	*	*	*	*	*
21.7	*9	*7	*	*	*5	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
21.71	*6	*5	*	*	*	*	*	*	*	*	*
21.8	22	15	*7	*	11	*7	*	*6	*	*9	*
21.81	*5	*	*	*	*	*	*	*	*	*	*
21.88	9	*6	*	*	*	*	*	*	*	*	*
22	36	18	18	*	13	13	*6	13	*6	10	*8
22.1	*5	*	*	*	*	*	*	*	*	*	*
22.2	*7	*	*	*	*	*	*	*	*	*	*
22.6	15	*8	*8	*	*6	*5	*	*5	*	*	*
22.63	11	*6	*	*	*	*	*	*	*	*	*
23	37	22	14	*7	21	*7	*	12	*	10	11
23.0	13	*8	*5	*	*8	*	*	*	*	*	*
23.09	13	*8	*5	*	*8	*	*	*	*	*	*
23.1	16	10	*6	*	9	*	*	*6	*	*	*
23.19	16	10	*6	*	9	*	*	*6	*	*	*
24	*8	*6	*	*	*	*	*	*	*	*	*
25	12	*8	*	*	*	*	*	*	*	*	*
26	18	*8	10	*	*	*5	*8	*6	*	*7	*
26.3	14	*7	*7	*	*	*	*6	*	*	*6	*
26.31	*6	*	*	*	*	*	*	*	*	*	*
26.32	*6	*	*	*	*	*	*	*	*	*	*
27	48	25	23	17	17	*6	*7	10	9	17	12
27.0	*6	*	*	*	*	*	*	*	*	*	*
27.5	21	11	10	*6	*9	*	*	*5	*5	*5	*5
27.51	11	*6	*5	*	*7	*	*	*	*	*	*
27.6	10	*6	*	*9	*	*	*	*	*	*	*
27.62	*5	*	*	*5	*	—	—	*	*	*	*
28	49	25	24	30	14	*	*	24	*	14	*6
28.0	*6	*	*	*	*	*	*	*	*	*	*
28.2	13	*6	*7	*	*7	*	*	*6	*	*	*
28.3	23	13	10	20	*	—	*	13	*	*7	*
29	28	20	*8	*	*8	*8	9	*9	*	10	*
29.1	9	*6	*	*	*	*	*	*	*	*	*
29.11	*7	*	*	*	*	*	*	*	*	*	*
29.4	12	10	*	*	*	*	*	*	*	*	*
30–34	1,038	582	456	57	172	298	511	270	217	343	208
30	12	9	*	*	*	*	*5	*	*	*	*
30.0	*5	*	*	*	*	*	*	*	*	*	*
30.09	*5	*	*	*	*	*	*	*	*	*	*
31	143	88	55	21	24	42	57	46	25	40	31
31.1	63	40	23	*	12	20	29	18	13	16	16
31.2	17	10	*8	*	*	*	*8	*	*	*6	*
31.29	17	10	*7	*	*	*	*8	*	*	*6	*
31.4	53	33	21	14	*6	15	17	21	*7	16	10
31.42	44	26	18	14	*6	11	13	19	*7	12	*7
31.43	*8	*6	*	*	*	*	*	*	*	*	*
32	69	35	35	*	*6	23	39	17	15	22	16
32.2	24	13	11	*	*	*7	11	*7	*6	*6	*
32.29	18	*9	9	*	*	*6	9	*5	*5	*	*
32.3	*7	*	*	*	*	*	*	*	*	*	*
32.4	30	12	17	*	*	*8	21	*7	*7	*8	*8
32.5	*6	*	*	—	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
— Quantity zero.NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
33	357	208	149	21	48	108	181	94	72	118	72
33.2	351	205	146	20	46	105	179	93	71	116	70
33.22	56	33	23	*	10	17	25	17	9	13	16
33.23	45	26	19	11	*6	13	16	15	9	12	10
33.24	137	82	55	*	19	39	77	35	29	48	25
33.26	26	13	13	*	*	*8	17	*	*6	11	*
33.27	62	34	28	*	*7	19	37	16	13	23	10
33.28	19	13	*6	*	*	*8	*6	*	*	*7	*
34	457	242	215	13	94	120	229	109	103	158	87
34.0	179	108	71	*8	51	50	70	39	40	65	35
34.03	*6	*	*	*	*	*	*	*	*	*	*
34.04	142	84	58	*7	41	39	55	32	33	53	23
34.09	24	15	*9	*	*8	*8	*7	*	*	*8	*8
34.2	51	25	26	*	*6	14	30	15	12	14	9
34.21	18	10	*7	*	*	*5	*8	*5	*	*6	*
34.22	12	*6	*7	—	*	*6	*6	*	*	*	*
34.24	11	*6	*6	—	*	*	9	*	*	*	*
34.5	14	10	*	*	*	*6	*	*	*	*	*
34.51	10	*7	*	*	*	*	*	*	*	*	*
34.6	*7	*	*	*	*	*	*	*	*	*	*
34.9	186	84	102	*	26	41	116	44	43	64	34
34.91	158	73	85	*	21	33	101	37	38	54	30
34.92	23	9	14	—	*	*8	11	*7	*	*8	*
35–39	5,444	3,231	2,213	164	544	1,882	2,853	1,209	1,285	1,942	1,007
35	116	66	50	25	14	27	50	37	25	33	20
35.1	10	*	*5	*	*	*	*	*	*	*	*
35.12	*5	*	*	*	*	*	*	*	*	*	*
35.2	69	41	28	*	*8	19	41	18	19	20	12
35.21	*9	*6	*	*	*	*	*6	*	*	*	*
35.22	41	27	14	*	*	12	24	10	11	14	*6
35.24	14	*6	*8	*	*	*	*8	*	*	*	*
35.3	9	*6	*	*	*	*	*	*	*	*	*
35.33	*7	*	*	*	*	*	*	*	*	*	*
35.6	*6	*	*	*	*	*	*	*	*	*	*
35.7	*8	*	*	*	*	*	*	*	*	*	*
35.71	*5	*	*	*	*	*	*	*	*	*	*
36	1,268	878	390	*	57	550	659	269	348	416	236
36.0	666	452	214	*	37	307	322	118	183	231	135
36.01	412	276	137	*	25	191	195	78	113	136	85
36.02	16	11	*	*	*	*5	*9	*	*	*6	*
36.05	54	36	18	—	*	23	30	*6	14	19	15
36.06	177	125	52	—	*8	84	85	31	50	66	31
36.1	598	424	175	*	20	242	335	150	164	183	101
36.11	50	32	18	—	*	21	25	14	17	13	*6
36.12	116	75	41	*	*	46	67	29	34	34	19
36.13	114	82	32	—	*	44	66	25	30	38	21
36.14	68	54	15	*	*	26	41	16	17	23	13
36.15	231	165	67	*	9	96	125	60	65	70	36
36.16	11	*9	*	—	*	*7	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
37	1,727	1,018	709	23	130	628	945	379	397	679	271
37.0	11	*6	*	*	*	*	*	*	*	*	*
37.1	12	*8	*	*	*	*	*	*	*	*	*
37.12	10	*7	*	*	*	*	*	*	*	*	*
37.2	1,241	754	488	16	105	524	596	257	295	495	194
37.21	49	29	20	*	*6	24	17	10	*6	23	10
37.22	897	556	342	*	72	402	422	168	220	365	146
37.23	215	119	96	10	12	70	123	58	47	83	26
37.25	*6	*	*	*	*	*	*	*	*	*	*
37.26	62	39	22	*	11	21	29	16	18	17	10
37.27	*7	*	*	*	*	*	*	*	*	*	*
37.3	29	15	13	*	*6	10	10	*9	*6	9	*5
37.34	18	*9	9	*	*	*6	*7	*5	*	*6	*5
37.6	61	41	20	*	*	28	31	16	15	19	10
37.61	56	38	18	*	*	25	30	14	14	18	10
37.7	186	94	92	*	*6	27	152	47	38	75	27
37.71	37	17	19	*	*	*	32	10	*7	15	*
37.72	81	41	40	*	*	10	69	20	16	35	11
37.76	14	*7	*7	*	*	*	12	*	*	*	*
37.78	40	24	16	*	*	9	27	10	*9	14	*6
37.8	155	77	78	*	*	19	132	37	28	65	25
37.80	9	*	*5	*	—	*	*9	*	*	*5	*
37.81	19	9	10	*	*	*	16	*	*	*8	*
37.82	23	11	11	*	*	*	20	*5	*	10	*
37.83	81	42	39	*	*	11	68	19	16	34	12
37.85	*5	*	*	*	*	*	*	*	*	*	*
37.86	*5	*	*	*	*	*	*5	*	*	*	*
37.87	10	*	*6	*	*	*	*7	*	*	*	*
37.9	26	19	*7	*	*	11	13	*6	*8	*7	*
37.94	19	15	*	*	*	*8	9	*	*7	*	*
38	1,117	583	534	82	188	266	580	253	265	375	224
38.0	46	18	28	*	*	10	30	13	*8	20	*
38.03	10	*	*7	*	*	*	*7	*	*	*7	*
38.08	24	10	14	—	*	*	18	*8	*5	*9	*
38.1	162	91	71	*	*	35	123	39	38	61	25
38.12	130	77	54	—	*	27	102	32	31	50	17
38.18	23	11	12	*	*	*5	16	*5	*	*7	*7
38.2	11	*	*7	*	*	*	*7	*	*	*	*
38.21	*8	*	*6	*	*	*	*5	*	*	*	*
38.3	*6	*	*	*	*	*	*	*	*	*	*
38.4	56	44	12	*	*	11	42	13	12	22	9
38.44	30	24	*6	—	*	*	25	*8	*7	10	*
38.45	*9	*5	*	*	*	*	*	*	*	*	*
38.46	*6	*6	*	—	*	*	*5	*	*	*	*
38.48	*9	*8	*	*	*	*	*6	*	*	*	*
38.5	*8	*	*6	—	*	*	*	*	*	*	*
38.59	*6	*	*	—	*	*	*	*	*	*	*
38.6	10	*6	*	*	*	*	*6	*	*	*	*
38.7	31	14	18	*	*	*6	22	10	*8	9	*
38.8	33	19	13	10	12	*	*8	10	*5	12	*5

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
38.85	11	*7	*	*8	*	*	*	*	*	*	*
38.86	*6	*	*	*	*	*	*	*	*	*	*
38.9	754	380	374	68	155	193	338	159	188	240	167
38.91	125	70	54	21	16	26	61	27	45	32	21
38.92	11	*7	*	10	*	*	*	*	*	*	*
38.93	496	238	258	33	110	128	225	110	115	157	114
38.94	10	*	*5	*	*	*	*	*	*	*	*
38.95	104	55	48	*	26	35	43	16	20	44	23
38.99	*6	*	*	*	*	*	*	*	*	*	*
39	1,217	687	530	32	155	411	620	272	250	439	256
39.1	*5	*	*	*	*	*	*	*	*	*	*
39.2	185	102	83	*	16	53	114	40	40	69	35
39.25	23	14	9	—	*	10	12	*6	*5	*9	*
39.27	58	29	29	*	11	18	28	10	11	22	16
39.29	95	57	39	*	*	24	68	23	23	37	13
39.3	20	12	*8	*	*9	*6	*	*	*	*7	*7
39.31	15	*9	*6	*	*7	*	*	*	*	*	*
39.4	89	46	43	*	20	30	38	17	12	38	22
39.42	22	9	13	—	*6	*7	*9	*	*	*8	*7
39.43	*8	*	*	—	*	*	*	*	*	*	*
39.49	56	31	25	*	11	20	25	11	*8	24	14
39.5	117	64	53	*5	14	35	62	29	22	44	21
39.50	60	34	25	*	*6	19	33	17	11	22	10
39.51	12	*	*8	—	*	*6	*	*	*	*	*
39.52	10	*7	*	*	*	*	*8	*	*	*6	*
39.57	*9	*	*	*	*	*	*5	*	*	*	*
39.59	15	*8	*6	*	*	*	*8	*	*	*6	*5
39.6	362	240	122	18	17	133	193	89	96	120	58
39.61	322	215	107	16	16	119	171	80	84	104	53
39.62	9	*6	*	—	*	*	*	*	*	*	*
39.63	*8	*	*	—	*	*	*	*	*	*	*
39.64	21	14	*7	*	*	*7	13	*	*	11	*
39.9	436	216	220	*	76	151	205	91	73	159	112
39.93	*7	*	*	*	*	*	*	*	*	*	*
39.95	413	205	209	*	73	143	195	86	69	152	106
39.97	*6	*	*	—	*	*	*	*	*	*	*
40–41	348	169	179	19	62	115	152	93	77	111	66
40	194	93	101	*	24	80	86	49	44	62	39
40.1	44	20	24	*	*8	18	17	14	10	15	*6
40.11	43	20	23	*	*7	18	17	14	10	14	*6
40.2	49	19	30	*	*8	19	20	14	12	16	*8
40.23	11	*	10	—	*	*	*	*	*	*	*
40.24	*7	*	*	*	*	*	*	*	*	*	*
40.29	27	13	15	*	*	10	12	*6	*8	*9	*
40.3	67	34	33	*	*5	27	34	16	12	21	18
40.4	12	*7	*5	—	*	*6	*6	*	*	*	*
40.41	11	*6	*5	—	*	*6	*5	*	*	*	*
40.5	20	13	*7	*	*	9	*9	*	*7	*6	*
40.53	*8	*7	*	—	*	*	*	*	*	*	*
41	154	76	78	15	38	36	66	45	33	49	27
41.0	13	*7	*6	*	*7	*	*	*	*	*	*
41.04	*8	*	*	*	*5	*	*	*	*	*	*
41.3	108	54	54	9	19	25	54	33	25	34	17
41.31	108	54	54	9	19	25	54	33	25	34	17
41.5	27	13	15	*	10	*6	10	*7	*5	*8	*7

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
42–54	4,976	2,152	2,824	218	1,206	1,247	2,305	1,135	1,182	1,729	930
42	149	80	69	12	20	39	78	36	31	51	31
42.2	82	45	37	10	12	20	40	21	16	28	17
42.23	76	42	34	*8	12	19	38	18	16	26	16
42.3	26	17	*9	*	*6	10	10	*6	*	10	*6
42.33	25	16	*8	*	*6	10	*9	*6	*	9	*6
42.9	29	11	18	*	*	*	22	*5	*9	10	*5
42.92	29	11	18	*	*	*	22	*5	*9	10	*5
43	216	110	105	21	18	34	143	48	43	85	39
43.1	170	82	88	10	13	23	124	35	32	71	31
43.11	134	65	69	*	10	15	105	28	23	59	24
43.19	36	17	19	*6	*	*8	19	*8	*9	12	*7
43.3	12	10	*	11	*	*	*	*	*	*	*
43.4	*9	*5	*	—	*	*	*	*	*	*	*
43.41	*7	*	*	—	*	*	*	*	*	*	*
43.7	*9	*5	*	—	*	*	*5	*	*	*	*
44	168	86	82	10	35	43	81	29	36	64	38
44.0	9	*5	*	*	*	*	*	*	*	*	*
44.1	28	12	16	*	*	*7	16	*	*6	14	*
44.13	17	*7	10	*	*	*	10	*	*	*9	*
44.14	9	*	*	*	*	*	*	*	*	*	*
44.2	11	*6	*6	*	*	*	*7	*	*	*	*
44.29	10	*	*	*	*	*	*6	*	*	*	*
44.3	17	*6	11	*	*7	*	*7	*	*	*6	*6
44.39	14	*6	*8	*	*	*	*7	*	*	*6	*
44.4	62	37	25	*	10	15	37	11	16	20	15
44.42	13	*8	*5	—	*	*	*6	*	*	*	*
44.43	43	26	17	*	*5	10	28	*7	10	14	12
44.6	36	17	19	*8	10	11	*7	*7	*8	16	*5
44.66	25	12	13	*6	*6	*8	*	*	*5	11	*
44.69	*5	*	*	*	*	*	*	*	*	*	*
45	1,828	827	1,001	26	258	440	1,103	421	448	659	300
45.0	*9	*5	*	*	*	*	*5	*	*	*	*
45.1	844	398	446	12	133	212	487	187	202	310	145
45.13	382	188	194	*	61	95	222	83	90	139	69
45.14	*8	*	*	*	*	*	*	*	*	*	*
45.16	450	203	247	*8	70	113	260	102	108	167	72
45.2	493	201	293	*5	63	111	315	115	119	184	75
45.22	*5	*	*	—	*	*	*	*	*	*	*
45.23	243	100	143	*	20	52	171	52	53	95	43
45.24	73	31	41	*	11	17	44	15	23	25	*9
45.25	167	65	102	*	32	39	94	46	38	61	23
45.3	13	*8	*	*	*	*	*	*	*	*	*
45.33	*7	*	*	*	*	*	*	*	*	*	*
45.4	106	49	56	*	*	19	82	26	26	37	16
45.42	87	42	45	—	*	16	69	23	22	31	11
45.43	13	*6	*8	*	*	*	11	*	*	*	*
45.6	61	27	34	*	9	16	33	16	17	19	9
45.62	59	26	32	*	9	15	33	15	16	18	*9
45.7	226	101	125	*	28	57	138	52	61	78	35
45.72	13	*	*8	*	*5	*	*	*	*	*	*
45.73	69	31	37	*	*5	14	49	15	22	23	*9
45.74	10	*	*8	*	*	*	*6	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
45.75	29	11	19	*	*	9	17	*8	*9	10	*
45.76	79	39	40	*	11	21	47	19	19	26	15
45.79	23	11	12	*	*	*7	13	*5	*5	10	*
45.8	*8	*5	*	*	*	*	*	*	*	*	*
45.9	66	30	36	*	12	15	37	17	14	22	12
45.91	11	*	*7	*	*	*	*6	*	*	*	*
45.93	26	12	14	*	*	*6	15	*6	*7	*8	*
45.94	19	*9	10	*	*	*	12	*	*	*6	*
46	208	106	102	13	47	46	103	54	51	71	33
46.0	16	*7	*9	*	*	*	*9	*	*	*5	*
46.03	11	*5	*6	*	*	*	*7	*	*	*	*
46.1	48	22	26	*	*7	*9	31	14	12	15	*7
46.10	22	10	12	*	*	*	15	*6	*6	*7	*
46.11	13	*8	*6	*	*	*	*7	*	*	*	*
46.13	13	*	*8	*	*	*	*8	*	*	*	*
46.2	10	*6	*	*	*	*	*	*	*	*	*
46.3	29	18	11	*	*	*7	15	*7	*9	*8	*6
46.32	*7	*	*	*	*	*	*	*	*	*	*
46.39	22	14	*8	*	*	*6	11	*5	*7	*6	*
46.4	12	*6	*6	*	*	*	*6	*	*	*	*
46.5	38	18	20	*	10	*9	17	*8	10	14	*6
46.51	*7	*	*	*	*	*	*	*	*	*	*
46.52	31	15	17	*	*7	*7	15	*6	*9	11	*
46.7	35	17	17	*	14	*8	10	10	*7	13	*
46.73	15	*7	*8	*	*5	*	*5	*	*	*5	*
46.75	*8	*	*	*	*	*	*	*	*	*	*
46.79	*5	*	*	*	*	*	*	*	*	*	*
46.8	16	9	*7	*	*	*	10	*	*	*7	*
46.81	*7	*	*	*	*	*	*	*	*	*	*
46.85	*7	*	*	*	*	*	*6	*	*	*	*
47	328	163	165	71	176	53	27	59	82	102	85
47.0	263	146	117	64	144	36	18	47	61	80	75
47.1	63	16	47	*6	31	17	*9	12	20	21	10
48	118	61	57	*5	26	24	63	35	29	32	22
48.2	45	25	20	*	10	*8	24	13	12	11	*8
48.23	24	15	9	*	*7	*	12	*8	*5	*6	*5
48.24	17	*8	9	*	*	*	10	*	*6	*	*
48.3	17	*9	*9	*	*	*	11	*6	*	*	*
48.36	13	*7	*6	*	*	*	*8	*	*	*	*
48.5	10	*	*5	—	*	*	*5	*	*	*	*
48.6	24	10	14	*	*	*5	14	*8	*	*7	*5
48.63	15	*6	*8	—	*	*	10	*6	*	*	*
48.69	*7	*	*	—	*	*	*	*	*	*	*
48.7	*6	*	*	*	*	*	*	*	*	*	*
48.8	11	*8	*	*	*	*	*	*	*	*	*
48.81	10	*8	*	*	*	*	*	*	*	*	*
49	57	34	23	*	25	19	10	9	17	19	12
49.0	9	*6	*	*	*	*	*	*	*	*	*
49.01	*6	*5	*	*	*	*	*	*	*	*	*
49.2	*6	*5	*	*	*	*	*	*	*	*	*
49.21	*6	*5	*	*	*	*	*	*	*	*	*
49.3	*6	*	*	*	*	*	*	*	*	*	*
49.4	21	12	*9	*	10	*7	*	*	*7	*7	*
49.46	16	*9	*7	*	*8	*6	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
50	112	53	60	*8	21	37	46	28	23	40	22
50.1	82	39	43	*5	12	24	40	19	18	30	14
50.11	53	26	28	*	*9	16	25	14	12	17	9
50.12	28	13	15	*	*	*8	14	*	*6	13	*
50.2	*6	*	*	*	*	*	*	*	*	*	*
50.6	*7	*	*	*	*	*	*	*	*	*	*
50.61	*6	*	*	*	*	*	*	*	*	*	*
50.9	*8	*	*6	*	*	*	*	*	*	*	*
50.91	*7	*	*	*	*	*	*	*	*	*	*
51	693	244	449	*7	196	205	285	164	152	239	138
51.1	78	34	45	*	22	19	38	20	17	23	18
51.10	67	29	38	*	18	17	31	18	15	19	16
51.2	448	142	306	*	136	147	160	112	97	151	88
51.22	142	61	80	*	27	42	71	29	34	47	31
51.23	307	81	226	*	109	105	90	83	63	104	57
51.3	13	*	*8	*	*	*	*7	*	*	*5	*
51.32	*5	*	*	—	*	*	*	*	*	*	*
51.36	*6	*	*	*	*	*	*	*	*	*	*
51.4	21	*6	15	*	*5	*	11	*	*6	*8	*
51.41	15	*	11	*	*	*	*8	*	*	*7	*
51.5	11	*6	*5	*	*	*	*8	*	*	*	*
51.51	11	*6	*	*	*	*	*8	*	*	*	*
51.8	104	43	60	*	27	26	50	20	25	38	20
51.84	*5	*	*	—	*	*	*	*	*	*	*
51.85	48	18	30	*	14	12	22	*9	11	19	*9
51.87	19	9	10	*	*	*	12	*	*	*6	*
51.88	26	10	16	*	*7	*6	13	*6	*6	10	*
51.9	11	*	*6	—	*	*	*8	*	*	*	*
51.98	*8	*	*	—	*	*	*6	*	*	*	*
52	34	17	17	*	10	*8	16	*8	*7	11	*8
52.1	10	*	*6	—	*	*	*8	*	*	*	*
52.11	*7	*	*	—	*	*	*6	*	*	*	*
52.9	10	*5	*	*	*	*	*	*	*	*	*
52.93	*7	*	*	—	*	*	*	*	*	*	*
53	229	117	112	15	40	65	110	65	55	74	35
53.0	62	50	12	*6	10	12	34	21	11	18	11
53.00	13	10	*	*	*	*	*8	*	*	*7	*
53.01	*8	*7	*	*	*	*	*5	*	*	*	*
53.02	13	*9	*	*	*	*	*	*	*	*	*
53.03	12	11	*	*	*	*	*6	*5	*	*	*
53.04	*8	*7	*	—	*	*	*6	*	*	*	*
53.05	*8	*6	*	*	*	*	*	*	*	*	*
53.1	15	13	*	*6	*	*	*6	*6	*	*5	*
53.12	*5	*	*	*	*	—	*	*	*	*	*
53.2	*6	*	*	—	*	*	*	*	*	*	*
53.4	46	17	30	*	10	19	16	12	12	15	*7
53.49	42	15	27	*	10	16	14	9	10	15	*7
53.5	47	15	32	*	12	12	23	10	13	18	*7
53.51	27	*8	19	*	*6	*7	14	*6	*7	11	*
53.59	20	*7	13	*	*6	*	9	*	*6	*7	*
53.6	41	16	25	*	*	16	20	11	11	13	*6
53.61	29	12	17	*	*	12	14	*9	*8	*8	*
53.69	12	*	*8	—	*	*	*6	*	*	*5	*
53.7	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
— Quantity zero.NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
54	838	255	582	27	335	235	240	178	209	282	168
54.0	16	*9	*8	*	*	*	*6	*	*	*	*
54.1	60	24	36	*	21	13	23	15	18	15	13
54.11	38	17	21	*	14	*9	14	10	12	*7	*9
54.12	*5	*	*	*	*	*	*	*	*	*	*
54.19	17	*5	12	*	*6	*	*7	*	*	*6	*
54.2	176	42	134	*	94	41	37	37	42	55	42
54.21	124	22	101	*	81	26	12	25	28	39	32
54.23	27	*9	18	*	*	*8	13	*6	*8	*7	*6
54.25	17	*6	11	*	*7	*	*6	*	*	*5	*
54.3	16	*	12	*	*6	*	*6	*	*	*6	*
54.4	50	*9	41	*	16	19	14	12	12	14	11
54.5	338	69	269	*	149	84	101	66	85	134	53
54.6	*9	*5	*	*	*	*	*	*	*	*	*
54.61	*6	*	*	*	*	*	*	*	*	*	*
54.7	9	*5	*	*	*	*	*	*	*	*	*
54.9	163	89	75	10	39	65	49	36	40	48	40
54.91	92	49	43	*	22	38	29	21	21	30	20
54.93	12	*6	*6	*	*	*	*	*	*	*	*
54.95	15	11	*	*6	*	*	*	*	*	*	*
54.98	37	19	18	*	*9	16	11	*8	12	*7	11
55–59	1,016	510	506	34	245	298	439	257	234	343	183
55	127	66	61	*8	40	46	32	34	22	48	23
55.0	25	10	14	*	*9	*7	*8	*7	*	*8	*5
55.03	15	*7	*8	*	*	*	*6	*	*	*	*
55.2	28	16	12	*	13	9	*	*8	*	12	*
55.23	24	13	10	*	12	*7	*	*8	*	11	*
55.5	39	22	16	*	*8	14	15	*9	*7	15	*8
55.51	36	20	16	*	*7	12	15	*8	*7	14	*7
55.6	15	*8	*7	*	*5	*9	*	*	*	*6	*
55.69	15	*8	*7	*	*5	*9	*	*	*	*6	*
55.8	*9	*	*	*	*	*	*	*	*	*	*
55.87	*8	*	*	*	*	*	*	*	*	*	*
55.9	*5	*	*	*	*	*	*	*	*	*	*
56	106	63	43	9	36	35	25	25	22	41	17
56.0	42	28	14	—	19	15	*8	*7	10	19	*6
56.2	*6	*	*	*	*	*	*	*	*	*	*
56.3	26	16	10	—	11	10	*6	*6	*	11	*
56.31	23	14	9	—	10	*8	*	*5	*	10	*
56.5	*9	*6	*	*	*	*	*	*	*	*	*
56.51	*8	*6	*	—	*	*	*	*	*	*	*
56.7	15	*5	10	*8	*	*	*	*5	*	*	*
56.74	14	*	*9	*7	*	*	*	*	*	*	*
57	503	285	218	12	88	118	285	140	124	152	88
57.0	21	15	*6	*	*	*6	13	*7	*8	*	*
57.1	43	16	28	*	10	11	20	*8	9	19	*7
57.17	16	*	12	*	*	*6	*6	*	*	*6	*
57.18	20	*8	12	*	*5	*	9	*	*	*9	*
57.19	*7	*	*	*	*	*	*	*	*	*	*
57.3	210	130	79	*5	36	49	119	54	61	68	28
57.32	189	118	71	*	33	43	108	48	58	61	23
57.33	18	10	*7	*	*	*5	10	*	*	*6	*
57.4	60	47	14	*	*	11	48	19	12	20	9
57.49	60	47	14	*	*	11	48	19	12	20	9
57.7	*9	*7	*	—	*	*	*5	*	*	*	*
57.71	*8	*6	*	—	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
57.8	25	*	20	*	10	*7	*6	*7	*6	*8	*
57.81	9	*	*8	—	*6	*	*	*	*	*	*
57.89	9	*	*9	*	*	*	*	*	*	*	*
57.9	126	60	65	*	28	29	68	41	24	28	33
57.91	*8	*8	—	—	*	*	*6	*	*	*	*
57.94	112	47	64	*	27	25	59	38	22	22	30
58	60	45	15	*	*8	15	34	16	14	21	9
58.4	*7	*	*	*	*	*	*	*	*	*	*
58.5	11	9	*	*	*	*	*7	*	*	*	*
58.6	29	22	*7	*	*	*8	18	*7	*9	10	*
59	220	52	169	*	72	84	63	42	52	80	46
59.0	9	*	*7	*	*	*	*	*	*	*	*
59.02	*6	*	*6	*	*	*	*	*	*	*	*
59.4	*9	—	*9	—	*	*	*	*	*	*	*
59.5	38	—	38	*	13	18	*6	*	*8	17	10
59.6	*6	—	*6	—	*	*	*	*	*	*	*
59.7	51	*	50	*	11	24	15	*6	13	19	13
59.79	49	—	49	—	11	23	14	*	13	18	13
59.8	101	46	55	*	41	29	31	25	27	34	14
60-64	302	302	—	29	15	66	192	74	78	97	53
60	223	223	—	—	*	52	169	49	60	73	40
60.1	14	14	—	—	*	*	11	*	*	*6	*
60.11	14	14	—	—	*	*	11	*	*	*6	*
60.2	145	145	—	—	*	23	121	33	40	45	27
60.21	9	9	—	—	*	*	*8	*	*	*	*
60.29	136	136	—	—	*	22	113	29	39	43	25
60.5	48	48	—	—	*	23	25	10	12	17	10
61	*8	*8	—	*	*	*	*	*	*	*	*
61.0	*5	*5	—	*	*	*	*	*	*	*	*
62	18	18	—	*	*	*	*7	*6	*	*	*
62.3	*6	*6	—	*	*	*	*	*	*	*	*
62.4	*5	*5	—	—	*	*	*	*	*	*	*
62.41	*5	*5	—	—	*	*	*	*	*	*	*
63	*8	*8	—	*	*	*	*	*	*	*	*
64	45	45	—	21	*	*8	11	12	10	15	*9
64.0	27	27	—	20	*	*	*	*8	*6	*8	*5
64.9	13	13	—	*	*	*7	*	*	*	*5	*
64.97	*8	*8	—	—	*	*	*	*	*	*	*
65-71	2,096	—	2,096	9	1,272	575	240	358	418	875	446
65	574	—	574	*	291	220	59	95	124	226	128
65.2	48	—	48	*	41	*	*	12	10	17	10
65.29	46	—	46	*	39	*	*	12	*9	16	10
65.3	17	—	17	*	13	*	*	*	*	*	*
65.4	74	—	74	*	52	14	*8	12	18	28	16
65.5	*9	—	*9	—	*	*	*	*	*	*	*
65.51	*7	—	*7	—	*	*	*	*	*	*	*
65.6	375	—	375	*	144	185	45	55	81	155	84
65.61	362	—	362	*	136	182	44	54	79	150	79
65.62	13	—	13	—	*8	*	*	*	*	*	*
65.8	31	—	31	—	21	*8	*	*6	*	12	*9
65.9	*9	—	*9	*	*8	*	—	*	*	*	*
65.91	*8	—	*8	*	*7	*	—	*	*	*	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
66	426	—	426	*	419	*	*	59	65	206	95
66.0	12	—	12	—	12	—	*	*	*	*	*
66.02	10	—	10	—	10	—	*	*	*	*	*
66.2	19	—	19	—	19	*	—	*	*	*7	*
66.29	15	—	15	—	15	—	—	*	*	*6	*
66.3	324	—	324	—	323	*	—	39	47	163	74
66.32	181	—	181	—	181	*	—	21	27	85	48
66.39	142	—	142	—	142	*	—	18	20	78	27
66.4	10	—	10	*	*8	*	*	*	*	*	*
66.5	*7	—	*7	—	*6	*	*	*	*	*	*
66.6	40	—	40	*	38	*	*	*6	*7	20	*7
66.61	*7	—	*7	*	*6	*	*	*	*	*	*
66.62	22	—	22	—	22	—	—	*	*	11	*
66.69	9	—	9	*	*8	*	*	*	*	*6	*
66.7	*6	—	*6	*	*6	—	—	*	*	*	*
67	19	—	19	—	11	*	*	*5	*	*6	*
68	646	—	646	*	349	226	69	111	139	266	130
68.1	17	—	17	—	*8	*	*	*5	*	*7	*
68.12	*8	—	*8	—	*	*	*	*	*	*	*
68.16	*7	—	*7	—	*	*	*	*	*	*	*
68.2	37	—	37	—	32	*	*	11	*6	11	*9
68.29	37	—	37	—	31	*	*	11	*6	11	*9
68.3	9	—	9	*	*	*	*	*	*	*	*
68.4	393	—	393	*	201	155	37	63	86	167	77
68.5	183	—	183	—	102	56	26	27	41	77	38
69	137	—	137	*	112	17	*7	42	22	52	20
69.0	83	—	83	*	66	12	*	28	12	30	13
69.02	45	—	45	*	45	*	—	12	*7	18	*8
69.09	36	—	36	*	19	12	*	15	*	12	*
69.1	*6	—	*6	—	*	*	*	*	*	*	*
69.19	*6	—	*6	—	*	*	*	*	*	*	*
69.2	*7	—	*7	—	*5	*	*	*	*	*	*
69.22	*6	—	*6	—	*	*	*	*	*	*	*
69.5	26	—	26	*	25	*	*	10	*	*9	*
69.52	19	—	19	*	18	*	—	*8	*	*6	*
69.9	13	—	13	—	11	*	*	*	*	*7	*
69.93	*5	—	*5	—	*5	*	—	*	*	*	*
70	264	—	264	*	75	95	91	39	56	107	62
70.2	*7	—	*7	*	*	*	*	*	*	*	*
70.3	13	—	13	*	10	*	*	*	*	*	*
70.32	*6	—	*6	*	*5	*	*	*	*	*	*
70.33	*7	—	*7	*	*	*	*	*	*	*	*
70.5	151	—	151	—	35	62	54	23	33	61	34
70.50	76	—	76	—	14	29	33	11	14	33	18
70.51	40	—	40	—	*9	20	12	*7	12	12	*9
70.52	35	—	35	—	12	13	9	*	*6	16	*7
70.7	40	—	40	*	10	14	14	*	*6	19	10
70.77	28	—	28	*	*5	10	13	*	*	13	*8
70.79	*6	—	*6	*	*	*	*	*	*	*	*
70.9	40	—	40	—	10	16	14	*	*9	18	10
70.92	40	—	40	—	10	16	14	*	*8	18	10

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
71	31	—	31	*	15	*8	*7	*6	*7	11	*7
71.3	*8	—	*8	*	*6	*	*	*	*	*	*
71.7	11	—	11	*	*	*	*	*	*	*	*
71.79	*8	—	*8	—	*	*	*	*	*	*	*
72–75	6,540	—	6,540	18	6,519	*	—	1,215	1,422	2,061	1,841
72	446	—	446	*	443	*	—	60	96	145	145
72.0	20	—	20	*	20	—	—	*	*	12	*
72.1	110	—	110	*	108	—	—	16	25	50	18
72.2	*8	—	*8	—	*8	—	—	*	*	*	*
72.21	*7	—	*7	—	*7	—	—	*	*	*	*
72.4	*6	—	*6	—	*6	—	—	*	*	*	*
72.5	*7	—	*7	—	*7	—	—	*	*	*	*
72.7	293	—	293	*	292	*	—	38	61	74	119
72.71	200	—	200	*	200	*	—	29	40	51	80
72.79	93	—	93	*	92	—	—	9	21	23	39
73	3,143	—	3,143	9	3,133	*	—	553	664	1,072	854
73.0	729	—	729	*	727	*	—	120	187	185	237
73.01	103	—	103	*	103	—	—	17	23	26	37
73.09	626	—	626	*	624	*	—	103	165	158	201
73.1	*7	—	*7	*	*7	—	—	*	*	*	*
73.4	389	—	389	*	388	—	—	63	91	103	132
73.5	1,031	—	1,031	*	1,028	*	—	177	160	420	273
73.51	*9	—	*9	—	*9	—	—	*	*	*	*
73.59	1,022	—	1,022	*	1,019	*	—	176	158	419	270
73.6	977	—	977	*	974	*	—	189	222	359	207
74	837	—	837	*	834	*	—	139	159	333	206
74.0	14	—	14	*	14	—	—	*	*	*6	*
74.1	804	—	804	*	801	*	—	135	156	319	195
74.9	17	—	17	—	17	—	—	*	*	*7	*8
74.99	17	—	17	—	17	—	—	*	*	*7	*8
75	2,115	—	2,115	*	2,109	*	—	464	503	512	636
75.1	22	—	22	*	21	—	—	*	*	*6	9
75.3	955	—	955	*	952	*	—	265	245	149	295
75.32	311	—	311	*	309	*	—	66	99	53	93
75.34	584	—	584	*	582	*	—	187	122	89	187
75.35	58	—	58	*	58	*	—	11	24	*8	15
75.4	48	—	48	—	48	—	—	*8	11	16	13
75.5	*8	—	*8	—	*8	—	—	*	*	*	*
75.51	*7	—	*7	—	*7	—	—	*	*	*	*
75.6	1,053	—	1,053	*	1,050	*	—	182	234	324	313
75.61	45	—	45	*	45	—	—	*7	15	13	9
75.62	120	—	120	*	120	—	—	23	25	42	30
75.69	887	—	887	*	886	*	—	152	193	269	274
75.7	23	—	23	—	23	—	—	*	*	13	*
76–84	3,134	1,546	1,589	165	960	841	1,168	713	748	1,079	594
76	94	61	33	*6	73	10	*	22	20	29	23
76.6	22	*9	13	*	19	*	*	*5	*7	*6	*
76.62	*6	*	*	*	*6	*	*	*	*	*	*
76.64	*6	*	*	*	*5	*	*	*	*	*	*
76.65	*6	*	*	*	*6	*	—	*	*	*	*
76.7	47	37	9	*	38	*	*	10	*8	16	13
76.72	*6	*	*	*	*5	*	*	*	*	*	*
76.75	*7	*6	*	*	*5	*	—	*	*	*	*
76.76	18	16	*	*	15	*	*	*	*	*7	*
76.79	*8	*5	*	*	*6	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
76.9	14	9	*	*	10	*	*	*	*	*	*
76.92	*6	*	*	*	*	*	*	*	*	*	*
77	306	150	156	22	109	103	73	76	68	105	57
77.1	*6	*	*	*	*	*	*	*	*	*	*
77.2	*8	*	*6	*	*	*	*	*	*	*	*
77.27	*6	*	*	*	*	*	*	*	*	*	*
77.3	17	*8	*9	*6	*	*6	*	*6	*	*	*
77.37	*7	*	*	*	*	*	*	*	*	*	*
77.4	17	*9	*9	*	*	*	9	*	*6	*6	*
77.49	*8	*	*	*	*	*	*5	*	*	*	*
77.5	22	*5	17	*	*6	*6	10	*	*	*7	*7
77.51	*6	*	*5	—	*	*	*	*	*	*	*
77.56	*7	*	*5	—	*	*	*	*	*	*	*
77.6	43	24	19	*	14	14	12	13	10	15	*5
77.65	*7	*	*	*	*	*	*	*	*	*	*
77.67	*6	*	*	*	*	*	*	*	*	*	*
77.68	*6	*	*	*	*	*	*	*	*	*	*
77.69	10	*	*5	*	*	*	*	*	*	*	*
77.7	119	61	57	*6	52	45	16	26	26	47	20
77.79	107	56	51	*	48	41	14	23	23	43	18
77.8	63	31	32	*	18	23	21	17	15	18	13
77.81	18	9	*9	*	*	*8	*6	*	*5	*	*
77.88	10	*	*6	—	*	*	*	*	*	*	*
77.89	20	10	10	*	*	*8	*8	*	*5	*7	*
77.9	*7	*	*	*	*	*	*	*	*	*	*
78	203	93	110	21	67	57	58	51	44	70	38
78.0	41	19	22	*	16	15	*8	*9	*8	17	*8
78.05	*8	*	*	*	*	*	*	*	*	*	*
78.07	12	*5	*7	*	*	*6	*	*	*	*6	*
78.09	*8	*	*	*	*	*	*	*	*	*	*
78.1	25	15	10	*	9	*6	*6	*6	*5	10	*
78.13	*8	*	*6	*	*	*	*	*	*	*	*
78.17	*9	*6	*	*	*	*	*	*	*	*	*
78.4	*5	*	*	*	*	*	*	*	*	*	*
78.5	56	22	35	*7	17	9	23	16	11	19	11
78.55	32	*9	23	*5	*	*	19	*8	*6	10	*8
78.57	*9	*	*	*	*	*	*	*	*	*	*
78.59	*8	*	*	*	*	*	*	*	*	*	*
78.6	60	29	31	*	19	17	19	14	16	19	11
78.65	22	11	11	*	*	*	11	*6	*7	*6	*
78.67	9	*	*	*	*	*	*	*	*	*	*
78.69	17	9	*8	*	*7	*7	*	*	*	*6	*
78.9	9	*	*6	—	*	*	*	*	*	*	*
78.99	*8	*	*5	—	*	*	*	*	*	*	*
79	678	304	374	63	208	124	283	150	166	233	130
79.0	91	47	44	19	29	15	28	27	22	26	15
79.01	11	*	*8	*	*	*	*6	*	*	*	*
79.02	27	12	15	*6	*6	*	12	*8	*6	*8	*
79.05	11	*6	*	*5	*	*	*	*	*	*	*
79.06	30	18	11	*5	12	*8	*5	*8	*8	10	*
79.1	71	33	38	12	16	12	32	12	22	22	15
79.11	10	*	*5	*6	*	*	*	*	*	*	*
79.12	*8	*	*	*	*	*	*	*	*	*	*
79.15	38	12	26	*	*5	*6	24	*6	12	12	*8
79.16	10	*8	*	*	*	*	*	*	*	*	*
79.2	*8	*6	*	*	*	*	*	*	*	*	*
79.3	432	170	262	17	128	82	205	92	104	157	78

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
79.31	32	11	21	*5	*8	*8	10	*7	*7	10	*7
79.32	33	18	15	*	17	*6	*6	*7	*8	9	*9
79.34	12	11	*	*	*5	*	*	*	*	*	*
79.35	193	52	141	*	16	17	158	42	51	72	28
79.36	133	59	75	*	64	39	28	26	30	52	25
79.37	10	*8	*	*	*7	*	*	*	*	*	*
79.39	13	*7	*6	*	*8	*	*	*	*	*	*
79.6	40	30	10	*9	20	*7	*	*7	10	14	10
79.62	10	*7	*	*	*	*	*	*	*	*	*
79.64	*6	*	*	*	*	*	*	*	*	*	*
79.66	15	11	*	*	9	*	*	*	*	*7	*
79.7	28	13	14	*	*8	*5	12	*6	*6	*9	*7
79.75	17	*6	11	*	*	*	9	*	*	*6	*
79.8	*6	*	*	*	*	*	*	*	*	*	*
80	464	265	199	*8	212	167	77	99	95	182	88
80.0	10	*6	*	—	*	*	*6	*	*	*	*
80.1	26	16	10	*	12	*5	*7	*5	*	10	*7
80.16	12	*8	*	*	*5	*	*	*	*	*	*
80.2	41	24	17	*	24	11	*	13	10	11	*7
80.21	12	*6	*6	—	*5	*	*	*	*	*	*
80.26	28	17	11	*	18	*6	*	9	*8	*7	*
80.4	16	*7	9	*	*	*5	*	*7	*	*	*
80.46	9	*	*7	*	*	*	*	*	*	*	*
80.5	288	164	124	*	134	117	36	49	59	128	51
80.51	285	162	123	*	132	116	35	49	58	127	51
80.6	24	12	12	*	14	*8	*	*7	*	*7	*5
80.7	14	*6	*9	*	*6	*	*	*	*	*	*
80.76	*9	*	*	*	*	*	*	*	*	*	*
80.8	33	23	10	*	12	10	10	*8	*7	11	*7
80.81	*6	*	*	—	*	*	*	*	*	*	*
80.86	14	10	*	*	*	*	*5	*	*	*	*
80.9	10	*7	*	*	*	*	*	*	*	*	*
80.91	*6	*	*	—	*	*	*	*	*	*	*
81	966	415	551	15	187	254	509	209	260	325	173
81.0	187	96	91	*6	78	78	26	32	42	83	30
81.02	87	47	40	*	35	45	*7	13	20	46	10
81.03	*7	*	*	*	*	*	*	*	*	*	*
81.05	11	*6	*5	*	*	*	*	*	*	*	*
81.06	*8	*	*	*	*	*	*	*	*	*	*
81.07	16	*7	9	—	*	*7	*	*	*6	*6	*
81.08	41	19	22	*	17	15	*8	*8	*7	15	10
81.09	11	*7	*	*	*7	*	*	*	*	*6	*
81.1	16	*6	10	*	*5	*7	*	*	*	*	*
81.11	*7	*	*	*	*	*	*	*	*	*	*
81.2	10	*	*6	*	*5	*	*	*	*	*	*
81.4	70	36	33	*	47	10	10	20	15	22	13
81.40	*6	*	*5	*	—	*	*	*	*	*	*
81.45	35	20	16	*	31	*	*	10	*9	11	*
81.47	19	11	*8	*	*9	*6	*	*7	*	*5	*
81.5	536	191	345	*	19	116	400	111	155	173	98
81.51	138	57	81	—	*8	35	95	29	42	40	27
81.52	103	25	78	—	*	*9	93	23	28	31	21
81.53	30	14	16	—	*	9	17	10	*5	11	*
81.54	245	88	157	—	*	59	181	46	75	82	42
81.55	20	*7	12	*	*	*	14	*	*	*8	*
81.8	69	35	34	*	18	25	25	18	20	18	14
81.80	*5	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
81.81	9	*	*7	—	*	*	*6	*	*	*	*
81.82	*7	*	*	*	*5	*	*	*	*	*	*
81.83	45	25	20	—	11	20	14	13	13	11	*9
81.9	75	44	31	*6	15	14	40	21	21	22	11
81.91	50	30	20	*5	*9	*8	28	14	13	15	*8
81.92	14	*6	*8	*	*	*	10	*	*	*	*
82	35	20	14	*	19	*8	*	9	*8	12	*5
82.0	10	*	*6	*	*	*	*	*	*	*	*
82.4	11	*8	*	*	*8	*	*	*	*	*	*
82.45	*5	*	*	*	*	*	*	*	*	*	*
83	216	129	88	25	65	67	60	57	49	59	51
83.0	13	9	*	*	*6	*	*	*	*	*	*
83.09	*6	*	*	*	*	*	*	*	*	*	*
83.1	26	15	11	9	*8	*	*	*8	*5	*7	*5
83.13	*7	*	*	*	*	*	*	*	*	*	*
83.14	13	10	*	*	*6	*	*	*	*	*	*
83.2	12	*7	*	*	*	*	*	*	*	*	*
83.21	11	*6	*	*	*	*	*	*	*	*	*
83.3	14	*9	*5	*	*	*	*	*5	*	*	*
83.39	11	*6	*	*	*	*	*	*	*	*	*
83.4	18	11	*8	*	*6	*6	*6	*5	*	*5	*
83.45	9	*6	*	*	*	*	*	*	*	*	*
83.5	10	*6	*	*	*	*	*	*	*	*	*
83.6	73	45	27	*	19	28	23	14	20	21	18
83.63	51	29	22	*	*6	24	20	10	14	14	12
83.64	14	11	*	*	*9	*	*	*	*	*	*
83.65	*7	*	*	*	*	*	*	*	*	*	*
83.7	*8	*	*	*	*	*	*	*	*	*	*
83.8	31	18	13	*8	9	*7	*7	10	*6	*8	*7
83.82	*7	*5	*	*	*	*	*	*	*	*	*
83.85	10	*5	*	*6	*	*	*	*	*	*	*
83.88	*6	*	*	*	*	*	*	*	*	*	*
83.9	12	*6	*5	*	*	*	*	*	*	*	*
84	171	108	63	*	20	51	99	39	40	64	29
84.0	15	13	*	*	*6	*7	*	*	*	*5	*
84.01	14	12	*	*	*	*6	*	*	*	*	*
84.1	141	85	56	*	11	41	90	34	33	52	22
84.11	44	27	18	—	*	15	26	12	12	15	*6
84.12	16	11	*	*	*	*6	9	*5	*	*6	*
84.15	42	27	15	—	*	12	25	11	*8	15	*8
84.17	36	18	18	—	*	*7	28	*6	10	15	*6
84.3	13	*8	*	*	*	*	*7	*	*	*6	*
85-86	1,290	572	718	78	402	360	450	323	281	442	245
85	210	*	205	*	54	83	73	50	47	68	44
85.1	18	*	17	—	*	*6	*9	*5	*	*6	*
85.11	*7	*	*7	—	*	*	*	*	*	*	*
85.12	*9	*	*8	—	*	*	*	*	*	*	*
85.2	37	*	36	—	*7	14	16	12	*6	10	*9
85.21	25	*	24	—	*5	10	10	*9	*	*8	*
85.23	10	*	10	—	*	*	*6	*	*	*	*
85.3	22	*	21	*	16	*	*	*9	*	*	*
85.32	20	*	19	*	15	*	*	*8	*	*	*
85.4	89	*	87	*	10	36	42	15	26	32	16
85.41	13	—	13	—	*	*	*7	*	*	*	*
85.43	72	*	71	*	*8	31	34	12	23	26	11

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
85.5	*7	*	*6	—	*	*	*	*	*	*	*
85.7	12	—	12	—	*	*6	*	*	*	*	*
85.8	*6	*	*6	*	*	*	*	*	*	*	*
85.9	13	—	13	*	*	*7	*	*	*	*	*
85.95	*6	—	*6	*	*	*	*	*	*	*	*
86	1,080	568	513	78	347	278	378	272	234	374	201
86.0	300	148	152	32	91	82	96	76	61	102	61
86.01	13	*6	*7	*	*	*	*	*	*	*	*
86.04	107	58	49	10	40	27	29	26	22	33	26
86.05	24	10	13	*	10	*7	*	*5	*	*8	*7
86.06	*9	*6	*	*	*	*	*	*	*	*	*
86.07	122	55	67	13	25	33	50	35	23	44	19
86.09	25	13	12	*	*7	*8	*7	*7	*	*8	*
86.1	31	16	15	*	*7	*7	15	11	*6	9	*
86.11	31	16	14	*	*7	*7	15	11	*6	9	*
86.2	390	219	172	19	103	108	160	101	90	138	61
86.22	279	163	116	13	79	80	108	72	58	106	43
86.23	15	*9	*6	*	*	*7	*6	*	*5	*6	*
86.27	31	14	17	*	*	*5	24	13	10	*	*
86.28	61	32	29	*	19	16	22	13	16	20	13
86.3	61	19	41	*	24	15	19	13	12	21	14
86.4	14	*6	*8	*	*	*	*7	*	*	*5	*
86.5	160	98	62	11	76	26	47	35	40	50	35
86.59	159	97	62	11	75	26	47	35	40	50	35
86.6	65	38	27	*6	23	16	20	18	14	22	11
86.69	49	28	22	*	16	12	18	14	12	15	*8
86.7	28	15	13	*	*8	*8	10	*7	*5	12	*
86.74	15	*8	*7	*	*	*	*6	*	*	*6	*
86.8	30	*8	22	*	12	13	*	*6	*	13	*6
86.83	14	*	13	*	*6	*7	*	*	*	*	*
86.89	*7	*	*	*	*	*	*	*	*	*	*
87–99	12,604	6,155	6,449	841	2,687	3,263	5,813	3,791	2,515	3,434	2,864
87	1,233	587	646	59	283	305	586	331	207	328	367
87.0	519	253	267	29	99	104	287	172	77	115	155
87.03	516	250	266	29	98	102	287	171	77	113	155
87.2	45	24	21	*	14	17	14	*5	10	22	*8
87.21	35	19	17	*	10	14	10	*	*6	20	*
87.4	216	112	104	9	42	54	110	39	36	42	98
87.41	100	58	43	*	21	28	49	36	19	26	19
87.44	108	51	57	*7	19	24	58	*	16	14	76
87.49	*6	*	*	*	*	*	*	*	*	*	*
87.5	165	51	114	*	51	51	60	37	34	61	33
87.51	*6	*	*	*	*	*	*	*	*	*	*
87.53	152	45	107	*	48	48	54	35	31	56	30
87.6	96	44	52	*7	15	21	53	30	14	24	27
87.61	19	10	9	*	*	*	11	*6	*	*5	*
87.62	38	17	21	*	*7	*8	18	14	*	10	11
87.63	12	*6	*6	*	*	*	*6	*	*	*	*
87.64	19	*9	11	*	*	*	14	*	*	*	*6
87.7	186	101	85	9	61	55	61	45	35	63	43
87.73	56	30	26	*	25	18	12	18	*9	16	14
87.74	96	54	43	*	27	31	36	21	21	38	16
87.76	*6	*	*	*	*	*	*	*	*	*	*
87.77	*7	*	*	*	*	*	*	*	*	*	*
87.79	12	*7	*5	*	*	*	*	*	*	*	*8

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
88	3,824	1,989	1,835	96	594	1,249	1,885	972	809	1,183	860
88.0	268	129	139	9	71	71	116	87	42	58	81
88.01	263	126	137	9	70	70	114	87	42	57	77
88.1	23	11	13	*	*7	*	10	*	*	*9	11
88.19	19	*8	11	*	*5	*	*8	*	*	*7	9
88.2	11	*	*7	*	*	*	*7	*	*	*	*7
88.3	122	62	61	*7	35	30	50	41	12	40	29
88.38	113	58	55	*5	34	27	46	40	11	35	27
88.4	385	213	172	*6	40	119	220	87	98	146	54
88.40	*6	*	*	*	*	*	*	*	*	*	*
88.41	117	54	63	*	15	35	66	21	34	47	15
88.42	96	58	38	*	*8	29	58	20	23	40	13
88.43	17	*8	*8	*	*	*7	*6	*6	*	*	*
88.45	11	*	*7	*	*	*	*	*	*	*6	*
88.47	17	11	*6	*	*	*	11	*	*	*	*
88.48	95	60	35	*	*	27	63	29	22	31	13
88.49	23	12	11	*	*	10	*9	*	*	9	*
88.5	1,580	952	627	10	131	693	745	315	390	631	244
88.50	*6	*	*	*	*	*	*	*	*	*	*
88.51	*7	*	*	*	*	*	*	*	*	*	*
88.53	654	393	260	*	56	292	305	129	156	265	103
88.54	33	19	15	*	*	11	16	*7	*8	15	*
88.55	15	*9	*6	*	*	*6	*7	*	*	*7	*
88.56	765	457	308	*	64	335	365	159	203	289	115
88.57	95	64	31	*	*	43	46	12	15	50	18
88.6	26	14	12	*	*7	*8	10	*6	*8	*6	*6
88.66	11	*7	*	*	*	*	*5	*	*	*	*
88.67	*9	*	*6	*	*	*	*	*	*	*	*
88.7	1,177	501	676	43	252	264	617	359	203	240	374
88.71	108	50	58	*	*5	19	79	27	22	18	40
88.72	558	270	288	17	57	139	345	178	101	132	147
88.73	13	*5	*8	*	*	*	*9	*	*	*	*5
88.74	43	18	25	*	15	11	17	9	*6	16	11
88.75	78	39	40	10	19	16	33	25	9	16	28
88.76	185	70	115	10	64	44	68	56	25	22	82
88.77	101	44	58	*	16	26	59	36	23	14	28
88.78	61	—	61	*	59	—	*	16	10	10	25
88.79	29	*6	24	*	14	*7	*6	11	*	*8	*6
88.9	232	104	128	16	48	58	109	74	52	52	55
88.91	128	58	71	11	26	28	63	37	31	30	30
88.93	58	28	31	*	13	15	28	19	15	13	12
88.94	12	*	*8	*	*	*	*	*	*	*	*
88.97	29	12	16	*	*	11	13	11	*	*6	*9
89	1,239	600	639	65	190	337	647	372	213	262	393
89.0	*6	*	*	*	*	*	*	*	*	—	*
89.1	125	57	69	27	29	22	47	54	13	32	27
89.14	98	45	52	17	20	19	42	42	11	22	23
89.15	*6	*	*	*	*	*	*	*	*	*	*
89.19	16	*6	10	*6	*7	*	*	*6	*	*7	*
89.2	31	12	19	*	*8	*	17	10	*9	10	*
89.22	14	*7	*6	*	*	*	11	*6	*	*	*
89.26	9	—	9	*	*	*	*	*	*	*	*
89.3	56	25	30	11	11	16	17	15	18	9	13
89.37	28	11	17	*	*8	*9	9	*7	*8	*	10
89.38	14	*8	*5	*	*	*	*	*6	*6	*	*
89.39	11	*	*7	*	*	*	*	*	*	*5	*
89.4	154	81	73	*	24	64	66	58	21	35	40

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
89.41	61	34	26	*	11	29	21	21	*6	11	23
89.44	92	46	46	*	13	35	44	37	15	24	17
89.5	527	254	274	12	66	148	301	135	75	104	213
89.50	24	9	14	*	*	*	17	16	*	*	*
89.51	10	*5	*	*	*	*	*6	*	—	—	9
89.52	184	89	95	*6	25	53	100	*9	29	36	111
89.54	299	146	153	*	36	87	172	106	42	63	88
89.59	*8	*	*	*	*	*	*6	*	*	*	*
89.6	337	168	170	12	49	81	195	96	76	71	94
89.61	11	*7	*	*	*	*	*	*	*	*	*
89.62	20	10	11	*	*	*	12	*7	*	*	*
89.64	80	43	36	*	*7	22	51	27	24	15	14
89.65	202	95	107	*7	34	45	116	54	41	47	59
89.68	19	10	10	*	*	*	9	*	*	*	13
90	30	12	17	*	*6	*5	15	*	*7	18	*
90.5	14	*6	*9	*	*	*	*8	*	*	12	*
90.53	*6	*	*	—	*	*	*	*	—	*6	—
90.8	*8	*	*5	—	*	*	*	*	*	*	*
90.81	*8	*	*	—	*	*	*	*	*	*	*
91	13	*6	*7	*	*	*	*8	*	*	*8	*
91.3	10	*	*6	*	*	*	*6	*	*	*6	*
92	403	170	233	*8	61	114	221	144	64	82	113
92.0	135	60	74	*	22	43	66	48	15	27	45
92.02	20	*7	14	*	*	*6	9	*	*	*	10
92.03	14	*7	*7	*	*	*	*7	*6	*	*	*
92.04	17	*7	9	*	*	*	*7	*	*	*	*7
92.05	80	38	42	*	*8	30	42	31	9	17	23
92.1	164	66	97	*	25	44	92	61	27	25	50
92.14	56	22	34	*	*7	12	35	22	9	9	15
92.15	84	34	50	*	13	26	45	26	15	14	29
92.19	12	*6	*6	*	*	*	*8	*6	*	*	*
92.2	102	43	59	*	14	26	60	33	22	29	17
92.23	*8	*	*	—	*	*	*5	*	*	*	*
92.24	24	10	15	*	*	*7	14	*7	*6	*	*8
92.27	11	*	*8	—	*	*	*5	*	*	*	*
92.29	52	24	28	*	*6	13	31	17	11	17	*8
93	1,477	675	802	189	235	278	775	471	335	354	317
93.0	17	*8	*9	*	*	*	*8	*	*8	*	*
93.01	*8	*	*	*	*	*	*	*	*6	*	*
93.08	*7	*	*	*	*	*	*	*	*	*	*
93.1	17	*7	10	*	*	*	11	10	*	*	*
93.2	89	36	52	*	10	20	57	42	21	*7	19
93.22	73	30	42	*	*6	16	50	34	17	*	17
93.24	*7	*	*	*	*	*	*	*5	*	*	*
93.26	*6	*	*	—	*	*	*	*	*	*	*
93.3	329	138	190	*5	36	68	219	96	79	99	54
93.32	18	10	*7	*	*	*6	*6	*	*	10	*
93.38	10	*	*7	*	*	*	*7	*	*	*6	*
93.39	293	121	172	*	30	57	203	90	76	78	49
93.4	23	10	13	*	*	*	15	*5	*5	*8	*
93.46	15	*	10	*	*	*	13	*	*	*	*
93.5	93	49	44	14	35	15	28	34	20	24	14
93.53	20	11	*9	*6	*6	*	*	*8	*6	*	*
93.54	15	*7	*8	*	*7	*	*	*	*	*	*
93.55	*6	*	*	*	*	*	—	*	*	*	*
93.57	16	9	*6	*	*7	*	*	*5	*	*7	*
93.59	28	13	15	*	*8	*5	13	12	*7	*6	*

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
93.6	*5	*	*	—	*	*	*	*	*	*	*
93.7	30	15	15	*	*	*6	21	*7	13	*6	*5
93.75	27	13	15	*	*	*6	18	*7	12	*	*
93.8	258	107	151	*	54	42	158	91	71	69	27
93.81	37	15	22	*	15	*6	15	*7	10	21	*
93.83	213	89	124	*	37	33	139	83	61	45	24
93.89	*6	*	*	*	*	*	*	*	*	*	*
93.9	616	304	313	159	85	117	255	182	113	131	190
93.90	28	15	12	*6	*	*7	11	*8	*8	*7	*
93.94	266	135	131	89	38	47	92	87	31	68	80
93.96	261	123	138	51	32	51	127	64	57	43	96
93.99	52	26	26	12	10	9	21	23	13	11	*5
94	708	421	287	39	462	146	61	331	195	137	46
94.0	*9	*	*	*	*	*	*	*	*	*5	*
94.1	*6	*	*	*	*	*	*	*	—	*	*
94.2	145	69	76	*7	81	32	24	92	21	19	13
94.22	9	*	*	*	*5	*	*	*6	*	*	*
94.23	18	10	*8	*	10	*5	*	14	—	*	*
94.25	84	42	43	*	52	19	*8	59	12	*6	*8
94.27	33	12	20	—	13	*6	14	12	*9	11	*
94.3	92	44	47	11	56	16	9	53	32	*6	*
94.38	32	13	19	*	19	*8	*	32	—	*	—
94.39	51	27	24	*7	32	*7	*	12	32	*6	*
94.4	106	53	52	18	63	14	10	46	39	18	*
94.42	12	*6	*6	*6	*	*	*	*	*7	*	—
94.44	82	40	42	12	49	12	9	34	32	16	*
94.46	*5	*	*	—	*	*	*	*	—	—	*
94.6	349	246	103	*	250	81	16	137	100	83	28
94.61	13	10	*	*	11	*	*	*6	*6	*	*
94.62	125	100	26	*	70	44	11	47	40	27	12
94.63	43	30	13	—	28	13	*	13	13	10	*7
94.64	*6	*	*	*	*6	*	*	*	*	*	*
94.65	53	29	24	—	43	*9	*	30	*7	11	*
94.66	16	*8	*8	—	13	*	*	*6	*	*	*
94.67	12	*8	*	*	11	*	—	*	*8	*	*
94.68	48	33	15	*	42	*5	—	25	11	11	*
94.69	33	23	9	—	27	*5	*	*5	9	17	*
95	*8	*6	*	*	*	*	*	*	*	*	*
96	1,183	587	596	121	244	250	568	328	239	385	231
96.0	497	254	243	45	88	114	250	132	97	172	97
96.01	*5	*	*	*	*	*	*	*	*	*	*
96.04	363	190	172	39	53	82	188	99	76	122	66
96.05	11	*5	*6	*	*	*	*	*	*	*5	*
96.07	91	46	44	*	23	19	46	28	18	19	27
96.08	9	*	*6	*	*	*	*	*	*	*6	*
96.09	*7	*	*	*	*	*	*	*	*	*	*
96.1	*8	*	*6	*	*	*	*	*	*	*7	*
96.2	*6	*	*	*	*	*	*	*	*	*	*
96.3	31	14	17	*	17	*5	*6	*5	10	*9	*7
96.33	24	12	12	*	16	*	*	*	*9	*7	*5
96.38	*6	*	*	*	*	*	*	*	*	*	*
96.4	51	*6	45	*	44	*	*	*7	10	15	20
96.49	45	*	43	*	42	*	*	*	*8	12	20
96.5	42	26	16	*	17	*8	15	9	*8	15	10
96.56	15	10	*5	*	*	*	*8	*	*	*	*
96.59	22	13	*9	*	12	*	*	*	*	11	*
96.6	78	31	48	*7	*7	11	54	45	10	17	*6

* Figure does not meet standard of reliability or precision.
 — Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1996—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
96.7	469	250	219	62	66	107	233	127	102	150	90
96.70	15	*	10	*	*	*	*8	*	*	*8	*
96.71	304	162	142	37	45	70	152	85	70	93	57
96.72	150	83	67	23	21	33	73	41	31	49	29
97	74	43	31	9	17	18	31	21	13	24	15
97.0	22	13	9	*	*	*	14	*7	*	*7	*5
97.02	17	*9	*8	*	*	*	11	*	*	*5	*
97.3	*7	*	*	*	*	*	*	*	*	*	*
97.5	*7	*	*	*	*	*	*	*	*	*	*
97.6	12	*7	*	*	*	*	*	*	*	*6	*
97.62	9	*5	*	*	*	*	*	*	*	*	*
97.8	15	*8	*7	*	*	*	*	*	*	*	*
97.89	*7	*	*	*	*	*	*	*	*	*	*
98	28	18	11	*6	*9	*6	*8	*9	*6	*9	*
98.0	11	*6	*	*	*	*	*	*	*	*	*
98.02	9	*5	*	*	*	*	*	*	*	*	*
98.5	10	*6	*	*	*	*	*	*	*	*	*
98.51	10	*6	*	*	*	*	*	*	*	*	*
99	2,384	1,040	1,343	241	581	554	1,008	807	424	641	511
99.0	810	349	461	35	129	180	466	201	154	261	194
99.00	10	*5	*	*	*	*	*6	*	*	*	*
99.02	51	20	31	*	*7	16	27	11	12	11	17
99.03	24	13	11	*	*	*7	13	*	*	10	*9
99.04	632	262	370	23	96	128	385	160	118	213	141
99.05	44	21	23	*6	13	13	12	12	*8	12	13
99.07	41	21	20	*	*	13	21	10	10	11	10
99.1	317	120	197	28	114	69	106	138	50	70	59
99.11	60	*	60	*	60	—	—	11	11	17	21
99.14	11	*	*7	*	*	*	*	*5	*	*	*
99.15	85	38	46	17	20	18	30	31	22	19	13
99.17	41	18	24	*	10	14	15	22	*	10	*5
99.18	27	12	15	*	*9	*	10	21	*	*6	*
99.19	92	47	45	*	13	30	48	48	10	15	19
99.2	1,014	441	573	128	296	262	328	405	167	230	212
99.21	321	133	189	54	120	59	89	147	39	72	64
99.23	64	25	39	*8	15	13	28	30	11	14	*8
99.24	*5	*	*	*	*	*	*	*	*	*	—
99.25	290	134	156	32	51	107	100	109	51	81	49
99.29	327	145	182	34	105	80	109	115	64	58	89
99.4	10	*	10	*	10	—	—	*	*	*	*
99.47	*6	—	*6	—	*6	—	—	*	*	*	*
99.5	*8	*	*	*6	*	*	*	*	*	*	*
99.55	*7	*	*	*5	*	—	—	*	*	*	*
99.6	134	78	56	*	*8	32	92	33	36	40	24
99.60	43	23	19	*	*	*7	32	10	10	14	*9
99.61	23	14	*8	*	*	*7	14	*	*	*9	*
99.62	61	36	25	*	*	17	40	16	19	17	*8
99.7	10	*5	*5	*	*	*	*	*	*	*	*
99.71	*8	*	*	*	*	*	*	*	*	*	*
99.8	41	25	16	37	*	*	*	*6	*6	16	13
99.83	35	22	13	34	*	*	—	*	*5	13	12
99.9	36	17	19	*	14	*6	11	18	*	12	*
99.99	33	15	19	*	14	*6	10	18	*	11	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

Appendix

Technical Notes

Abstract Form

Figure I shows the medical abstract form used for the 1996 National Health Discharge Survey (NHDS). The hospital or facility number, medical record number, date of birth, and patient ZIP code are confidential information and are not released to the public.

Population Estimates

The population estimates used to compute rates, shown in [table I](#), are for the U.S. civilian population, including institutionalized persons, as of July 1, 1996. These estimates were provided by the U.S. Bureau of the Census and are consistent with population estimates published in *Current Population Reports*. Population estimates are shown for four geographic regions.

Region	States included
Northeast	Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, and Pennsylvania
Midwest	Michigan, Ohio, Illinois, Indiana, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas
South	Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, Florida, Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, and Texas
West	Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, Nevada, Washington, Oregon, California, Hawaii, and Alaska

Table I. Civilian population, by sex, age, and geographic region: United States, 1996

[Population estimates consistent with Series P-25, *Current Population Reports*, U.S. Bureau of the Census]

Sex, age, and geographic region	Population in thousands
Total	263,998
Sex	
Male	128,691
Female	135,306
Age	
Under 15 years	57,708
15–44 years	118,734
45–64 years	53,694
65 years and over	33,861
Geographic region	
Northeast	51,520
Midwest	61,983
South	92,373
West	58,122

Reliability of Estimates

The statistics presented in this report are based on sample surveys, and therefore may differ from the figures that would be obtained if a complete census had been taken. The standard error is primarily a measure of sampling variability that occurs by chance because only a sample rather than an entire universe is surveyed. Standard errors for the estimates from the NHDS were calculated with SUDAAN software, which takes into account the complex sample designs. A description of the software and the approach it uses has been published (14).

The chances are about 32 in 100 that an estimate from the samples would differ from a complete census by more than the standard error. The chances are 5 in 100 that the difference would be more than twice the standard error and about 1 in 100 that the difference would be more than 2.5 times as large as the standard error. The relative standard error (RSE) expresses the standard error as a percent of the estimate and can be multiplied by the estimate to obtain the standard error.

To derive approximate standard errors for the estimates in this report,

first calculate the approximate RSE of the estimate using the parameters shown in [table II](#). Then multiply the approximate RSE by the estimate. The approximate RSE of an estimate X may be estimated from the formula:

$$RSE(X) = \sqrt{a+b/X}$$

where *a* and *b* are defined in [table II](#). When multiplied by 100, RSE(X) is expressed as a percent of X.

The approximate RSE for a percent *p* in which the numerator is a subclass of the denominator may be calculated directly using the formula:

$$RSE(p) = \sqrt{b(1-p)/(pX)}$$

where 100*p* is the percent of interest, X is the denominator of the percent, and *b* is the parameter *b* in the formula for approximating the RSE (X). The values for *b* are given in [table II](#). When multiplied by 100, RSE(p) is expressed as a percent of p. This approximation is valid if the RSE of the denominator is less than 5 percent or the RSE's of the numerator and denominator are both less than 10 percent (15,16).

The approximate RSE of a population-based rate is equivalent to the approximate RSE of the numerator (obtained from [table II](#)), if the denominator is an estimate produced by the U.S. Bureau of the Census for the total population or one or more of the age-sex groups of the total population.

The approximate RSE of a ratio (X/Y) in which the numerator (X) and the denominator (Y) are both estimated from the same survey, but the numerator is not a subclass of the denominator, is calculated using the formula:

$$RSE(X/Y) = \sqrt{[RSE(X)]^2 + [RSE(Y)]^2}$$

When multiplied by 100, the RSE (X/Y) is expressed as a percent of the ratio X/Y. This approximation is valid if the RSE of the denominator is less than 5 percent or the RSE's of both the numerator and denominator are both less than 10 percent (15,16).

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
U.S. PUBLIC HEALTH SERVICE
CENTERS FOR DISEASE CONTROL
NATIONAL CENTER FOR HEALTH STATISTICS

A. PATIENT IDENTIFICATION

	Month	Day	Year
4. Date of admission			
5. Date of discharge			
6. Residence ZIP Code			

7. Date of birth

Month	

 -

Day	

 -

Year		

8. Age (Complete only if date of birth not given)

--	--	--

 Units

{	☐ Years
	☐ Months
	☐ Days

12. Marital status (Mark (X) one)

1 ☐ Married	3 ☐ Widowed	5 ☐ Separated
2 ☐ Single	4 ☐ Divorced	6 ☐ Not stated

2 ☐ Died
3 ☐ Status not stated

*U.S.GPO-1996-750-074/20352

Figure I. Medical abstract for the National Hospital Discharge Survey, 1996

Table II. Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics by selected characteristics: United States, 1996

Characteristic	Number of discharges or first-listed diagnosis		Number of all-listed diagnoses		Number of days of care		Number of procedures	
	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>
Total	0.00135	355.278	0.00315	340.624	0.00422	961.321	0.00265	367.004
Sex								
Male	0.00153	328.232	0.00412	323.367	0.00594	1,104.936	0.00347	330.761
Female	0.00157	384.999	0.00174	322.104	0.00226	1,341.036	0.00278	331.207
Age								
Under 15 years	0.01700	229.443	0.01958	289.646	0.02006	722.724	0.01858	236.345
15–44 years	0.00158	322.959	0.00265	328.234	0.00317	997.933	0.00398	298.470
45–64 years	0.00147	321.327	0.00235	302.405	0.00323	1,186.155	0.00373	293.436
65 years and over	0.00157	340.854	0.00196	309.446	0.00296	1,593.926	0.00252	277.559
Region								
Northeast	0.00437	213.840	0.01157	277.233	0.00778	632.459	0.00668	234.728
Midwest	0.01253	384.988	0.00819	183.086	0.00783	701.464	0.00821	154.838
South	0.00358	350.427	0.00388	306.111	0.00671	923.876	0.00441	270.511
West	0.00519	389.105	0.00822	420.446	0.01139	1,093.364	0.00829	369.526

NOTE: The relative standard error (RSE) can be determined from the equation $RSE(X)=100 \sqrt{a + b/x}$.

Changes in ICD–9–CM Codes

Although the *International Classification of Diseases, 9th Revision, Clinical Modification* has been used for coding NHDS data since 1979, the classification undergoes annual updating. Assignment of new diagnostic and procedure codes, fourth and fifth digit expansion of codes, as well as code deletions, are contained in addenda developed by the ICD–9–CM Coordination and Maintenance Committee and approved by the Director of the NCHS and the Administrator of the Health Care Financing Administration. Addenda to the ICD–9–CM become effective October 1 of the affected year and have been released annually since 1986.

The data in this report were coded using the addenda for October 1, 1986–95. When data were reported using code changes introduced in the October 1, 1996, addenda, the new codes were converted to their previous code assignments so that the annual estimates would be based on consistent coding.

The [tables III](#) and [IV](#) show the new ICD–9–CM codes from the 1986–95 addenda, the date they were introduced, and the code to which the diagnosis or

procedure had been assigned prior to the development of the new code. Specific titles and more detailed information about the coding scheme can be found in appropriate volumes of the ICD–9–CM (13).

Table III. Changes in diagnosis codes

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
005.81	1995	005.8	204.11	1991	V10.61
005.89	1995	005.8	204.20	1991	204.2
008.00–008.09	1992	008.0	204.21	1991	V10.61
008.43–008.47	1992	008.49	204.80	1991	204.8
008.61–008.69	1992	008.6	204.81	1991	V10.61
041.00–041.09	1992	041.0	204.90	1991	204.9
041.10–041.19	1992	041.1	204.91	1991	V10.61
041.81–041.89	1992	041.8	205.00	1991	205.0
041.86	1995	041.84	205.01	1991	V10.62
042	1994	042.0–042.2, 042.9, 043.0–043.3, 043.9, 044.0, 044.9	205.10	1991	205.1
042.0–042.9	1986	279.19	205.11	1991	V10.62
043.0–043.9	1986	279.19	205.20	1991	205.2
044.0–044.9	1986	279.19	205.21	1991	V10.62
070.20–070.21	1991	070.2	205.30	1991	205.3
070.22	1994	070.20	205.31	1991	V10.62
070.23	1994	070.21	205.80	1991	205.8
070.30–070.31	1991	070.3	205.81	1991	V10.62
070.32	1994	070.30	205.90	1991	205.9
070.33	1994	070.31	205.91	1991	V10.62
070.41–070.49	1991	070.4	206.00	1991	206.0
070.44	1994	070.41	206.01	1991	V10.63
070.51–070.59	1991	070.5	206.10	1991	206.1
070.54	1994	070.51	206.11	1991	V10.63
077.98–077.99	1993	077.9	206.20	1991	206.2
078.10–078.11, 078.19	1993	078.1	206.21	1991	V10.63
078.88	1993	078.89	206.80	1991	206.8
079.4	1993	079.8	206.81	1991	V10.63
079.50–079.53, 079.59	1993	079.8	206.90	1991	206.9
079.81	1995	079.89	206.91	1991	V10.63
079.88–079.89	1993	079.8	207.00	1991	207.0
079.98–079.99	1993	079.9	207.01	1991	V10.69
088.81, 088.89	1989	088.8	207.10	1991	207.1
088.82	1993	088.89	207.11	1991	V10.69
099.40–099.49	1992	099.4	207.20	1991	207.2
099.50–099.59	1992	078.89	207.21	1991	V10.69
112.84–112.85	1992	112.89	207.80	1991	207.8
114.4–114.5	1993	114.3	207.81	1991	V10.69
176.0–176.9	1991	173.0–173.9	208.00	1991	208.0
203.00	1991	203.0	208.01	1991	V10.60
203.01	1991	V10.79	208.10	1991	208.1
203.10	1991	203.1	208.11	1991	V10.60
203.11	1991	V10.79	208.20	1991	208.2
203.80	1991	203.8	208.21	1991	V10.60
203.81	1991	V10.79	208.80	1991	208.8
204.00	1991	204.0	208.81	1991	V10.60
204.01	1991	V10.61	208.90	1991	208.9
204.10	1991	204.1	208.91	1991	V10.60
			237.70–237.72	1990	237.7
			250.02	1993	250.90
			250.03	1993	250.91
			250.12	1993	250.10
			250.13	1993	250.11
			250.22	1993	250.20
			250.23	1993	250.21
			250.32	1993	250.30
			250.33	1993	250.31
			250.42	1993	250.40
			250.43	1993	250.41
			250.52	1993	250.50
			250.53	1993	250.51
			250.62	1993	250.60
			250.63	1993	250.61
			250.72	1993	250.70

Table III. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
250.73	1993	250.71	410.40–410.42	1989	410.4
250.82	1993	250.80	410.50–410.52	1989	410.5
250.83	1993	250.81	410.60–410.62	1989	410.6
250.92	1993	250.90	410.70–410.72	1989	410.7
250.93	1993	250.91	410.80–410.82	1989	410.8
			410.90–410.92	1989	410.9
278.00–278.01	1995	278.0	411.81	1989	410.9
			411.89	1989	411.8
283.10–283.11,283.19	1993	283.1	414.00–414.01	1994	414.0
305.1	1994	305.10,305.11,305.12, 305.13 (delete code)	414.02–414.03	1994	996.03
			415.11	1995	997.3 & 415.1
312.81–312.82,381.89	1994	312.8	415.19	1995	415.1
320.81–320.89	1992	320.8	429.71	1989	410.0–410.9
			429.79	1989	410.0–410.9
333.92–333.93	1994	333.99	433.00–433.01	1993	433.0
			433.10–433.11	1993	433.1
337.20–337.22,337.29	1993	337.9	433.20–433.21	1993	433.2
			433.30–433.31	1993	433.3
342.00–342.02	1994	342.0	433.80–433.81	1993	433.8
342.10–342.12	1994	342.1	433.90–433.91	1993	433.9
342.80–342.82	1994	342.9			
342.90–342.92	1994	342.9	434.00–434.01	1993	434.0
			434.10–434.11	1993	434.1
344.00–344.04,344.09	1994	344.0	434.90–434.91	1993	434.9
344.30–344.32	1994	344.3			
344.40–344.42	1994	344.4	435.3	1995	435.0 & 435.1
344.81,344.89	1993	344.8	437.7	1992	780.9
			440.20–440.22	1992	440.2
345.00–345.01	1989	345.0			
345.10–345.11	1989	345.1	440.23	1993	440.20 & 707.1 or 707.8 or 707.9
345.40–345.41	1989	345.4	440.24	1993	440.20 & 785.4
345.50–345.51	1989	345.5	440.29	1993	440.20
345.60–345.61	1989	345.6			
345.70–345.71	1989	345.7	440.30–440.32	1994	996.1
345.80–345.81	1989	345.8			
345.90–345.91	1989	345.9	441.00–441.03	1994	441.0
			441.6	1993	441.1 & 441.3
346.00–346.01	1992	346.0	441.7	1993	441.2 & 441.4
346.10–346.11	1992	346.1			
346.20–346.21	1992	346.2	446.20–446.21,446.29	1990	446.2
346.80–346.81	1992	346.8			
346.90–346.91	1992	346.9	451.82–451.84	1993	451.89
			458.2	1995	997.9 & 458.9
355.71	1993	354.4	482.30–482.39	1992	482.3
355.79	1993	355.7	482.81–482.89	1992	482.8
371.82	1992	371.89	483.0	1992	483
			483.8	1992	483
374.87	1990	374.89	491.20–491.21	1991	491.2
403.00–403.01	1989	403.0	493.20	1989	493.90
403.10–403.11	1989	403.1	493.21	1989	493.91
403.90–403.91	1989	403.9			
			512.1	1994	997.3
404.00–404.03	1989	404.0			
404.10–404.13	1989	404.1	518.81	1987	799.1
404.90–404.93	1989	404.9	518.82–518.89	1987	518.8
410.00–410.02	1989	410.0	524.00–524.09	1992	524.0
410.10–410.12	1989	410.1	524.10–524.19	1992	524.1
410.20–410.22	1989	410.2	524.60–524.69	1991	524.6
410.30–410.32	1989	410.3	524.70–524.79	1992	524.8

Table III. Changes in diagnosis codes—Con

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
530.10–530.11, 530.19	1993	530.1	702.11,702.19	1994	702.1
530.81	1993	530.1	704.02	1993	704.09
530.82–530.84, 530.89	1993	530.8	709.00–709.01,709.09	1994	709.0
535.00–535.01	1991	535.0	710.5	1992	288.3,729.1
535.10–535.11	1991	535.1	728.86	1995	729.4
535.20–535.21	1991	535.2	733.10–733.16, 733.19	1993	733.1
535.30–535.31	1991	535.3	738.10–738.19	1992	738.1
535.40–535.41	1991	535.4	747.60–747.64, 747.69	1993	747.6
535.50–535.51	1991	535.5	747.82	1993	747.89
535.60–535.61	1991	535.6	753.10–753.17,753.19	1990	753.1
536.3	1994	536.8	759.81–759.89	1989	759.8
537.82	1990	537.89	759.83	1994	759.89
537.83	1991	537.82	760.75	1991	760.79
556.0–556.6	1994	556	760.76	1994	760.79
556.8–556.9	1994	556	764.00–764.09	1988	764.0
562.02	1991	562.00	764.10–764.19	1988	764.1
562.03	1991	562.01	764.20–764.29	1988	764.2
562.12	1991	562.10	764.90–764.99	1988	764.9
562.13	1991	562.11	765.00–765.09	1988	765.0
569.60–569.61	1995	569.6	765.10–765.19	1988	765.1
569.69	1995	569.6	780.01–780.09	1992	780.0
569.84	1990	557.1	780.03	1993	780.01
569.85	1991	569.84	780.57	1992	780.51,780.53
593.70–593.73	1994	593.7	781.8	1994	781.9
596.51–596.53	1992	596.5	787.01–787.03	1994	787.0
596.54	1992	344.61	787.91	1995	558.9
596.55–596.59	1992	596.5	787.99	1995	787.9
599.81–599.89	1992	599.8	788.20–788.21, 788.29	1993	788.2
645.0	1991	645	788.30–788.39	1992	788.3
651.30–651.31,651.33	1989	651.00–651.01,651.03	788.41–788.43	1993	788.4
651.40–651.41,651.43	1989	651.10–651.11,651.13	788.61–788.62, 788.69	1993	788.6
651.50–651.51,651.53	1989	651.20–651.21,651.23	789.00–789.07, 789.09	1994	789.0
651.60–651.61,651.63	1989	651.80–651.81,651.83	789.30–789.37, 789.39	1994	789.3
654.20–654.21,654.23	1990	654.2,654.9	789.40–789.47, 789.49	1994	789.4
654.90–651.94	1990	654.2,654.9	789.60–789.67, 789.69	1994	789.6
657.0	1991	657	790.91	1993	790.9
659.60,659.61,659.63	1992	659.80–659.81,659.83	790.92	1993	286.9
665.10,665.11	1992	665.10,665.11,665.12,665.14 Note: The title for the subcategory 665.1 has been changed, making the fifth-digit subclassification, 665.12 and 665.14 invalid.	790.93, 790.99	1993	790.9
670.0	1991	670	795.71	1994	795.8 (delete code)
672.0	1991	672	795.79	1994	795.7
677	1994	There was no previous code assignment for this code.	795.8	1986	795.7
690.10	1995	690	864.05	1992	864.09
690.11	1995	691.8 & 704.8	864.15	1992	864.19
690.12	1995	691.8	909.5	1994	909.9
690.18	1995	690	925.1–925.2	1993	925
690.8	1995	690	989.81–989.84	1995	989.8
692.72–692.74	1992	692.79	989.89	1995	989.8
692.82–692.83	1992	692.89	995.60–995.69	1993	995.0
702.0–702.8	1991	702	996.04	1994	996.09

Table III. Changes in diagnosis codes—Con

Current code(s) assignment	Effective October 1	Previous code(s) assignment
996.51–996.59	1987	996.5
996.60–996.69	1989	996.6
996.70–996.79	1989	996.7
996.80–996.89	1987	996.8
996.85	1990	999.8
997.00–997.01	1995	997.0
997.02	1995	997.9 & 430–434, 436
997.09	1995	997.0
997.91	1995	997.9
997.99	1995	997.9
998.81–998.82, 998.89	1994	998.8
V03.81–V03.82, V03.89	1994	V03.8
V05.3–V05.4	1993	V05.8
V06.5–V06.6	1994	V06.8
V07.31, V07.39	1994	V07.3
V07.4	1992	V07.8
V08	1994	044.9, 795.8 (delete code)
V09.0–V09.91	1993	There were no previous code assignments for these codes.
V12.00–V12.03, V12.09	1994	V12.0
V12.50–V12.52	1995	V12.5
V12.59	1995	V12.5
V12.70–V12.72, V12.79	1994	V12.7
V13.00–V13.01, V13.09	1994	V13.0
V15.82	1994	305.13 (delete code)
V15.84–V15.86	1995	V15.89
V25.43	1992	V25.49
V25.5	1992	V25.8
V29.0–V29.8	1992	V71.8
V29.9	1992	V71.9
V30.00–V30.01	1989	V30.0
V31.00–V31.01	1989	V31.0
V32.00–V32.01	1989	V32.0
V33.00–V33.01	1989	V33.0
V34.00–V34.01	1989	V34.0
V35.00–V35.01	1989	V35.0
V36.00–V36.01	1989	V36.0
V37.00–V37.01	1989	V37.0
V39.00–V39.01	1989	V39.0
V43.60–V43.66, V43.69	1994	V43.6
V43.81–V43.82	1995	V43.8
V43.89	1995	V43.8
V45.00	1994	V45.89

Table III. Changes in diagnosis codes—Con

Current code(s) assignment	Effective October 1	Previous code(s) assignment
V45.01	1994	V45.0
V45.02, V45.09	1994	V45.89
V45.51	1994	V45.5
V45.52, V45.59	1994	V45.89
V45.82	1994	V45.89
V45.83	1995	V45.89
V49.60–V49.67	1994	V49.5
V49.70–V49.77	1994	V49.5
V50.41–V50.42, V50.49	1994	V50.8
V53.31	1994	V53.3
V53.32, V53.39	1994	V53.9
V56.1	1995	V58.89
V57.21–V57.22	1994	V57.2
V58.41, V58.49	1994	V58.4
V58.61	1995	V67.51
V58.69	1995	V67.51
V58.81, V58.89	1994	V58.8
V58.82	1995	V58.89
V59.01–V59.02	1995	V59.0
V59.09	1995	V59.0
V59.6	1995	V59.8
V65.40–V65.45, V65.49	1994	V65.4
V69.0–V69.3	1994	No previous code assignments for these codes.
V69.8–V69.9	1994	
V72.81–V72.85	1993	V72.8
V73.88–V73.89	1993	V73.8
V73.98–V73.99	1993	V73.9
E854.8	1995	E858.8
E869.4	1994	E869.8
E880.1	1995	E884.9
E884.3–E884.4	1995	E884.2
E884.5–E884.6	1995	E884.9
E906.5	1995	E906.3
E908.0–E908.4	1995	E908
E908.8–E908.9	1995	E908
E909.0–E909.4	1995	E909
E909.8–E909.9	1995	E909
E920.5	1995	E920.4
E924.2	1995	E924.0
E968.5	1995	E968.8

Table IV. Changes in procedure codes

Current code(s) assignment	Effective October 1	Previous code(s) assignment
02.96	1992	89.19
03.90	1987	03.99 (Insertion of catheter)
05.25	1995	39.7 (code deleted)
11.75	1989	11.79
11.76	1989	11.62
20.96–20.98	1986	20.95
22.12	1988	22.11
26.12	1988	26.11
29.31	1991	83.02
29.32	1991	29.3
29.33	1991	29.3
29.39	1991	29.3
31.45	1988	31.43–31.44
31.95	1989	31.75
32.01	1989	32.0
32.09	1989	32.0
32.22	1995	32.29, 32.9
32.28	1989	32.29
33.27	1987	33.22 + 33.27
33.28	1987	33.27
33.29	1987	33.28–33.29
33.50	1995	33.5
33.51	1995	33.5
33.52	1995	33.5
33.6	1990	33.5 + 37.5
34.05	1994	34.99
35.84	1988	35.82
35.96	1986	35.03
36.00–36.03	1986	36.0
36.04	1986	39.97
36.05	1987	36.01
36.05	1986	36.01 ¹ , 36.02
36.06	1995	36.01, 36.02, 36.03, 36.05
36.09	1986	36.0
36.09	1991	36.00 (code deleted)
37.26–37.27	1988	37.29
37.34	1988	37.33
37.65	1995	37.62
37.66	1995	37.62
37.70 (Leads only)	1987	(Leads/Device) 37.70
37.71–37.72 (Leads only)	1987	(Leads/Device) 37.74
37.73 (Leads only)	1987	(Leads/Device) 37.73
37.74 (Leads only)	1987	(Leads/Device) 37.76
37.75 (Leads only)	1987	(Leads/Device) 37.89
37.76 (Leads only)	1987	(Leads/Device) 37.81
37.77 (Leads only)	1987	(Leads/Device) 37.83–37.84
37.78	1987	37.71–37.72
37.79	1987	86.09
37.80–37.87	1992	89.49 ²

Table IV. Changes in procedure codes

Current code(s) assignment	Effective October 1	Previous code(s) assignment
37.80 (Device only)	1987	(Leads/Device) 37.73–37.77
37.81 (Device only)	1987	(Leads/Device) 37.73–37.77
37.82 (Device only)	1987	(Leads/Device) 37.73–37.77
37.83 (Device only)	1987	(Leads/Device) 37.73–37.77
37.85–37.87	1987	37.85
37.89	1987	37.86 + 37.89
37.94–37.98	1986	37.99
38.22	1986	38.29
38.44 (Abdominal aorta only)	1986	38.44 (Entire aorta)
38.45 (Thoracic aorta added)	1986	38.44–38.45
38.95	1989	38.93
39.28	1991	39.29
39.50	1995	39.59
39.65	1988	39.61
39.66	1990	39.65
41.00–41.03	1988	41.0
41.04	1994	99.79
42.25	1988	42.24
42.33	1989	42.32, 42.39
42.33	1990	42.91
43.11	1989	43.1
43.19	1989	43.1, 43.2
43.41	1989	43.41, 43.49
44.21	1986	44.2
44.22	1986	44.99
44.29	1986	44.2
44.43	1989	43.49, 45.32
44.44	1989	38.86
44.49	1989	43.0
44.93–44.94	1986	44.99
45.16	1988	45.14 (45.15 before 1987)
45.30	1989	45.31, 45.32
45.42	1988	45.41
45.43	1989	45.49
45.75 (Hartmann resection added)	1988	48.66 (code deleted)
45.95	1987	45.93
46.13	1992	46.12 (code deleted)
46.32	1989	46.39
46.85	1989	46.99
48.36	1995	45.42
49.31	1989	49.3
49.39	1989	49.3
51.10	1989	51.97
51.11	1989	51.11, 51.97
51.14	1989	51.12
51.15	1989	51.97
51.22	1991	51.21 (code deleted), 51.22
51.23	1991	51.22
51.64	1989	51.69
51.84–51.88	1989	51.97
51.97	1986	52.91, 51.99, or 51.82
51.98	1986	51.99

Table IV. Changes in procedure codes—Con

Current code(s) assignment	Effective October 1	Previous code(s) assignment
52.13	1989	51.97,52.91
52.14	1989	52.11
52.21	1989	52.2
52.22	1989	52.2
52.93	1989	52.93 + 52.91
52.94	1989	52.09
52.97	1989	52.91
52.98	1989	52.91
52.99	1989	52.93,52.94,52.99
54.24	1987	54.23
54.25	1993	54.98
55.03–55.04	1986	55.02
56.33–56.34	1987	56.33
56.35	1987	45.12
57.17–57.18	1989	57.21
57.22	1989	57.22,57.82
58.31	1990	58.3
58.39	1990	58.3
58.93	1986	57.99
59.72	1995	59.79
59.96	1986	59.95
60.21	1995	60.2
60.29	1995	60.2
60.95	1991	60.99
64.97	1986	64.95
66.01	1992	66.0
66.02	1992	66.73
68.15	1987	68.14
68.16	1987	68.13
68.9	1992	68.4
74.3	1992	69.11 (code deleted)
77.56	1989	77.89,78.49,81.18
77.57	1989	77.89,80.48,81.18,83.85
77.58	1989	77.59,81.18
78.10	1991	78.40
78.11	1991	78.41
78.12	1991	78.42
78.13	1991	78.43
78.14	1991	78.44
78.15	1991	78.45
78.16	1991	78.46
78.17	1991	78.47
78.18	1991	78.48
78.19	1991	78.49
78.20	1991	78.10,78.20,78.30
78.21	1991	78.11,78.31
78.22	1991	78.12,78.22,78.32
78.23	1991	78.13,78.23,78.33
78.24	1991	78.14,78.34
78.25	1991	78.15,78.25,78.35
78.27	1991	78.17,78.27,78.37
78.28	1991	78.18,78.38

Table IV. Changes in procedure codes—Con

Current code(s) assignment	Effective October 1	Previous code(s) assignment
78.29	1991	78.11,78.16,78.19,78.29,78.39
78.39	1991	78.31
78.90 ³	1987	78.40
78.91 ³	1987	78.41
78.92 ³	1987	78.42
78.93 ³	1987	78.43
78.94 ³	1987	78.44
78.95 ³	1987	78.45
78.96 ³	1987	78.46
78.97 ³	1987	78.47
78.98 ³	1987	78.48
78.99 ³	1987	78.49
80.50–80.59	1986	80.5
81.03	1989	81.02
81.04–81.05	1989	81.03,81.04,81.05
81.06–81.07	1989	81.06,81.07
81.08	1989	81.06,81.07,81.08
81.09	1989	81.08
81.40	1989	81.69
81.51	1989	81.51,81.59
81.52	1989	81.61,81.62,81.63,81.64
81.53	1989	81.51,81.59,81.61,81.62, 81.63,81.64
81.54–81.55	1989	81.41
81.56	1989	81.48
81.57	1989	81.31,81.39
81.59	1989	81.39
81.72	1989	81.79
81.73–81.74	1989	81.86
81.75	1989	81.87
81.79	1989	81.79,81.87
81.80	1989	81.81
81.97	1992	81.59
85.95	1987	85.99
85.96	1987	85.99
86.06	1987	86.09
86.07	1990	86.09
86.27	1986	86.22–86.23
86.28	1988	86.22
86.93	1987	86.89
88.90	1986	88.39
88.91	1986	89.15
88.92	1986	89.39
88.93	1986	89.15
88.94	1986	89.39
88.95	1986	89.29
88.97	1989	88.99
88.98	1989	88.90
88.99	1986	89.39
89.10	1989	89.15
89.17–89.18	1988	89.15
89.19	1989	89.15
89.50	1991	89.54
92.3	1995	01.59, 04.07, 07.63, 07.68
93.90	1988	93.92
94.61–94.69	1989	94.25

Table IV. Changes in procedure codes—Con

Current code(s) assignment	Effective October 1	Previous code(s) assignment
96.6	1986	96.35
96.70	1991	93.92 (code deleted)
96.71	1991	93.92 (code deleted)
96.72	1991	93.92 (code deleted)
97.05	1989	51.97
98.51–98.52	1989	59.96 (code deleted)
98.59	1989	59.96 (code deleted)
99.00	1995	99.02
99.15	1986	99.29
99.28	1994	99.25

Table IV. Changes in procedure codes—Con

Current code(s) assignment	Effective October 1	Previous code(s) assignment
99.71–99.79 ⁴	1988	99.07
99.85	1987	93.35
99.86	1987	93.39
99.88	1988	99.83

¹Before October 1986 contents of current code 36.05 would have been assigned to 36.0.

²Code 89.49 deleted; this procedure is included in the code for pacemaker insertion/replacement.

³Codes 78.90–78.99 were retitled as "Insertion of bone growth stimulator" in October 1987; the previous contents of codes 78.90–78.99 were reassigned to codes 78.40–78.49.

⁴Codes 99.71–99.79 were deleted in October 1987; their contents were not transferred elsewhere. In the October 1988 revision, codes 99.71–99.79 were reclassified as "Therapeutic apheresis."

Vital and Health Statistics series descriptions

- SERIES 1. **Programs and Collection Procedures**—These reports describe the data collection programs of the National Center for Health Statistics. They include descriptions of the methods used to collect and process the data, definitions, and other material necessary for understanding the data.
- SERIES 2. **Data Evaluation and Methods Research**—These reports are studies of new statistical methods and include analytical techniques, objective evaluations of reliability of collected data, and contributions to statistical theory. These studies also include experimental tests of new survey methods and comparisons of U.S. methodology with those of other countries.
- SERIES 3. **Analytical and Epidemiological Studies**—These reports present analytical or interpretive studies based on vital and health statistics. These reports carry the analyses further than the expository types of reports in the other series.
- SERIES 4. **Documents and Committee Reports**—These are final reports of major committees concerned with vital and health statistics and documents such as recommended model vital registration laws and revised birth and death certificates.
- SERIES 5. **International Vital and Health Statistics Reports**—These reports are analytical or descriptive reports that compare U.S. vital and health statistics with those of other countries or present other international data of relevance to the health statistics system of the United States.
- SERIES 6. **Cognition and Survey Measurement**—These reports are from the National Laboratory for Collaborative Research in Cognition and Survey Measurement. They use methods of cognitive science to design, evaluate, and test survey instruments.
- SERIES 10. **Data From the National Health Interview Survey**—These reports contain statistics on illness; unintentional injuries; disability; use of hospital, medical, and other health services; and a wide range of special current health topics covering many aspects of health behaviors, health status, and health care utilization. They are based on data collected in a continuing national household interview survey.
- SERIES 11. **Data From the National Health Examination Survey, the National Health and Nutrition Examination Surveys, and the Hispanic Health and Nutrition Examination Survey**—Data from direct examination, testing, and measurement on representative samples of the civilian noninstitutionalized population provide the basis for (1) medically defined total prevalence of specific diseases or conditions in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics, and (2) analyses of trends and relationships among various measurements and between survey periods.
- SERIES 12. **Data From the Institutionalized Population Surveys**—Discontinued in 1975. Reports from these surveys are included in Series 13.
- SERIES 13. **Data From the National Health Care Survey**—These reports contain statistics on health resources and the public's use of health care resources including ambulatory, hospital, and long-term care services based on data collected directly from health care providers and provider records.
- SERIES 14. **Data on Health Resources: Manpower and Facilities**—Discontinued in 1990. Reports on the numbers, geographic distribution, and characteristics of health resources are now included in Series 13.
- SERIES 15. **Data From Special Surveys**—These reports contain statistics on health and health-related topics collected in special surveys that are not part of the continuing data systems of the National Center for Health Statistics.
- SERIES 16. **Compilations of Advance Data From Vital and Health Statistics**—Advance Data Reports provide early release of information from the National Center for Health Statistics' health and demographic surveys. They are compiled in the order in which they are published. Some of these releases may be followed by detailed reports in Series 10–13.
- SERIES 20. **Data on Mortality**—These reports contain statistics on mortality that are not included in regular, annual, or monthly reports. Special analyses by cause of death, age, other demographic variables, and geographic and trend analyses are included.
- SERIES 21. **Data on Natality, Marriage, and Divorce**—These reports contain statistics on natality, marriage, and divorce that are not included in regular, annual, or monthly reports. Special analyses by health and demographic variables and geographic and trend analyses are included.
- SERIES 22. **Data From the National Mortality and Natality Surveys**—Discontinued in 1975. Reports from these sample surveys, based on vital records, are now published in Series 20 or 21.
- SERIES 23. **Data From the National Survey of Family Growth**—These reports contain statistics on factors that affect birth rates, including contraception, infertility, cohabitation, marriage, divorce, and remarriage; adoption; use of medical care for family planning and infertility; and related maternal and infant health topics. These statistics are based on national surveys of women of childbearing age.
- SERIES 24. **Compilations of Data on Natality, Mortality, Marriage, Divorce, and Induced Terminations of Pregnancy**—These include advance reports of births, deaths, marriages, and divorces based on final data from the National Vital Statistics System that were published as supplements to the *Monthly Vital Statistics Report (MVSR)*. These reports provide highlights and summaries of detailed data subsequently published in *Vital Statistics of the United States*. Other supplements to the MVSR published here provide selected findings based on final data from the National Vital Statistics System and may be followed by detailed reports in Series 20 or 21.

For answers to questions about this report or for a list of reports published in these series, contact:

Data Dissemination Branch
National Center for Health Statistics
Centers for Disease Control and Prevention
6525 Belcrest Road, Room 1064
Hyattsville, MD 20782-2003
(301) 436-8500
E-mail: nchsquery@cdc.gov
Internet: www.cdc.gov/nchswwww/

**DEPARTMENT OF
HEALTH & HUMAN SERVICES**

Centers for Disease Control and Prevention
National Center for Health Statistics
6525 Belcrest Road
Hyattsville, Maryland 20782-2003

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

STANDARD MAIL (B)
POSTAGE & FEES PAID
CDC/NCHS
PERMIT NO. G-284